Replacing Random Signs by Neighborhood Counts

A deterministic early-enumeration theorem and a limitation

8 October 2026

Abstract

The cancellation needed for early enumeration has an exact formulation in terms of agreement and disagreement counts between signed neighborhoods. For a fixed family of learned inequalities, these pair counts determine the variance of their sum under a uniform completion; independence between neighborhoods is unnecessary. We formulate a general deterministic theorem using signed co-occurrence counts and counts of residual-gap profiles. A bipartite special case illustrates the mechanism, but does not improve on the simpler algorithm available for bipartite formulas. We also show why pairwise balance alone is insufficient for this particular summed-constraint generator: with zero initial gaps, it can enumerate a constant fraction of all assignments despite exactly orthogonal signed neighborhoods. These are sufficient conditions and a limitation of one algorithm, not a characterization of hard Max 2-SAT.

1 What independence means here

Independence in the random-sign model is a property of the joint distribution over formulas. Empirical counts inside one formula are different objects. Even exactly balanced one- and two-coordinate counts do not imply joint independence. For example, the three sign patterns (u, v, uv) over the four choices $(u, v) \in \{-1, +1\}^2$ have uniform one- and two-coordinate frequencies but only four of eight possible triples.

We should replace independence by the particular counting inequalities used by the algorithm. This note separates two jobs: making learned inequalities sufficiently restrictive, and preventing their bad values from aligning too strongly. Pair counts handle the second job directly. In the theorem below, literal-degree gaps supply the first job.

2 Exact agreement counts for a fixed constraint family

Consider fixed inequalities $L_i(y) \le k_i$ on r free variables, each counting one bad value on its support N_i. Write assignments as $y_w \in \{-1, +1\}$ and set

$$b_i(w) = \begin{cases} +1, & \text{if value } +1 \text{ is bad for } i \text{ at } w, \\ -1, & \text{if value } -1 \text{ is bad for } i \text{ at } w, \\ 0, & w \notin N_i. \end{cases}$$

Thus $L_i(y) = |N_i|/2 + \frac{1}{2}\sum_w b_i(w) y_w$. For two rows define the directly countable quantity

$$C_{ij} = A_{ij} - O_{ij} = \sum_w b_i(w) b_j(w), \tag{1}$$

where A_{ij} counts shared variables with agreeing bad values and O_{ij} counts shared variables with opposite bad values. In particular, $C_{ii} = |N_i|$.

For a selected family I, define

$$s_I = \sum_{i \in I} \left(\frac{|N_i|}{2} - k_i \right), \tag{2}$$

$$V_I = \sum_w \left(\sum_{i \in I} b_i(w) \right)^2 = \sum_{i \in I} |N_i| + 2 \sum_{\substack{i<j \\ i,j \in I}} C_{ij}. \tag{3}$$

The sum of the inequalities is $Q_I(y) \le -s_I$, where $Q_I(y) = \frac{1}{2} \sum_w (\sum_{i \in I} b_i(w)) y_w$.

Lemma 1 (A deterministic counting bound)**.** *If $s_I > 0$ and $V_I > 0$, the number of assignments satisfying the summed inequality is at most*

$$2^r \exp\left(-\frac{2 s_I^2}{V_I} \right). \tag{4}$$

If $s_I > 0$ and $V_I = 0$, that number is zero.

Proof. Use a uniform assignment only to count completions of this fixed family. Its coordinates y_w are independent signs, regardless of any dependencies between the rows b_i. Thus

$$\mathbb{E} e^{-\lambda Q_I} = \prod_w \cosh\left(\frac{\lambda}{2} \sum_{i \in I} b_i(w) \right) \le e^{\lambda^2 V_I / 8}.$$

Markov's inequality with $\lambda = 4 s_I / V_I$ gives (4). If $V_I = 0$, then Q_I is identically zero. □

This is stronger than an analogy with independence: the exact variance is $V_I/4$, and equation (3) expresses it using pair counts. Independence of the neighborhoods is not used. For example, when every $C_{ij} = 0$ for $i \ne j$, then $V_I = \sum_{i \in I} |N_i|$ for every selected subfamily, even if there are higher-order relations among rows.

For exact counts, let m_w^+ and m_w^- count bad occurrences at w, put $c_w = |m_w^+ - m_w^-|$, and $B = \sum_i k_i - \sum_w \min(m_w^+, m_w^-)$. Orient the cheaper value as zero. The summed inequality is $\sum_w c_w z_w \le B$, and its exact completion count is

$$\sum_{j=0}^{\lfloor B \rfloor} [x^j] \prod_w (1 + x^{c_w}),$$

interpreted as zero when $B < 0$. Thus even the full count is determined by a histogram of signed occurrence multiplicities, not by a claim of independence. The pair counts give the simpler sufficient estimate above.

3 A deterministic counting version of the full proof

Fix any simple signed interaction graph, with one clause per edge and no initial units, and fix an order and a depth $T = \lfloor \tau n \rfloor$, where $0 < \tau < 1$ is constant. Let $d \ge 1$ be a reference degree scale, h_i the initial literal-degree gap, and J_{iw} the product of the literal signs on an edge, zero off edges. Define the ordered interaction row $r_i(w) = J_{iw}$ for $w > i$, and zero otherwise, for $i \le T$. Its raw pair counts are

$$M_{ij} = \sum_{w > \max(i,j)} J_{iw} J_{jw} = \#\{\text{agreeing interaction signs}\} - \#\{\text{opposite interaction signs}\}.$$

Here M_{ii} is the number of later neighbors of i. Unlike C in Section 2, these rows have not been oriented by preferred values.

For a prefix $u \in \{-1,+1\}^T$, define its total gap

$$G(u) = \sum_{i \le T} \left| h_i - \sum_{j<i} J_{ij} u_j \right|.$$

This is defined on the full assignment cube without survival tests or optimization. Fix constants $K > 0$, $H \ge 0$, $g > 0$, and $\eta > 0$. The three conditions are

$$\sum_{i,j \le T} M_{ij} z_i z_j \le Kd \sum_{i \le T} z_i^2 \qquad \text{for all } z \in \mathbb{R}^T, \tag{5}$$

$$\sum_{i \le T} h_i^2 \le Hnd, \tag{6}$$

$$\#\{u \in \{-1,+1\}^T : G(u) < gn\sqrt{d}\} \le 2^T e^{-\eta n}. \tag{7}$$

The first is an aggregate condition on pair counts, not a bound on each pair separately. It can be checked by linear algebra on M; the stronger condition $\sum_j |M_{ij}| \le Kd$ for every i also suffices. The last condition counts exceptional gap profiles. It is not asserted to have a polynomial-time membership test. Replacing it by simpler input conditions is a remaining structural question.

Theorem 2 (Full deterministic counting criterion)**.** *Every fixed formula satisfying* (5)–(7) *is solved exactly by fixed-order branching through depth T, followed by enumeration of the summed learned inequality, in polynomial space and time*

$$O^*(2^T + 2^n e^{-cn}), \qquad c = \min\left\{\eta, \frac{g^2}{32(H+K)}\right\} > 0.$$

All constants are independent of d. The statement has no probability qualification over input formulas.

Proof. Use a uniform full assignment u only to count generated candidates. Let I consist of its nonpreferred decisions among the first T, $v_i = u_i[i \in I]$, and $\Delta_i = h_i - \sum_{j<i} J_{ij} u_j$. As in the early-enumeration note, set

$$S = \sum_{i \in I} (|\Delta_i|/2 + 1), \qquad Q = \frac{1}{2} \sum_{w=1}^{n} u_w \sum_{\substack{i<w \\ i \le T}} J_{iw} v_i.$$

Exactly those assignments with $S + Q \le 0$ pass their prefix's summed inequality. Consequently, their total number is $W(F) = 2^n \mathbb{P}_u[S + Q \le 0]$ for this fixed formula.

The coefficient of u_w in the inner sum for Q is determined before u_w is exposed. Its total squared coefficient is at most $Kd \sum_i v_i^2 \le KdT$ by (5). The conditional cosh bound, compensated by these predictable squared coefficients, therefore gives

$$\mathbb{P}[Q \le -b] \le e^{-2b^2/(KdT)}. \tag{8}$$

Writing R for the matrix of rows r_i, condition (5) is $\|R\|^2 \le Kd$. Its leading $T \times T$ submatrix is the transpose of the map from the prefix to its gap changes. Hence, for every prefix,

$$\sum_{i \le T} \Delta_i^2 \le 2 \sum_{i \le T} h_i^2 + 2KdT \le 2(H+K)nd.$$

Conditional on earlier decisions, $X_i = [i \in I]|\Delta_i|/2$ has mean $|\Delta_i|/4$ and second moment $\Delta_i^2/8$. Using $e^{-\lambda X_i} \le 1 - \lambda X_i + \lambda^2 X_i^2/2$ and the predictable second-moment sum at most $(H+K)nd/4$ gives

$$\mathbb{P}\left[\sum_{i \le T} X_i < G(u)/4 - b\right] \le e^{-2b^2/((H+K)nd)}. \tag{9}$$

These conditional arguments require independent assignment coordinates, which the counting experiment supplies; no input signs are random.

Take $b = gn\sqrt{d}/8$ in (8)–(9). Outside the gap exception in (7) and the shortfall exception in (9), $S \geq \sum_i X_i \geq gn\sqrt{d}/8$. If also $Q > -gn\sqrt{d}/8$, then $S+Q > 0$. A union bound therefore gives

$$\frac{W(F)}{2^n} \leq e^{-\eta n} + e^{-g^2 n/(32(H+K))} + e^{-g^2 n/(32K)} \leq 3e^{-cn},$$

where $T \leq n$ was used in the Q bound. The cost is $O^*(2^T + W(F))$, proving the result. □

Thus there is a precise counting replacement for the two roles of independence. Pair counts control collective agreement even after adaptive selection and orientation of rows. The gap-profile count controls how frequently the order fails to produce useful imbalance. The latter condition is a sufficient structural counting hypothesis, not a restatement that few assignments survive the algorithm.

4 A transparent bipartite special case

Let the formula have a simple bipartite interaction graph with parts L and R, of sizes p and q. There is one clause per edge and no initial unit clauses. Every vertex of L has degree d; degrees in R need not be equal. Let h_i be the positive literal degree minus the negative literal degree of $i \in L$, and let $t_i \in \{-1, +1\}$ be its preferred value, using $+1$ to break a tie.

For edge iw, let J_{iw} be the product of its two literal signs. The learned inequality in the nonpreferred branch of i counts the bad value

$$b_i(w) = -t_i J_{iw}, \qquad L_i(y) \leq \frac{d - |h_i|}{2} - 1. \tag{10}$$

Set $b_i(w) = 0$ off the support. All these rows and their pair counts C_{ij} are defined directly from the input. They do not depend on which branches are later taken, because variables in L have no edges to one another.

Fix constants $a > 0$ and $K \geq 1$. Impose the following two conditions:

$$|h_i| \geq a\sqrt{d} \qquad \text{for every } i \in L, \tag{11}$$

$$d|I| + 2 \sum_{\substack{i<j \\ i,j \in I}} C_{ij} \leq Kd|I| \qquad \text{for every } I \subseteq L. \tag{12}$$

Condition (12) bounds the total excess of agreements over disagreements in every selected family. It permits nonzero correlations and arbitrary higher-order dependencies.

One stronger, polynomial-time checkable condition is

$$\sum_{j \neq i} \max(C_{ij}, 0) \leq (K-1)d \quad \text{for every } i \in L. \tag{13}$$

Indeed, the off-diagonal sum in (12) is at most the sum of these positive row sums over $i \in I$. Another sufficient certificate is $C \preceq Kd\,\mathrm{Id}$, which can be checked by linear algebra. That certificate is an equivalent spectral formulation of a stronger quadratic bound; it is not a separate independence assumption. The proof needs only the subset inequality (12).

Theorem 3 (Counting conditions suffice)**.** *Every input satisfying* (11)–(12) *can be solved exactly in polynomial space and time*

$$O^*\left(2^p + 2^q(1 + e^{-a^2/(2K)})^p\right). \tag{14}$$

If $p/(p+q)$ is bounded away from zero and one, this is $O^*((2-\delta)^{p+q})$ for a fixed $\delta > 0$ independent of d. The stronger condition (13)*, together with* (11)*, defines a class whose membership can be verified in polynomial time when the bipartition is supplied.*

Proof. Branch on every variable of L, learning (10) when its nonpreferred value is used. At each resulting prefix freeze the learned family and enumerate solutions of its summed inequality. The learned constraints preserve the first optimum in preferred-child-first order: violating a learned constraint permits a no-worse single flip to the preferred value, producing an earlier optimum, a contradiction. Their sum therefore also preserves an optimum.

There is exactly one prefix for each set I of nonpreferred decisions. For $k = |I| > 0$, its shortfall and variance parameter satisfy

$$s_I = \sum_{i \in I} (|h_i|/2 + 1) \geq ka\sqrt{d}/2, \qquad V_I \leq Kdk.$$

Lemma 1 bounds its candidate count by $N(I) \leq 2^q e^{-a^2 k/(2K)}$. The same formula holds for $I = \varnothing$, whose count is 2^q. Summing over all prefixes gives

$$\sum_{I \subseteq L} N(I) \leq 2^q \sum_{k=0}^{p} \binom{p}{k} e^{-a^2 k/(2K)} = 2^q (1 + e^{-a^2/(2K)})^p.$$

After orienting cheaper values, a depth-first generator of the nonnegative budget inequality takes $\mathrm{poly}(n)(N(I)+1)$ time: every retained partial assignment extends by choosing cheaper values on the remaining variables. This proves (14); the traversal uses polynomial space. If $p = \tau n$, the two exponential bases are 2^τ and $2((1 + e^{-a^2/(2K)})/2)^\tau$, both uniformly below two when τ stays in a compact subinterval of $(0, 1)$. □

This special case is a theorem about the constraint-guided enumeration mechanism, not a new tractability result for bipartite Max 2-SAT. Every bipartite formula already has an $O^*(2^{\min(p,q)})$ algorithm: enumerate the smaller side and optimize the variables on the other side independently. The value of the special case is its direct translation from static neighborhood counts to a bound on every selected learned family. Theorem 2 is not restricted to bipartite graphs or graphs with a large independent set.

5 An explicit dense example

Take $p = q = d = 4^k$ for any integer $k \geq 1$, with the complete bipartite interaction graph. Let H be a $q \times q$ sign matrix satisfying $HH^{\mathsf{T}} = q\,\mathrm{Id}$. Such matrices follow from the recursion

$$H_1 = (1), \qquad H_{2r} = \begin{pmatrix} H_r & H_r \\ H_r & -H_r \end{pmatrix};$$

the stated identity follows by block multiplication. Choose $u \in \{-1, +1\}^q$ with $\sum_w u_w = \sqrt{q}$. For each edge iw, set the literal sign at i to u_w and that at w to $u_w H_{iw}$. This specifies all input clauses explicitly.

Every left gap is $\sqrt{q}$, the interaction sign matrix is H, and the bad-value matrix is $-H$. Consequently $C_{ij} = 0$ for $i \neq j$: each pair agrees on exactly half its neighbors and disagrees on the other half. Conditions (11)–(12) hold with $a = K = 1$. Since $n = 2q$, Theorem 3 gives

$$O^*\left(\left[\sqrt{2(1 + e^{-1/2})}\right]^n\right) = O^*(1.792502^n) \subseteq O^*(1.793^n).$$

The clause density is $q/2 = n/4$. There is no random-input qualification. This is an explicit witness that the class is nonempty at growing density, not a claim that this structured example requires exponential time or that the constants are optimal.

6 Why cancellation counts alone do not suffice

The role of the gap hypothesis is real for this particular algorithm. Take a matrix with p orthogonal sign rows of length q, where $q \geq 16p$, and make it the interaction matrix of a complete bipartite formula. Choose the own-literal signs at each left variable to be balanced, and choose the other endpoint signs to realize the desired interaction products. Then every $h_i = 0$, while every distinct pair has $C_{ij} = 0$ exactly. For example, take the first $p = q/16$ rows of the recursively defined matrix above, for powers of two $q \geq 32$. These instances have clause density proportional to their number of variables.

Proposition 4 (Failure of the summed generator with zero gaps)**.** *For these formulas, the algorithm that branches on L and then enumerates the summed inequality generates at least*

$$\frac{3}{32}(2^p - 1)2^q$$

complete assignments. Thus its work is $\Omega(2^{p+q})$ despite perfectly balanced signed pair-agreement counts.

Proof. For any nonempty set I of k nonpreferred decisions, the total shortfall is $s_I = k$ and $V_I = qk$. Under a uniform right assignment, Q_I is a symmetric sum of independent signs, of variance $v = qk/4$ and fourth moment at most $3v^2$. For completeness, the elementary second-moment inequality for $X \geq 0$ is

$$\mathbb{P}[X \geq \theta \mathbb{E}X] \geq (1-\theta)^2 \frac{(\mathbb{E}X)^2}{\mathbb{E}X^2}, \qquad 0 \leq \theta < 1.$$

It follows by bounding the contribution from $X < \theta \mathbb{E}X$ and applying Cauchy–Schwarz to the remaining contribution. Apply it to $X = Q_I^2$ and $\theta = k^2/v = 4k/q \leq 1/4$. Then

$$\mathbb{P}[|Q_I| \geq k] \geq \frac{(1 - 4k/q)^2}{3} \geq \frac{3}{16}.$$

Symmetry and $k > 0$ give $\mathbb{P}[Q_I \leq -k] \geq 3/32$. These are exactly the completions admitted by the summed inequality. Sum their counts over nonempty I. □

This does not show that the formulas are hard for other branching orders, domination, individual-constraint propagation, or aggregation. It shows that initial literal balance and pairwise neighborhood balance do not, by themselves, give the earlier switching guarantee for this generator. One must also obtain sufficiently restrictive constraints.

7 What remains for the general branching approach

For a fixed residual constraint family, equations (1)–(3) and Lemma 1 already apply to arbitrary inputs. The remaining global task is to guarantee a large shortfall relative to V_I on sufficiently many prefixes, accounting for the cost of all other prefixes. In the bipartite theorem, static gaps and the binomial count of nonpreferred decisions settle this exactly. In a general formula, branching changes gaps, supports, and which rows are selected. Initial pair counts cannot simply be reused after these restrictions.

This gives a concrete form of the proposed dichotomy. Either learning creates a large total shortfall while the selected neighborhoods have controlled net agreement, and enumeration saves an exponential factor; or one must address insufficient gaps or a selected family with large positive agreement excess. The latter is a precisely countable alignment of bad values, but it is not yet a proof of a small separator or an aggregation rule with a global running-time bound.

The counts also give exact transitions for a fixed summed inequality. Write its current shortfall as s and let $B_w = \sum_i b_i(w)$, so $V = \sum_w B_w^2$. Assigning $y_w = \varepsilon$ changes these parameters to

$$s' = s + \frac{\varepsilon B_w}{2}, \qquad V' = V - B_w^2.$$

Thus the effect of choosing a variable can be read from occurrence counts directly. A large $|B_w|$ gives a large separation between the two children's shortfalls and removes a large contribution to V. This does not itself shrink the number of completions of the inherited inequality: their counts partition between the children. Further savings must come from additional constraints, domination, or another valid optimization argument.

Verification. The companion program `explore_counting_conditions.py` verifies the agreement identities, the upper bound and preservation of an optimum on small explicit instances, and the zero-gap lower bound using exact completion counts. All 548 prefix checks passed, including 65,792 full-assignment identity checks on the small positive-gap examples. Another 2,560 complete-assignment checks on signed complete graphs verified the ordered-count identities and quadratic bounds used in Theorem 2. These checks support the implementation; the theorems follow from the proofs above.

Related working note. The algorithm and output-sensitive budget generator are those of *Max 2-SAT by Early Constraint-Guided Enumeration*, 8 October 2026, local source `max2sat_early_enumeration.tex`. The present note replaces a probabilistic assumption by sufficient deterministic conditions for a narrower input class. It does not assert that the conditions are necessary or characterize typical random signings.