A Checkable Replacement for the Gap-Profile Condition

Addendum to *Replacing Random Signs by Neighborhood Counts*

8 October 2026

Abstract

The full deterministic counting criterion of [1] assumes three conditions: an aggregate bound on ordered pair counts, a bound on the energy of the initial gaps, and a bound on the number of exceptional gap profiles. The last condition has no evident polynomial-time membership test. We show that the first condition, together with a lower bound on initial gaps and earlier-neighbor counts that can be checked in linear time, implies the third. The criterion of [1] therefore holds for a class of formulas whose membership can be verified in polynomial time. The proof uses the observation of [1] that the pair-count condition bounds the map from prefixes to gap changes, together with Talagrand's concentration inequality for convex Lipschitz functions.

1 Setting

We use the notation of [1, Section 3]. The input is a simple signed interaction graph on n vertices, with one clause per edge and no initial unit clauses, a fixed order of the vertices, and a depth $T = \lfloor \tau n \rfloor$ with $0 < \tau < 1$. Let $d \geq 1$ be a reference degree scale, h_i the initial literal-degree gap of x_i, and J_{iw} the product of the two literal signs on the edge iw, with $J_{iw} = 0$ off edges.

Let R be the $T \times n$ matrix with $R_{iw} = J_{iw}$ for $w > i$ and $R_{iw} = 0$ otherwise, so that $M = RR^{\mathsf{T}}$ is the matrix of ordered pair counts $M_{ij} = \sum_{w > \max(i,j)} J_{iw} J_{jw}$. Let L_T be the $T \times T$ matrix with $(L_T)_{ij} = J_{ij}$ for $j < i$ and 0 otherwise. For a prefix $u \in \{-1, +1\}^T$ the gap of x_i is $h_i - (L_T u)_i$, and the total gap is

$$G(u) = \sum_{i \leq T} |h_i - (L_T u)_i|.$$

Finally, D_i^- is the number of neighbors $j < i$ of i. The three conditions of [1] are

$$z^{\mathsf{T}} M z \leq K d \, \|z\|^2 \qquad \text{for all } z \in \mathbb{R}^T, \tag{C1}$$

$$\sum_{i \leq T} h_i^2 \leq H n d, \tag{C2}$$

$$\#\{u \in \{-1, +1\}^T : G(u) < g n \sqrt{d}\} \leq 2^T e^{-\eta n}. \tag{C3}$$

We replace (C3) by

$$\sum_{i \leq T} \max\Big\{ |h_i|, \sqrt{D_i^- / 3} \Big\} \;\geq\; 2g \, n \sqrt{d}. \tag{C3$'$}$$

Condition (C1) can be checked by computing the largest eigenvalue of M; conditions (C2) and (C3′) can be checked in linear time.

2 The result

Proposition 1. *Assume* (C1) *and* (C3′), *and* $n \geq 256\pi\tau K/g^2$. *Then* (C3) *holds with* $\eta = g^2/(128\tau K)$.

Proof. Lipschitz constant. As observed in [1], L_T^{T} is the leading $T \times T$ block of R. A submatrix has norm at most that of the matrix, and (C1) says $\|R\|^2 = \|M\| \leq Kd$. Hence $\|L_T\| \leq \sqrt{Kd}$. The function G is convex on $\mathbb{R}^T$, as a sum of absolute values of affine functions, and

$$|G(u) - G(u')| \leq \sum_{i \leq T} |(L_T(u - u'))_i| \leq \sqrt{T}\, \|L_T\|\, \|u - u'\|_2.$$

So G is ℓ-Lipschitz with $\ell = \sqrt{TKd}$.

Mean. Let u be uniform. For $i \leq T$, the sum $X_i = (L_T u)_i = \sum_{j<i} J_{ij} u_j$ has D_i^- terms, each an independent uniform sign, so it is symmetric. By Jensen's inequality, $\mathbb{E}|h_i - X_i| \geq |h_i|$. By symmetry, $\mathbb{E}|h_i - X_i| = \frac{1}{2}\mathbb{E}(|h_i - X_i| + |h_i + X_i|) \geq \mathbb{E}|X_i|$. Since $\mathbb{E}X_i^4 \leq 3(D_i^-)^2$, Hölder's inequality $\mathbb{E}X_i^2 \leq (\mathbb{E}|X_i|)^{2/3}(\mathbb{E}X_i^4)^{1/3}$ gives $\mathbb{E}|X_i| \geq \sqrt{D_i^-/3}$, as in [2]. By (C3′),

$$\mathbb{E}G \;\geq\; \sum_{i \leq T} \max\Big\{|h_i|, \sqrt{D_i^-/3}\Big\} \;\geq\; 2g\, n\sqrt{d}.$$

Concentration. Let f be convex and ℓ-Lipschitz on $\mathbb{R}^T$, and let the coordinates of u be independent with values in $[-1, 1]$. Then Talagrand's inequality gives $\mathbb{P}(|f(u) - m| \geq s) \leq 4e^{-s^2/(16\ell^2)}$ for every $s \geq 0$, where m is a median of $f(u)$ [5]. The inequality is usually stated for $[0, 1]$ with exponent $-s^2/4$ for $\ell = 1$ [4, Chapter 4]; the factor 16 comes from rescaling to $[-1, 1]$. Integrating the tail gives $|\mathbb{E}f - m| \leq 8\sqrt{\pi}\,\ell$. Now $\ell^2 = TKd \leq \tau n K d$, and the assumption on n gives $8\sqrt{\pi}\,\ell \leq \frac{1}{2} g n \sqrt{d}$. Therefore

$$\mathbb{P}(G(u) < gn\sqrt{d}) \leq \mathbb{P}(G(u) \leq m - \tfrac{1}{2}gn\sqrt{d}) \leq 4\exp\Big(-\frac{g^2 n}{64\tau K}\Big) \leq \exp\Big(-\frac{g^2 n}{128\tau K}\Big).$$

The last step uses $g^2 n/(128\tau K) \geq 2\pi > \ln 4$, which also follows from the assumption on n. Multiplying by 2^T gives (C3). □

Corollary 2 (A criterion checkable in polynomial time)**.** *Every fixed formula that satisfies* (C1), (C2) *and* (C3′), *with* $n \geq 256\pi\tau K/g^2$, *is solved exactly by the algorithm of [1, Theorem 2]. That algorithm branches in the fixed order through depth* T *and then enumerates the summed learned inequality. It uses polynomial space and time*

$$O^*(2^T + 2^n e^{-cn}), \qquad c = \min\Big\{\frac{g^2}{128\tau K},\ \frac{g^2}{32(H + K)}\Big\}.$$

Membership in the class defined by (C1), (C2) *and* (C3′) *can be verified in polynomial time.*

Proof. Proposition 1 gives (C3) with $\eta = g^2/(128\tau K)$; then apply [1, Theorem 2]. □

3 Remarks

Remark 3 (The two sources of gap). The two terms in (C3′) are the two sources of shortfall seen so far: initial gaps, used in the bipartite theorem of [1], and earlier neighbors, used in the random-sign proofs [2]. The zero-gap formulas of [1, Section 6], with the left side first, have $h_i = 0$ and $D_i^- = 0$ for every $i \leq T$. So (C3′) fails there, as it must.

Remark 4 (The examples of [1]). For the Hadamard example of [1, Section 5], with the left side first and $T = n/2$, the conditions hold with $K = 1$, $H = \frac{1}{2}$ and $g = \frac{1}{4}$. Here G is constant, equal to $\frac{1}{2}n\sqrt{d}$. The corollary then gives $c = 1/1024$, which is far weaker than the bound $O^*(1.793^n)$ of the bipartite theorem; the general criterion is not tuned. On random signed graphs in the natural order with $T = n/2$, the conditions hold with $K \approx 2.4$, $H \approx 0.5$ and $g \approx 0.21$, independently of the density (Table 1). The constants of Proposition 1 are conservative: at these sizes its bound is still above one, while the observed standard deviation of G is about $0.01\, n\sqrt{d}$. That random signed graphs with $d \geq C \log n$ satisfy (C1) with bounded K follows from norm bounds for random matrices with independent entries [3]; we do not use this here.

n	d	K	H	g	$\mathbb{E}G$	sd of G	min of G	row-sum K
400	19.6	2.50	0.50	0.203	0.435	0.015	0.381	9.6
400	81.2	2.29	0.55	0.216	0.461	0.015	0.405	9.0
400	199.4	2.42	0.50	0.211	0.452	0.014	0.400	8.6
800	100.6	2.43	0.48	0.200	0.429	0.011	0.395	12.2
800	400.7	2.41	0.49	0.206	0.442	0.011	0.405	12.1

Table 1: Random signed graphs in the natural order, $T = n/2$. Here K, H and g are the smallest constants for which (C1), (C2) and (C3′) hold. The values of G are in units of $n\sqrt{d}$, over 4000 uniform prefixes. The last column is the constant needed by the row-sum certificate $\max_i \sum_j |M_{ij}| \leq Kd$.

Remark 5 (Certificates for (C1)). The stronger condition $\sum_j |M_{ij}| \leq Kd$ for every i, mentioned in [1], fails for random signed graphs. The required K grows like $\sqrt{n}$: in the runs of Table 1 it is about 9 for $n = 400$ and 12 for $n = 800$, at every density (last column). For a class that contains typical formulas, condition (C1) has to remain an aggregate quadratic-form condition.

Remark 6 (Adaptive orders). The proposition concerns a fixed order, which defines D_i^-, R and G. For an adaptive first phase the order depends on the prefix, so G is no longer a single convex function of u, and the argument does not apply as it stands.

Checks. `explore_counting_addendum.py` computes the quantities of Table 1, confirms that no sampled prefix falls below $gn\sqrt{d}$, and evaluates the Hadamard and zero-gap examples of [1].

References

[1] *Replacing Random Signs by Neighborhood Counts: a deterministic early-enumeration theorem and a limitation*, working note, 8 October 2026. Local source: `max2sat_counting_conditions.tex`.

[2] *Max 2-SAT by Early Constraint-Guided Enumeration*, working note, 8 October 2026. Local source: `max2sat_early_enumeration.tex`.

[3] A. S. Bandeira and R. van Handel. Sharp nonasymptotic bounds on the norm of random matrices with independent entries. *The Annals of Probability* 44(4):2479–2506, 2016.

[4] M. Ledoux. *The Concentration of Measure Phenomenon*. Mathematical Surveys and Monographs 89, American Mathematical Society, 2001.

[5] M. Talagrand. An isoperimetric theorem on the cube and the Khintchine–Kahane inequalities. *Proceedings of the American Mathematical Society* 104(3):905–909, 1988.