Max 2-SAT and the Domination Rule: Results and Ideas So Far

5 October 2026

Working notes, updated

Abstract

We collect the current state of our work on deterministic polynomial-space algorithms for Max 2-SAT analyzed in the number n of variables. Two earlier notes give an additive deficit measure, which certifies savings for bounded deficits, and an exponential saving for formulas in which a linear number of variables have an absolute linear polarity gap. For DPLL with the domination rule we add a *level budget*: for every static variable order, the number of leaves is at most one plus the number of predecessor assignments that leave a variable undominated, summed over all variables. For formulas with at most one binary clause per pair this gives $O^*(2^{\max\{1-\eta,\, H((1-\varepsilon)/2)\}n})$, and a circulant family shows that no DPLL algorithm whose only simplification is domination can do better on the gap term. We then study an extension of domination: when branching on x, the branch against the literal ℓ closer to domination keeps only the assignments under which ℓ would not become dominated, recorded as a *pruning clause* $\langle L_x \le p_\ell - 1\rangle$. The search stays exact when these clauses are nested and combined with domination; with them the gap bound holds for every static order; and with disjoint supports their savings multiply exactly. With overlapping supports two things limit the savings: correlation, measured by the signed Gram matrix of the clauses, and timing, since a clause is decided only once most of its support is assigned. Finally we record a spectral view, domination for sets of variables, and two special cases that separate the remaining difficulties.

1 Goal and context

Goal. A deterministic algorithm that solves Max 2-SAT in time $O^*(c^n)$ for a constant $c < 2$, using polynomial space, for formulas of *every* clause density. Williams' algorithm runs in time $O^*(2^{\omega n/3})$ but needs exponential space [3]; beating 2^n in polynomial space is open [3, 4], already for Max-Cut. The known polynomial-space bounds in terms of n depend on the average degree d and save only $O((\log d)/d) \cdot n$ [5, 6, 7]. They give c^n only for constant density.

Approach. Identify classes of formulas of unbounded density that DPLL with the domination rule solves in time c^n, understand which proof techniques capture the saving, and then extend the rule. The first class: a linear number of variables with an absolute linear polarity gap [2].

2 Setting and notation

We follow [1, 2]. A formula is *canonical* if it consists of unit clauses, kept with multiplicity, and binary clauses on distinct variables such that every unordered pair carries zero clauses,

one clause, or two clauses complementary in both positions. Let $B \in \{1, 2\}$ bound the number of binary clauses per pair. Exact pair reductions [2, Sec. 2] bring every unweighted 2-CNF without repeated binary clauses into canonical form. They preserve the score up to an additive offset and preserve the effective gaps defined below. If every variable has distinct neighbors in the input, then $B = 1$.

For a literal ℓ, let $d_1(\ell)$ and $d_2(\ell)$ count its occurrences in unit and binary clauses. Its *deficit* is

$$p_\ell = d_2(\neg\ell) - d_1(\ell) + d_1(\neg\ell).$$

For a variable v put $D_v = d_2(v) + d_2(\neg v)$ and $\Delta_v = d_2(v) - d_2(\neg v) + 2\big(d_1(v) - d_1(\neg v)\big)$. Then

$$p_v = \frac{D_v - \Delta_v}{2}, \qquad p_{\neg v} = \frac{D_v + \Delta_v}{2}, \qquad p_v + p_{\neg v} = D_v. \tag{1}$$

For a partial assignment σ, $F|_\sigma$ denotes substitution: delete satisfied clauses and shorten the others, with no further simplification. Canceling opposite unit clauses never changes a deficit and is ignored below. We write $H(t) = -t\log_2 t - (1-t)\log_2(1-t)$ for the binary entropy.

Lemma 1 (Domination [1, 2])**.** *If $p_\ell \le 0$, some optimal assignment makes ℓ true.*

Lemma 2 (Local update [2])**.** *Let $c_{xv} \in \{0, 1, 2\}$ be the number of binary clauses on $\{x, v\}$. Assigning x lowers exactly one of $p_v, p_{\neg v}$ by c_{xv}, for every unassigned $v \neq x$; which one depends on the value of x and on the clause signs. Non-neighbors are unaffected.*

Corollary 3 (Monotonicity)**.** *Deficits never increase under assignments. In particular, if v is dominated in $F|_\sigma$, then it is dominated in $F|_{\sigma\tau}$ for every assignment τ to further variables other than v.*

DPLL with domination. *Closure* fixes dominated variables until none is left. If no variable remains, the node is a leaf; otherwise a branching rule picks a variable and both values are explored recursively. Let $L(F)$ be the number of leaves, with unary closure steps contracted, so that every internal node has two children. The *static rule* A_π for an order π always branches on the first unassigned variable of π. Depth-first search uses polynomial space, and the running time is $O^*(L(F))$.

3 Results of the earlier notes

Theorem 4 (Literal-deficit measure [1])**.** *Let $B = 1$ and $q_v = \min\{p_v, p_{\neg v}\}$. Branching on a variable of maximum q_v and using the measure $\mu(F) = \sum_v \big(1 - \frac{1}{3q_v}\big)$ over the undominated variables gives $L(F) \le 2^{\mu(F)}$, that is, time $O^*\big(2^{n - \frac{1}{3}\sum_v 1/q_v}\big)$.*

The saving comes only from variables of bounded deficit. For a linear gap, $q_v = \Theta(n)$ and the certified saving is $O(1)$; see Section 7.2 for why additive measures cannot do better here.

Theorem 5 (Absolute linear gap [2])**.** *Let the input be an unweighted 2-CNF without unit clauses and without repeated binary clauses, and suppose at least ηn variables satisfy $|d_2(v) - d_2(\neg v)| \ge \varepsilon n$ in the input. Then Max 2-SAT can be solved deterministically in polynomial space and time $O^*(2^{(1-\kappa)n})$ with $\kappa = \min\{\eta, \varepsilon^2/(4B^2)\}$.*

The proof reserves a set $L \subseteq S$ of size at most $\lfloor \varepsilon^2 n/(4B^2) \rfloor$ and enumerates all assignments to $H = V \setminus L$. Hoeffding's inequality and a union bound show that all of L is dominated except for a $2^{-\gamma n}$ fraction of these assignments, with $\gamma > |L|/n$. The search-work budget $W(F; H, L) = 2^h + 2^s \sum_{v \in L} N_v(F; H)$ also covers domination applied during the search.

Remark 6 (Constant). With $|L| = \alpha n$ the margin is $(\varepsilon - B\alpha)n/2$, so the failure exponent is $\gamma(\alpha) = (\varepsilon - B\alpha)^2/(2B^2 \ln 2)$ and the saving is $\min\{\eta, \alpha, \gamma(\alpha)\}$. Choosing $\alpha = \gamma(\alpha)$ gives about $\varepsilon^2/(2B^2 \ln 2) \approx 0.72\,\varepsilon^2/B^2$ for small ε. Theorem 11 below removes the loss $B\alpha$ and the restriction to two blocks.

4 The level budget

Let $\pi = (x_1, \ldots, x_r)$ be an order of the variables of F and $P_k = \{x_1, \ldots, x_{k-1}\}$. Define

$$N_k(F, \pi) = \big|\{\sigma \in \{0,1\}^{P_k} : p_{x_k}(F|_\sigma) > 0 \text{ and } p_{\neg x_k}(F|_\sigma) > 0\}\big|,$$
$$B_\pi(F) = 1 + \sum_{k=1}^{r} N_k(F, \pi). \tag{2}$$

Thus N_k counts the assignments to the predecessors of x_k under which x_k is dominated in neither polarity.

Lemma 7 (Level budget)**.** *For every canonical formula F and every order π, $L_{A_\pi}(F) \le B_\pi(F)$.*

Proof. Since every internal node has two children, $L_{A_\pi}(F) = 1 +$ (number of internal nodes). Let u be an internal node that branches on x_k. The rule A_π branches on the first unassigned variable of π, so all of P_k is assigned at u; let $\sigma_u \in \{0,1\}^{P_k}$ record these values, whether they were branched on or fixed by closure. The formula at u arises from $F|_{\sigma_u}$ by assigning further variables that closure fixed. Since x_k is not dominated at u, Corollary 3 shows that x_k is dominated in neither polarity in $F|_{\sigma_u}$.

It remains to show that $u \mapsto \sigma_u$ is injective on the internal nodes branching on x_k. Two distinct such nodes u, u' are incomparable, because x_k is still unassigned at both. Their lowest common ancestor w is an internal node branching on some x_j. Since w is a proper ancestor of u, the variable x_k is unassigned at w and $j \neq k$; as A_π picks the first unassigned variable, $j < k$. The nodes lie below different children of w, so $\sigma_u(x_j) \neq \sigma_{u'}(x_j)$. Hence at most $N_k(F, \pi)$ internal nodes branch on x_k. □

Remark 8 (Two blocks). For $\pi = (H, L)$ with $L = (v_1, \ldots, v_s)$, positions in H contribute at most $2^h - 1$. By Corollary 3, $N_{h+j}(F, \pi) \le 2^{j-1} N_{v_j}(F; H)$. Hence $B_\pi(F) \le 2^h + \sum_j 2^{j-1} N_{v_j}(F; H) \le W(F; H, L)$: the budget of [2, Sec. 7] is the two-block case, with the factor 2^s replaced by 2^{j-1}.

Lemma 9 (Computing N_k)**.** *Let U be the set of neighbors of $x = x_k$ in P_k, put $c_u = c_{ux}$ and $K = \sum_{u \in U} c_u$, and let a_j be the coefficient of z^j in $\prod_{u \in U}(1 + z^{c_u})$. Then*

$$N_k(F, \pi) = 2^{k-1-|U|} \sum_j a_j \,[\, p_x - j > 0 \,]\,[\, p_{\neg x} - (K - j) > 0 \,].$$

In particular N_k and $B_\pi(F)$ are computable in polynomial time.

Proof. By Lemma 2, each $u \in U$ lowers p_x by c_u for one of its values and $p_{\neg x}$ by c_u for the other; predecessors outside U do not affect x. Thus a_j counts the assignments to U that lower p_x by exactly j, and then $p_{\neg x}$ is lowered by $K - j$. □

Proposition 10 (A measure for adaptive branching)**.** *For a formula F let $\Phi(F) = \min_\pi B_\pi(F)$ over all orders of its variables, and $\Phi(F) = 1$ if none remain.*

1. *If F' arises from F by fixing a dominated variable, then $\Phi(F') \le \Phi(F)$.*

2. *If x is undominated in F, then for every order π' of the remaining variables, $B_{(x,\pi')}(F) = B_{\pi'}(F|_{x=0}) + B_{\pi'}(F|_{x=1})$.*

Consequently, the algorithm that after closure branches on the first variable of an order minimizing B_π has at most $\Phi(F)$ leaves.

Proof. (2) The first position contributes $N_1 = 1$ because x is undominated. For a later position, split the assignments to the predecessors according to the value of x; this gives $N_k(F,(x,\pi')) = N_{k-1}(F|_{x=0},\pi') + N_{k-1}(F|_{x=1},\pi')$. Summing, $B_{(x,\pi')}(F) = 2 + \sum_j (N_j(F|_{x=0},\pi') + N_j(F|_{x=1},\pi'))$, which is the claim.

(1) Let y be fixed to b and let π be any order of the variables of F; write $\pi \setminus y$ for its restriction. For a variable before y in π, the predecessor set is unchanged, and the extra assignment $y = b$ can only lower its deficits. For a variable after y, the count in F' equals the number of assignments to its predecessors in F with $y = b$ that leave it undominated. The term of y disappears. Hence $B_{\pi\setminus y}(F') \le B_\pi(F)$, and minimizing over π gives $\Phi(F') \le \Phi(F)$.

For the consequence, use induction on the number of variables. Closure does not increase Φ by (1). If no variable remains, $L = 1 \le \Phi$. Otherwise let (x,π') minimize B_π for the closed formula G; then x is undominated, and $L(G) = L(G|_{x=0}) + L(G|_{x=1}) \le \Phi(G|_{x=0}) + \Phi(G|_{x=1}) \le B_{\pi'}(G|_{x=0}) + B_{\pi'}(G|_{x=1}) = B_{(x,\pi')}(G) = \Phi(G)$. □

Φ is a measure of the formula alone. It is not obviously computable, but for every concrete order the bound $L_{A_\pi} \le B_\pi$ is certified in polynomial time by Lemma 9, so orders can be optimized heuristically per instance.

5 A tight bound for the absolute linear gap class

Theorem 11. *Let F be canonical with $B = 1$ on n variables, let $\varepsilon \in (0,1]$, and let S be a set of variables with $|\Delta_v(F)| \ge \varepsilon n$ for all $v \in S$. For every order π that lists $V \setminus S$ before S,*

$$L_{A_\pi}(F) \le 2^{\,n-|S|} + |S| \cdot 2^{\,H((1-\varepsilon)/2)\,n}.$$

Hence, if $|S| \ge \eta n$, DPLL with domination solves Max 2-SAT on F in polynomial space and time

$$O^*\big(2^{\max\{1-\eta,\, H((1-\varepsilon)/2)\}\,n}\big).$$

For $\eta = 1$ every order works.

The saving in the exponent is $\min\{\eta, 1-H((1-\varepsilon)/2)\}$, and $1-H((1-\varepsilon)/2) = \varepsilon^2/(2\ln 2) + O(\varepsilon^4)$. By gap preservation [2, Lemma 3], the hypothesis holds for the class of Theorem 5 whenever canonical form has no complementary pairs.

Proof. Use Lemma 7. Positions in $V \setminus S$ contribute at most $\sum_{k \le n-|S|} 2^{k-1} < 2^{n-|S|}$. Fix a position k with $x = x_k \in S$, rename so that $\Delta := \Delta_x(F) \ge \varepsilon n$, and let K and R be the numbers of neighbors of x before and after position k. Then $D_x = K + R$ and $p_x = (K + R - \Delta)/2$ by (1). If $\Delta \ge D_x$, then x is dominated under every σ and $N_k = 0$; assume $\Delta < D_x$.

For uniform σ on P_k, the number X of neighbors in P_k whose value lowers p_x is $\mathrm{Bin}(K, \frac{1}{2})$, by Lemma 2 with $B = 1$. An undominated x needs $p_x - X > 0$, so

$$N_k \le 2^m \Pr\big[X \le K/2 - g/2\big], \qquad m := k-1 \ge K, \quad g := \Delta - R \le K.$$

Case $m < (1-\varepsilon)n$. Then $N_k \le 2^m \le 2^{(1-\varepsilon)n} \le 2^{H((1-\varepsilon)/2)n}$, since $H(t) \ge 2t$ on $[0, \frac{1}{2}]$.

Case $m \ge (1-\varepsilon)n$. Since $R \le n - k = n - 1 - m$, we have $g \ge m + 1 - (1-\varepsilon)n > 0$. The binomial tail bound $\Pr[\mathrm{Bin}(K, \frac{1}{2}) \le K/2 - t] \le 2^{-K\varphi(t/K)}$ with $\varphi(u) = 1 - H(\frac{1}{2} - u)$

gives $N_k \le 2^{m-K\varphi(g/(2K))}$. As φ is convex with $\varphi(0) = 0$, $\varphi(u)/u$ is nondecreasing, so $K \mapsto K\varphi(g/(2K))$ is nonincreasing; with $K \le m$,

$$N_k \le 2^{\,mH(\frac{1}{2}-\frac{g}{2m})} \le 2^{\,mH(c/m)}, \qquad c := \tfrac{(1-\varepsilon)n}{2},$$

using that H increases on $[0, \frac{1}{2}]$ and $\frac{1}{2} - \frac{g}{2m} \le \frac{(1-\varepsilon)n-1}{2m} \le \frac{c}{m} \le \frac{1}{2}$. Finally $\psi(m) = mH(c/m)$ satisfies $\psi'(m) = -\log_2(1 - c/m) > 0$, so $\psi(m) \le \psi(n) = nH((1-\varepsilon)/2)$. Summing over the at most $|S|$ positions of S proves the bound. □

The two monotonicity facts in the proof have a simple meaning: for a fixed gap, more neighbors mean more noise, so the worst case is that all predecessors are neighbors; and an earlier position saves on the factor 2^{k-1} but loses more in the margin, so the last position dominates.

Remark 12 ($B = 2$). With complementary pairs, $X = \sum_u c_u Z_u$ with $c_u \in \{1, 2\}$. Hoeffding's inequality with $Q \le B^2 m$ and margin $(\Delta - R')/2$, where $R' \le B(n-1-m)$ is the weighted number of later neighbors, gives $N_k \le 2^{\,m-(\varepsilon n - B(n-1-m))^2/(2B^2 m \ln 2)}$. The exponent is decreasing in $n-1-m$ on the relevant range, so the last position dominates again and $L_{A_\pi}(F) \le 2^{n-|S|} + |S|\, 2^{(1-\varepsilon^2/(2B^2 \ln 2))n}$. This improves $\varepsilon^2/(4B^2)$ by the factor $2/\ln 2 \approx 2.9$. The exact rate function for $B = 2$ has not been worked out.

ε	Thm. 5 ($1 - \varepsilon^2/4$)	Rem. 6 (optimized)	Thm. 11 ($H(\frac{1-\varepsilon}{2})$)	circulant, $n = 3201$
0.2	0.990	0.977	0.971	0.969
0.4	0.960	0.924	0.881	0.879
0.6	0.910	0.852	0.722	0.720

Table 1: Exponents (in units of n) for $B = 1$ and η large. The last column is the exact $\frac{1}{n}\log_2$ of the tree size on the family of Theorem 13, which every branching rule attains.

6 A matching lower bound for domination-based DPLL

For odd n and an integer c with $(n-1)/4 \le c < (n-1)/2$, let $C_{n,c}$ contain, for each pair $i < j$, the clause $(x_i \vee x_j)$ if the circular distance $\min\{j - i, n - j + i\}$ is at most c, and $(\neg x_i \vee \neg x_j)$ otherwise. Every variable has $p_v = a := n - 1 - 2c$ and $p_{\neg v} = b := 2c$, hence the same gap $\Delta = 4c - (n-1)$.

Theorem 13. *Every DPLL algorithm on $C_{n,c}$ whose simplifications are domination fixings and opposite-unit cancellations has exactly $\binom{n-1}{a}$ leaves, for every branching rule. Choosing c with $\Delta \ge \varepsilon n$ and $\Delta = \varepsilon n + O(1)$, where $0 < \varepsilon < 1$, gives $2^{H((1-\varepsilon)/2)n - O(\log n)}$ leaves, while every variable has gap at least εn and $B = 1$.*

Proof. Every pair carries exactly one clause, of the form $(u \vee v)$ or $(\neg u \vee \neg v)$. Assigning $u := 0$ turns $(u \vee v)$ into the unit (v) and satisfies $(\neg u \vee \neg v)$; in both cases p_v drops by one. Symmetrically, $u := 1$ lowers $p_{\neg v}$ by one. Hence at a node where z zeros and o ones have been assigned, by branching or by closure, every unassigned variable has deficits $(a - z, b - o)$, and $a - z + b - o$ equals its current degree. If $z < a$ and $o < b$, no variable is dominated and the node branches, with children of types $(z+1, o)$ and $(z, o+1)$; as soon as $z = a$ or $o = b$, closure fixes all remaining variables and the node is a leaf. Thus the number of leaves $\lambda(a', b')$ below a node with deficits (a', b') satisfies $\lambda(a', b') = \lambda(a'-1, b') + \lambda(a', b'-1)$ and $\lambda(0, \cdot) = \lambda(\cdot, 0) = 1$, so $\lambda(a, b) = \binom{a+b}{a}$ with $a + b = n - 1$. The asymptotic form follows from $\binom{N}{K} \ge 2^{NH(K/N)}/(N+1)$. □

Remark 14 (Exactness of the budget). On $C_{n,c}$ the predecessors of x_k are all its neighbors, and x_k is undominated in $F|_\sigma$ if and only if σ has fewer than a zeros and fewer than b ones. Hence $B_\pi(C_{n,c}) = 1 + \sum_{z<a,\, o<b} \binom{z+o}{z} = \binom{a+b}{a}$ for every order: the level budget is exact on this family.

Consequences. For formulas with $B = 1$ the gap term $H((1-\varepsilon)/2)$ is optimal for domination alone. Any extension of the domination rule that aims at a better exponent must fire on $C_{n,c}$; this family is the benchmark for the next step. Note that $C_{n,c}$ is strongly aligned: all clauses are $(u \vee v)$ or $(\neg u \vee \neg v)$, so one value of each assigned variable helps all remaining variables at once.

7 Branching rules and measures: what we learned

7.1 Maximum-degree branching

Remark 15 (Sketch, not verified numerically). For $\eta < 1$, maximum-degree branching can lose the whole saving. Let the gap variables S form a clique with balanced signs inside S. Join S to every variable of $R = V \setminus S$ with signs biased at the S-end, so that every $v \in S$ has gap $\varepsilon'|R| \geq \varepsilon n$, and balanced at the R-end. Let R be dense with balanced signs but sparser than S. Then S has the larger degree, and each branching on S lowers every degree in $S \cup R$ by one, so all of S is branched before any of R. A variable of S has deficit $(|S| - 1 + (1-\varepsilon')|R|)/2$ in its gap direction, and during this phase only its $|S|-1$ neighbors in S are assigned; so it is never dominated if $|S| - 1 < (1-\varepsilon')|R|$. Afterwards the variables of R have unit imbalance $O(\sqrt{n})$ against degree $\Theta(n)$, so the remaining search is essentially exhaustive and the total is $2^{n-o(n)}$. The last step is heuristic, and simulations with $n \leq 20$ are too small to show the effect. For $\eta = 1$ no adaptive rule is needed, since every static order is optimal by Theorem 11.

7.2 Additive measures

Observation 16. Let $W(p,q) = \log_2 \binom{p+q}{p}/(p+q+1)$ for $p, q \geq 1$ and $W = 0$ otherwise, and $\mu(F) = \sum_v W(p_v, p_{\neg v})$. Then $W \approx H(p/D) \approx 1 - (\Delta/D)^2/(2\ln 2)$ depends only on the variable's own deficits.

1. On $C_{n,c}$, at every node, 2^μ equals the number of leaves below, and the branching inequality $2^{\mu(F_0)} + 2^{\mu(F_1)} \leq 2^{\mu(F)}$ holds with equality by Pascal's rule.

2. On heterogeneous instances the inequality fails. With maximum-degree branching on random aligned instances with $n = 12$ it was violated in 28 of 357 branchings, with ratio up to 1.17.

The reason is that W is not monotone: moving toward balance raises the entropy, for example $W(1,8) > W(1,9)$. In the branch that is unfavorable for a gap variable its weight goes up, and nothing compensates when the branching variable itself carries no gap.

The same mechanism explains why the measure of Theorem 4 certifies only bounded deficits: an additive measure must certify the saving branch by branch, while for linear gaps the saving is a large-deviation effect spread over many branches.

7.3 The counting view

For any search tree, choose a child uniformly at random at every internal node and let β be the number of branchings on the resulting root-to-leaf walk. A leaf at branching depth β_ℓ is reached with probability $2^{-\beta_\ell}$, so

$$L = \mathbb{E}[2^\beta] = \sum_{T \subseteq V} \Pr[\text{every } v \in T \text{ is branched on}],$$

where the second form uses $2^\beta = \prod_v (1 + [v \text{ branched}])$. An additive measure with $2^{\mu(F_0)} + 2^{\mu(F_1)} \leq 2^{\mu(F)}$ makes $2^\beta 2^\mu$ a supermartingale along the walk, so it has to certify the inequality at every single node, for the worst sign pattern. The gain per neighbor is of order $1/D$, and the gains must compensate in both branches; this works on perfectly symmetric instances (Observation 16) but not in general. The level budget counts failures level by level instead, and is valid for arbitrary correlations between the neighbors' progress.

7.4 What the method needs from a reduction rule

Lemma 7 and Proposition 10 use only two properties of domination:

(R1) *Soundness:* applying the rule preserves some optimal assignment.

(R2) *Monotonicity:* if the rule fixes v in $F|_\sigma$, it fixes v in $F|_{\sigma\tau}$ for every further assignment τ that does not touch v.

For any rule with (R1) and (R2) that fixes single variables, the level budget holds verbatim with N_k counting the predecessor assignments under which the rule does not fire on x_k. Rules that fix several variables at once or impose relations between variables need an adapted version. Pruning clauses (Section 8) are not fixing rules, but they are monotone in the same sense: the number of true literals of a clause only grows, so a violated clause stays violated. Lemma 23 uses exactly this.

Remark 17 (Limits of domination). For the budget to certify a saving κn, every position $k > (1-\kappa)n$ must have $\Pr_\sigma[x_k \text{ undominated}] \leq 2^{(1-\kappa)n-k+1}$, which is exponentially small for the last positions. For $X \sim \mathrm{Bin}(K, \frac{1}{2})$ this requires $(\Delta - R)^2 = \Omega(nK)$ there; since $\Delta \leq D \leq Bn$, the variables placed last must have gap and degree linear in n. Variables with smaller gaps yield only polynomial savings. This is not an artifact of the analysis: $C_{n,c}$ with gap $\Delta = o(n)$ has $\binom{n-1}{a} = 2^{n-o(n)}$ leaves for every branching rule by Theorem 13. Going beyond absolute linear gaps needs a stronger rule.

8 Pruning clauses

Let x be an undominated variable and ℓ one of its literals; the search below uses the literal with the smaller deficit, the one closer to domination. The *pruning literals* of x with respect to ℓ are

- the literal $\neg m$ for every binary clause $(\ell \vee m)$, and
- the literal m for every binary clause $(\neg\ell \vee m)$.

We write L_x for this set, with multiplicities if $B = 2$. By Lemma 2, the pruning literals are exactly the neighbour literals whose truth lowers p_ℓ. In the notation of the original formulation, with $d_1(x) = k$, $d_1(\neg x) = 0$, $d_2(\neg x) = q$, $d_2(x) = r$ and $\ell = x$, we have $|L_x| = q + r$ and $p_\ell = q - k$.

Lemma 18 (Flip identity)**.** *Let τ assign all variables other than x, and let $c(\tau)$ be the number of literals of L_x that are true under τ. Then*

$$\text{score}(\tau,\ \ell \text{ true}) - \text{score}(\tau,\ \ell \text{ false}) = c(\tau) - p_\ell.$$

In particular, ℓ is dominated in $F|_\tau$ if and only if $c(\tau) \geq p_\ell$.

Proof. Making ℓ true changes the score of the unit clauses on x by $d_1(\ell) - d_1(\neg\ell)$. A clause $(\ell \vee m)$ gains one exactly when m is false, that is, when its pruning literal $\neg m$ is true. A clause $(\neg\ell \vee m)$ loses one exactly when m is false, that is, when its pruning literal m is false. Hence the difference is $d_1(\ell) - d_1(\neg\ell) + c(\tau) - d_2(\neg\ell) = c(\tau) - p_\ell$. □

The *pruning clause* of x is $P_x = \langle L_x \leq p_\ell - 1\rangle$: an assignment satisfies it if at most $p_\ell - 1$ literals of L_x are true. By Lemma 18, every complete assignment that makes ℓ false and violates P_x is at least as good after flipping x. So when we branch on x, the child with ℓ true is explored in full, and the child with ℓ false only has to cover the assignments that satisfy P_x.

Remark 19 (One clause, not two). The two kinds of progress add up. If s true pruning literals come from clauses $(\neg\ell \vee m)$ and t from clauses $(\ell \vee m)$, the condition is $s + t \leq p_\ell - 1$. Two separate clauses, one for each kind and each with the full budget $p_\ell - 1$, are implied and therefore sound, but weaker: they keep $s = t = p_\ell - 1$. Splitting the budget, $s \leq s_1$ and $t \leq s_2$ with $s_1 + s_2 = p_\ell - 1$, is not implied and would lose optimal assignments. The clause is only an upper bound: nothing forces any pruning literal to be true.

Soft and hard clauses. The clauses of F are soft: they define the score and never force a value; only domination fixes variables from them. A pruning clause is hard for the subtree in which it was created, because that subtree is responsible only for the assignments that satisfy it. Accordingly, a node is *cut* as soon as an active pruning clause has more true literals than its bound, and *propagation* sets every remaining literal of a pruning clause false once the number of its true literals equals its bound. Propagation is a shortcut: it fixes values on which the search would otherwise branch and immediately cut one side.

The search. Solve$(F, \mathcal{P})$ takes a formula F and a set $\mathcal{P}$ of active pruning clauses.

1. Repeat until nothing changes:
   (a) if some $\langle L \leq b\rangle \in \mathcal{P}$ has more than b true literals, return $-\infty$ (cut);
   (b) if some literal ℓ has $p_\ell \leq 0$, set ℓ true (domination);
   (c) if some $\langle L \leq b\rangle \in \mathcal{P}$ has exactly b true literals and an unassigned literal $\lambda \in L$, set λ false (propagation).
2. If no variable is left, return the score.
3. Choose a variable x, and let ℓ be its literal with the smaller deficit; $p_\ell \geq 1$ because x is undominated.
4. Return $\max\{\text{Solve}(F|_{\ell \text{ true}}, \mathcal{P}),\ \text{Solve}(F|_{\ell \text{ false}}, \mathcal{P} \cup \{P_x\})\}$.

The numbers of true literals change whenever a variable is set, by branching, domination or propagation.

Proposition 20 (Soundness)**.** Solve *returns the optimum for every branching rule and every choice of the pruned literal, provided each branching adds a clause to only one of its two children.*

Proof. A complete assignment α determines a path from the root: at a branching node follow the child that agrees with α, and at a fixing step (domination or propagation) continue if α takes the fixed value and stop otherwise. Call the child without a new clause, and the fixed value, the *preferred* side. For $\alpha \neq \beta$ let w be the first node at which their paths part, and write $\alpha \succ \beta$ if α takes the preferred side at w. This is a strict total order on complete assignments.

Let α^* be the $\succ$-largest optimal assignment, and suppose it is never evaluated. Then it disagrees with a domination step, or it violates a pruning clause on its path; disobeying a propagation step is such a violation. Let w be the shallowest node on its path that is either a domination step disagreeing with α^* or the creation node of a clause that α^* violates. If w fixes v by domination, the proof of Lemma 1 shows that flipping v to the fixed value does not decrease the score. If w branches on x, then α^* lies in its pruned child, and Lemma 18, applied to the formula at w, shows the same for flipping x. The flipped assignment agrees with α^* on every variable decided above w, so it follows the same path to w and takes the preferred side there. It is optimal and larger than α^*, a contradiction. □

Remark 21 (Surviving leaves). By Lemma 18, at every surviving leaf each variable that was branched to its pruned side is strictly better than its flip, given all other values. The surviving leaves are one-sided local optima.

Observation 22 (Benchmark). On $C_{n,c}$ pruning clauses do not change the search tree: it still has $\binom{n-1}{a}$ leaves, for every branching rule.

Proof. At a node with z zeros and o ones assigned, all unassigned variables have deficits $(a', b') = (a - z, b - o)$. Let $a' \leq b'$; then $\ell = x$ and the pruned child is $x = 0$. For every neighbour y the pruning literal is $\neg y$, both for $(x \vee y)$ and for $(\neg x \vee \neg y)$, so P_x allows at most $a' - 1$ further zeros. In that child all other variables have deficits $(a' - 1, b')$. If $a' - 1$ further zeros occur, closure fixes all remaining variables to 1 and the bound holds; propagation only repeats these fixings. If instead b' further ones occur first, closure fixes all remaining variables to 0 and may exceed the bound, but at a node that is a leaf anyway. So the rule turns some leaves into cuts and leaves the tree unchanged. The case $a' > b'$ is symmetric. □

9 How much do pruning clauses prune?

9.1 The level budget with pruning clauses

Lemma 23. *Let $B = 1$, let π be a static order, let Q be a set of variables, and fix for each $x \in Q$ a literal ℓ_x in advance. Run the static rule A_π with domination and cuts, without propagation, where branching on $x \in Q$ adds the pruning clause of x with respect to ℓ_x to the ℓ_x-false child, and other branchings add nothing. For $\sigma \in \{0,1\}^{P_k}$ and $x \in Q \cap P_k$ let $\mathrm{prog}_x(\sigma)$ be the number of neighbours of x in P_k whose value under σ lowers p_{ℓ_x}. Then at most $|G_k|$ internal nodes branch on x_k, where*

$$G_k = \{\sigma \in \{0,1\}^{P_k} : \text{ for every } x \in Q \cap P_k,\ \sigma \text{ makes } \ell_x \text{ true or } \mathrm{prog}_x(\sigma) \leq p_{\ell_x}(F)\}.$$

Proof. As in Lemma 7, the values σ_u of P_k at the internal nodes u branching on x_k are pairwise distinct; cuts do not affect that argument. Fix such a u and $x \in Q \cap P_k$. If x was branched or fixed to make ℓ_x true, then σ_u makes ℓ_x true. If x was branched to ℓ_x false at a node w, its clause is active and not violated at u. The neighbours assigned before w lowered p_{ℓ_x} from $p_{\ell_x}(F)$ to $p_{\ell_x}(w)$; the neighbours assigned between w and u with their lowering value are true literals of the clause, at most $p_{\ell_x}(w) - 1$ of them. So at most $p_{\ell_x}(F) - 1$ neighbours of x have their lowering value at u, and those in P_k are among them. If x was

fixed by domination to make ℓ_x false at a node v, then $p_{\neg\ell_x}(v) \le 0$, hence $p_{\ell_x}(v) \ge D_x(v)$, the number of neighbours unassigned at v. The neighbours assigned before v lowered p_{ℓ_x} by at most $p_{\ell_x}(F) - D_x(v)$, and the later ones add at most $D_x(v)$ because $B = 1$. So $\mathrm{prog}_x(\sigma_u) \le p_{\ell_x}(F)$. □

Theorem 24 (Order independence)**.** *Let F be canonical with $B = 1$ on n variables, let $\varepsilon \in (0, 1]$, and let S be a set of variables with $|\Delta_v(F)| \ge \varepsilon n$. Let ℓ_v be the literal of v in its gap direction, so that $p_{\ell_v}(F) = (D_v - |\Delta_v|)/2$. Run the algorithm of Lemma 23 with $Q = S$ and these literals, for an* arbitrary *static order π. Then*

$$L \le 1 + n \cdot 2^{\,n-|S|} + n|S| \cdot 2^{\,H((1-\varepsilon)/2)\,n}.$$

Proof. By Lemma 23, $L \le 1 + \sum_k |G_k|$. Take σ uniform on P_k. If $\sigma \in G_k$, then either σ makes every ℓ_x with $x \in S \cap P_k$ true, which has probability $2^{-|S\cap P_k|}$, or $\mathrm{prog}_x(\sigma) \le p_{\ell_x}(F)$ for some such x. Hence

$$|G_k| \le 2^{\,|P_k \setminus S|} + \sum_{x \in S\cap P_k} 2^{k-1} \Pr\Big[\mathrm{Bin}(K_x, \tfrac{1}{2}) \le \tfrac{K_x + R_x - \Delta}{2}\Big],$$

where K_x and R_x are the numbers of neighbours of x inside and outside P_k, $\Delta = |\Delta_x(F)| \ge \varepsilon n$, and $B = 1$ makes $\mathrm{prog}_x(\sigma)$ binomial. The first term is at most $2^{n-|S|}$. Variables with $\Delta \ge D_x$ are dominated at the root and fixed toward ℓ_x, so they do not occur in the second term. Otherwise put $m = k - 1 \ge K_x$ and $g = \Delta - R_x < K_x$; since $R_x \le n - m$, we have $g \ge m - (1 - \varepsilon)n$. If $m < (1 - \varepsilon)n$, the summand is at most $2^m \le 2^{H((1-\varepsilon)/2)n}$. Otherwise the computation in the proof of Theorem 11 applies with this lower bound on g and bounds the summand by $2^{\psi(m)} \le 2^{nH((1-\varepsilon)/2)}$. Summing over the at most n levels and $|S|$ variables proves the bound. □

Theorem 11 needed the gap variables last, and Remark 15 indicates that the order matters for domination alone. With pruning clauses on the gap side the order no longer matters; the exponent is the same, and by Theorem 13 it cannot be improved by domination-type arguments. Adding further pruning clauses, for variables outside S, can only cut more nodes. The theorem is stated without propagation, which is sound (Proposition 20) but whose effect on the count we have not analyzed.

9.2 Disjoint supports: the savings multiply

Proposition 25. *Let $B = 1$. Let $x_1, \ldots, x_m$ be variables at pairwise distance at least 3 in the constraint graph, so that their closed neighbourhoods $N[x_i] = \{x_i\} \cup N(x_i)$ are pairwise disjoint. Let $D_i = |N(x_i)|$, let ℓ_i be the literal of x_i with the smaller deficit p_i, assume $p_i \ge 1$, and let R be the remaining variables. Branch in the order $x_1, N(x_1), \ldots, x_m, N(x_m), R$, with pruning clauses only at the x_i, with cuts and propagation, and without domination. Then the number of leaves is exactly*

$$2^n \prod_{i=1}^{m} \frac{1 + \theta_i}{2}, \qquad \theta_i = 2^{-D_i} \sum_{j < p_i} \binom{D_i}{j}.$$

Proof. When x_i is branched, none of its neighbours is assigned: they lie in its own block, and the clauses of earlier blocks only involve earlier blocks. So its deficit is p_i on every path, and its clause bounds the number of true pruning literals among its D_i neighbours by $p_i - 1$. In the ℓ_i-true child the block is branched freely, giving 2^{D_i} assignments of $N(x_i)$. In the ℓ_i-false child, propagation sets the rest of the block as soon as the count reaches $p_i - 1$, so no cut occurs, and every assignment of $N(x_i)$ with at most $p_i - 1$ true pruning literals is reached

exactly once: $\theta_i 2^{D_i}$ of them. When block i opens, the clauses of earlier blocks are decided, so every leaf of block i continues into an identical subtree. Multiplying, with $2^{|R|}$ for R and $n = |R| + \sum_i (1 + D_i)$, gives the formula. □

For a balanced centre $\theta_i \approx \frac{1}{2}$, so each centre saves about $\log_2 \frac{4}{3} \approx 0.4$ bits. Two ingredients make the count exact: (a) disjoint supports, so the survival events are independent and the factors multiply; and (b) each clause is decided right after it is created, because its support is the next block, so propagation leaves no dead ends.

Remark 26 (What the example does and does not show). The example isolates the mechanism; it is not an advantage of the rule. Here the classical order is better: branching on all non-centres first leaves every centre isolated, domination fixes it, and each centre costs a factor 1 instead of $1 + \theta_i$; this only needs the centres to be pairwise non-adjacent. In general, a single pruning clause carries exactly the information whether ℓ is dominated once all neighbours of x are assigned, which placing x after its neighbourhood provides directly. The rule's advantage is that it provides this for every branched variable at once, whereas an order provides it only for an independent set. Moreover, disjoint closed neighbourhoods allow at most $n/(d+1)$ centres in a graph of minimum degree d, so linearly many disjoint clauses force bounded degree.

9.3 Overlapping supports

Lemma 27 (Joint survival). *Let $\langle L_i \le b_i \rangle$, $i \in T$, be pruning clauses, let X_i be the number of true literals of L_i under a uniformly random assignment, and let $\lambda_i \ge 0$. For a variable z let A_z^1 (resp. A_z^0) be the sum of the λ_i over the clauses in which the literal of z is true for $z = 1$ (resp. $z = 0$). Then*

$$\Pr[X_i \le b_i \text{ for all } i \in T] \le e^{\sum_i \lambda_i b_i} \prod_z \frac{e^{-A_z^1} + e^{-A_z^0}}{2}.$$

Proof. On the event, $\sum_i \lambda_i (b_i - X_i) \ge 0$, so its indicator is at most $\exp\bigl(\sum_i \lambda_i (b_i - X_i)\bigr)$. Take expectations: $\sum_i \lambda_i X_i = \sum_z \bigl(A_z^1 [z = 1] + A_z^0 [z = 0]\bigr)$ is a sum of independent terms, one per variable. □

Each variable contributes one factor, which depends only on how its memberships split between its two values. With one clause per variable the factors are $(1 + e^{-\lambda})/2$, which is ordinary stacking. A variable shared with opposite literals contributes $(e^{-\lambda_1} + e^{-\lambda_2})/2$, less than two independent memberships: better than stacking. A variable shared with the same literal contributes $(e^{-\lambda_1 - \lambda_2} + 1)/2$, more: worse than stacking; identical clauses cut no more than one. For example, $\langle \{u, z\} \le 0 \rangle$ survives with probability $\frac{1}{4}$; together with $\langle \{w, z\} \le 0 \rangle$ both survive with probability $\frac{1}{8} > \frac{1}{16}$, and together with $\langle \{w, \neg z\} \le 0 \rangle$ with probability 0. In general, survival events that share only same literals are positively correlated and those that share only opposite literals are negatively correlated, by the Harris inequality after orienting each shared variable.

Corollary 28 (Signed Gram matrix). *Let M be the signed incidence matrix of the clauses: $M_{iz} = 1$ if the literal of z in L_i is true for $z = 1$, $M_{iz} = -1$ if it is true for $z = 0$, and $M_{iz} = 0$ otherwise; let $C = MM^{\mathsf{T}}$. Up to fourth-order terms in the λ_i, the logarithm of the bound in Lemma 27 exceeds its value for clauses with disjoint supports by*

$$\frac{1}{8} \sum_{i \ne j} \lambda_i \lambda_j C_{ij},$$

where C_{ij} is the number of shared variables with the same literal minus the number with opposite literals.

Proof. $\log \frac{e^{-a}+e^{-b}}{2} = -\frac{a+b}{2} + \log\cosh\frac{a-b}{2} = -\frac{a+b}{2} + \frac{(a-b)^2}{8} + O((a-b)^4)$, and $A^1_z - A^0_z = \sum_i \lambda_i M_{iz}$. Summing the squares over z gives $\sum_{i,j} \lambda_i\lambda_j C_{ij}$, and the diagonal part $\sum_i \lambda_i^2 C_{ii}$ is the value for disjoint supports. □

So the obstruction is not density but aligned density. Dense overlaps with random signs give $C_{ij} \approx 0$ and stack like independent clauses; opposite signs give $C_{ij} < 0$; aligned dense overlaps give large C_{ij}, equivalently large singular values of M. The rows of M are signed neighbourhoods: up to the sign chosen by ℓ, M is the block of the signed adjacency matrix A of Section 10 between branched and unassigned variables. $C_{n,c}$ is the extreme case: after orientation all rows of M are equal.

9.4 Timing

Remark 29 (Heuristic). Lemma 27 bounds the surviving *leaves*; the tree also contains the nodes visited before the clauses are decided. A clause can be violated only once the unassigned part of its support is smaller than the imbalance of its variable. In balanced instances the imbalance grows like the square root of the number of assigned neighbours, so with large overlapping supports no order decides clauses early. On random root-to-leaf walks (random dense balanced instances with edge probability $\delta = \frac{1}{2}$, random static orders), the first cut occurred with 24, 42, 72 and 113 variables still unassigned for $n = 50, 100, 200, 400$, close to the estimate $(|\Delta| + \sqrt{2\delta n \ln n})/\delta$, which gives 36, 54, 81 and 121. So the tree reaches depth about $n - O(\sqrt{n \log n})$ before anything is cut, however few leaves survive. For independent clauses the saving is roughly $n \cdot (\text{imbalance})/(\text{degree})$: linear for linear gaps, about $n/\sqrt{d}$ for balanced instances of degree d, and about $\sqrt{n}$ for dense ones.

10 A spectral view: domination for sets

In spin notation $s_v \in \{\pm 1\}$, a clause is a constant plus linear terms plus one coupling; for instance $(x \vee y)$ has value $\frac{3}{4} + \frac{s_x+s_y}{4} - \frac{s_x s_y}{4}$. With $B = 1$ the score is

$$\text{const} + \sum_v h_v s_v + \tfrac{1}{2} s^{\mathsf{T}} K s, \qquad h_v = \Delta_v/4, \qquad K = A/4,$$

where A is the signed adjacency matrix of the constraint graph: $A_{uv} = -1$ for a clause $(u \vee v)$ or $(\neg u \vee \neg v)$, and $A_{uv} = +1$ for a mixed clause. The field of a variable is a quarter of its effective gap, and its sign gives the majority literal, the literal with the smaller deficit.

Lemma 30 (Set domination)**.** *Fix a set T of t variables and assign all others. Let H_i be the fields on T in the resulting formula, $s^0 = \operatorname{sign}(H)$ the majority literals, and K_{TT} the couplings inside T.*

1. *If s^* is optimal on T and S is the set where it differs from s^0, then $\sum_{i\in S} |H_i| \le \|K_{TT}\|(\sqrt{|S|\,t} + |S|)$.*
2. *If every variable of a set $S' \subseteq T$ is set against its field and is strictly better than its flip, then $\sum_{i\in S'} H_i^2 < \|K_{TT}\|^2\, t$.*

Proof. (1) Write $s^* = s^0 - 2v$, where v agrees with s^0 on S and is 0 elsewhere. The linear part changes by $-2\sum_{i\in S}|H_i|$, and the quadratic part by $-2v^{\mathsf{T}}K_{TT}s^0 + 2v^{\mathsf{T}}K_{TT}v \le 2\|K_{TT}\|(\sqrt{|S|t} + |S|)$. Optimality of s^* gives the claim. (2) Flipping x changes the score by $-2s_x(H_x + (K_{TT}s)_x)$. Strictly worse means $s_x(H_x + (K_{TT}s)_x) > 0$; with $s_x = -\operatorname{sign}(H_x)$ this gives $s_x(K_{TT}s)_x > |H_x|$. Summing squares, $\sum_{x\in S'} H_x^2 < \|K_{TT}s\|^2 \le \|K_{TT}\|^2 t$. □

In clause terms, part (1) reads $\sum_{i\in S}|\Delta_i| \le \|A_T\|(\sqrt{|S|t}+|S|)$. Single-variable domination, and a single pruning clause, is the ℓ_1 version $|\Delta_i| \le D_i$: the worst case over the neighbours. The operator norm allows for sign cancellations; for random signs $\|A_T\|$ is of the order of the square root of the degree rather than the degree, so only few variables can go against their majority literal at the same time. Part (2) certifies that many pruning clauses cannot survive together (Remark 21): it is a joint use of clauses. The same matrix appears in both routes, since the rows of M in Corollary 28 are, up to sign, rows of a block of A; large singular values spoil the stacking of pruning clauses and the set bound alike.

Expected consequence (not yet proved). For pseudo-random dense instances, let T be the last εn variables of the order. Most fields are then of order $\sqrt{d}$ while $\|A_T\|$ is of order $\sqrt{\varepsilon d}$, so by part (1) the optimum on T agrees with the majority literals up to a small fraction of T, and the search on T can be replaced by an enumeration of $2^{o(1)\cdot t}$ candidates, for a total of $2^{(1-\Omega(\varepsilon))n}$. A complete proof needs a small-ball estimate over the outer assignments, which fails for near-twin variables, whose fields are small together.

10.1 Two special cases that separate the difficulties

1. **Pseudo-random dense balanced instances.** Single-variable rules, domination and single pruning clauses, act only in the last $O(\sqrt{n\log n})$ levels (Remark 29) and save only $2^{O(\sqrt{n\log n})}$. The set bound should give c^n. This case isolates the need for joint certificates.

2. **Aligned structure.** $C_{n,c}$ defeats every single-variable rule (Theorem 13 and Observation 22) and the set bound as well, since $A = -(J - I)$ has the eigenvalue $-(n-1)$ in the all-ones direction. Yet it is easy, by the next proposition. This case isolates the need to exploit low-rank structure.

Proposition 31. *If every pair of variables carries exactly one clause, of the form $(x \vee y)$ or $(\neg x \vee \neg y)$, then Max 2-SAT is solvable in polynomial time.*

Proof. Since $[x\vee y] = x+y-xy$ and $[\neg x\vee\neg y] = 1-xy$, the score equals $N^- + \sum_v d_v^+ x_v - \binom{m}{2}$, where N^- is the number of clauses $(\neg x \vee \neg y)$, d_v^+ the number of clauses $(v \vee y)$, and m the number of true variables. For each m, the best assignment makes true the m variables with the largest d_v^+; try all m. □

The general case is a mixture: aligned parts, which are low-rank and should be handled structurally, and pseudo-random parts, which need joint certificates.

11 Numerical checks

The scripts are in this folder.

- `explore_dpll_measures.py` (about three minutes): Lemma 7 on 240 random canonical instances with $n \in \{8, 10, 12\}$, $B \in \{1, 2\}$ and random orders, $L \le B_\pi$ always, with equality attained; Theorem 13 and Remark 14 on $C_{n,c}$ for $n \le 19$, where the natural order, maximum-degree branching and random orders give identical trees with $\binom{n-1}{a}$ leaves, equal to B_π, and for $n = 3201$ the exponents of Table 1 are within 0.003 of $H((1-\varepsilon)/2)$; Observation 16, with no violation on $C_{n,c}$ and 28 of 357 branchings violated on random aligned instances.

- `explore_extended_rule.py` (seconds): Proposition 20 on 240 random canonical instances with $n \leq 12$, nested pruning clauses, domination, random unit clauses and complementary pairs, where the optimum found with the rule, with and without propagation, equals the brute-force optimum; Observation 22, with identical leaf counts on $C_{n,c}$ for $n \leq 19$. On small random families the rule removes 7–27% of the leaves without propagation and 29–43% with it.
- `explore_disjoint_blocks.py`: Proposition 25, exact equality on 12 instances.
- `explore_late_detection.py`: the measurements of Remark 29.

12 Open questions and next steps

1. **Overlaps, quantitatively.** Turn Lemma 27 and Corollary 28 into a bound on the whole tree, under a condition on the signed Gram matrix (for example a bounded aligned degree), together with the timing.
2. **Early joint detection.** Certify that many active pruning clauses cannot all survive long before any single one is violated. Lemma 30(2) is one such certificate; pairs of clauses with opposite literals are another.
3. **Special case 1.** Turn the set bound into a pruning rule inside DPLL and prove c^n for random dense balanced instances.
4. **Special case 2.** Exploit aligned, low-rank structure beyond Proposition 31, and combine it with item 3.
5. **Smaller items.** Adaptive orders such as maximum degree with pruning clauses; the exact rate for $B = 2$; a lower bound matching the term $1 - \eta$.

References

[1] A literal-deficit measure for Max 2-SAT. Working note, 22 September 2026 (`max2sat_literal_deficit_proof.tex`).

[2] Exact Max 2-SAT with an absolute linear polarity gap. Working note, 23 September 2026 (`max2sat_absolute_linear_gap_proof.tex`).

[3] R. Williams. A new algorithm for optimal 2-constraint satisfaction and its implications. *Theoretical Computer Science* 348(2–3):357–365, 2005.

[4] G. J. Woeginger. Open problems around exact algorithms. *Discrete Applied Mathematics* 156(3):397–405, 2008.

[5] A. Golovnev. New upper bounds for MAX-2-SAT and MAX-2-CSP w.r.t. the average variable degree. In *IPEC 2011*, LNCS 7112, pp. 106–117, 2012.

[6] A. Golovnev and K. Kutzkov. New exact algorithms for the 2-constraint satisfaction problem. *Theoretical Computer Science* 526:18–27, 2014.

[7] M. B. Hastings. Local maxima and improved exact algorithm for MAX-2-SAT. arXiv:1610.07100, 2016.