

Corrected low-dimensional temperature-transfer results. Direct FM samples at the target β are compared with AM-corrected samples. $E(\cdot)W_2$ is the Wasserstein–2 distance between energy distributions; marginal W_2 is the mean coordinate-wise one-dimensional Wasserstein–2 distance. Lower is better.

System	Reference	FM anchors	AM pair	FM $E(\cdot)W_2$	AM $E(\cdot)W_2$	FM marginal W_2	AM marginal W_2
DW-1 [†]	Legacy	[0.25, 1.50]	0.25 $\rightarrow$ 0.70	0.604	0.184	0.256	0.049
DW-2 [†]	Legacy	[0.25, 1.50]	0.25 $\rightarrow$ 0.70	1.941	0.151	0.395	0.031
MB-2*	Reference-v2	[0.25, 0.50, 0.75, 1.00, 1.50]	1.00 $\rightarrow$ 1.20	0.549	0.066	0.123	0.015

[†]Hash-bound legacy results: energy W_2 uses the frozen 2k protocol and dataset-generation provenance is incomplete. *Corrected single-seed pilot using independent 100k train/eval reference-v2 samples; energy W_2 uses 20k samples. Its independent reference-vs-reference energy floor is 0.0284 ± 0.0034 . The former MB-2 0.75 $\rightarrow$ 0.90 row is omitted because it used the rejected finite-step ULA reference.