

UNIVERSITÀ DEGLI STUDI DI TORINO
DIPARTIMENTO DI MATEMATICA GIUSEPPE PEANO

SCUOLA DI SCIENZE DELLA NATURA

Corso di Laurea Magistrale in Matematica



Tesi di Laurea Magistrale

Consistency results concerning ω_1 -trees

Relatore: Prof. Matteo Viale
Correlatore: Prof. Sy David Friedman

Candidato: Andrea Gadotti

ANNO ACCADEMICO 2015/16

To my parents

Acknowledgements

I wish to thank Prof. Sy D. Friedman for advising this work and hosting me at the Kurt Gödel Research Center for the past months. I am very grateful for living in an academic, social and historical context which could give me this opportunity.

I need to thank Prof. Matteo Viale, who supported my wish to write the thesis in Vienna.

I sincerely thank Stefan Hoffelner for helping me promptly whenever I asked and for putting up with my mathematical immaturity.

Ringrazio la mia famiglia per la libertà che mi ha donato. Un pensiero particolare va ai miei genitori: le loro scelte e il loro affetto mi hanno assicurato il privilegio di pensare al mio futuro sempre con ottimismo e serenità.

Ringrazio i miei amici: per quanto distanti ci troviamo, il legame che ci unisce non si spezza mai.

Contents

Introduction	viii
Notation	x
Preliminaries	xi
1 The tree property	1
1.1 Basic notions about trees	1
1.2 The tree property at ω_1	2
1.3 The tree property at higher cardinals	4
2 Suslin trees	7
2.1 Suslin's Hypothesis	7
2.2 Suslin trees in L	9
2.3 Forcing and Suslin trees	14
2.3.1 A forcing model where there is a Suslin tree	14
2.3.2 A forcing model where there are no Suslin trees	16
3 Kurepa trees	21
3.1 Kurepa's Hypothesis	21
3.2 Kurepa families in L	22
3.3 Forcing and Kurepa trees	24
3.3.1 A forcing model where there is a Kurepa tree	24
3.3.2 A forcing model where there are no Kurepa trees	27
4 Automorphisms of trees	31
4.1 A homogeneous Suslin tree	33
4.2 A rigid Suslin tree	38
Bibliography	41

Introduction

Trees play an important role in set theory, mainly because of their rather *concrete* nature. This is probably the reason why many fundamental tools and concepts in set theory (most notably, iterated forcing) originated from the study of trees. The starting point can be identified with the following well-known elementary fact, known as König’s lemma:

Every infinite tree with finite levels has an infinite branch.

For an infinite cardinal κ , a κ -tree is a tree of height κ . The *tree property at κ* , or $\text{TP}(\kappa)$, asserts that

Every tree of height κ with levels of size $< \kappa$ has a branch of length κ .

Any counterexample to $\text{TP}(\kappa)$ is called *Aronszajn κ -tree*. So König’s lemma says that there exists no Aronszajn ω -tree, or equivalently that $\text{TP}(\omega)$ is true.

In 1935, Kurepa conducted the first systematic study of trees, which included the first construction of an Aronszajn tree of height ω_1 . In the same work, he introduced the notions of *Suslin tree* and *Kurepa tree*. A Suslin tree (ST) is an ω_1 -tree with no uncountable antichains and no uncountable branches. A Kurepa tree (KT) is an ω_1 -tree with countable levels and at least \aleph_2 -many uncountable branches. Kurepa’s hypothesis (KH) is the conjecture “There exists a KT”. Kurepa characterized in terms of Suslin trees a famous problem originally formulated by Suslin, called *Suslin hypothesis* (SH), which states:

Every dense linearly ordered set without endpoints which is Dedekind complete and satisfies the countable chain condition¹ is order-isomorphic to the real line.

Kurepa translated SH into a purely combinatorial problem, by showing that SH holds iff there exists no ST. However, he couldn’t prove nor refute the existence of Suslin and Kurepa trees. In fact, no much progress was made about this problem until the advent of forcing. But after Cohen established the relative consistency of $\neg\text{CH}$ introducing the method of forcing in 1963, many people began to explore the possibility of applying the same techniques to SH and KH. Within a few years, the problem was settled, as both SH and KH were proved to be independent of ZFC. This thesis offers a detailed presentation of such consistency results, together with other interesting theorems about trees.

¹The *countable chain condition* says that every collection of mutually disjoint non-empty open intervals is countable.

In the first chapter we present Kurepa’s construction of an Aronszajn ω_1 -tree. Then we generalize this result to Aronszajn trees of height the successor of any regular cardinal, under the assumption that GCH holds.

The second chapter is the core of our work. After discussing some easy properties of Suslin trees that hold in ZFC, we prove the independence of SH. This is done by presenting Jech’s forcing model for \neg SH and Solovay-Tennenbaum’s model for SH. We also include a fascinating result by Jensen, which establishes the existence of a ST in Gödel’s constructible universe.

The third chapter is symmetrical to the second, but relative to Kurepa’s hypothesis and Kurepa trees. We shall mention that to “kill” Kurepa trees via forcing, we will need to assume that inaccessible cardinals exist. In fact, \neg KH is equiconsistent with the existence of an inaccessible cardinal.

The last chapter contains two “bonus” results: we show that using the diamond principle (\diamond), one can construct a *homogeneous* ST (i.e. a ST where any two nodes of the same height can be swapped by some automorphism) and a *rigid* ST (i.e. a ST with no non-trivial automorphisms). These are two enthralling theorems which are also interesting from a historical point of view, although they are much less known compared to the other results presented in the thesis.

Our work is inspired by Jech’s review about trees [6]. We expand both the mathematical and the historical content of Jech’s article, where proofs are explicitly defined as “sketched” by the author. This thesis is meant to be a clear and exhaustive introduction to trees, and proofs are filled with details. A peculiarity of our work is the historical perspective we adopted: although modern tools and notation are used, the main theorems are presented and proved in their original form, whereas the standard approach in textbooks is to derive them from later results. More specifically, some technical *focal* principles were formulated to generalize and strengthen the results presented in our thesis. This is for example the case of the diamond principle, introduced by Jensen and extracted from his proof that $V = L$ implies the existence of a ST. Usually, in textbooks it is proved that \diamond holds in L , and then it is noted that \neg SH easily follows from \diamond . Similarly, Martin’s axiom was formulated and studied by Martin by generalizing the methods (i.e. iterated forcing) introduced for the first time in the original work of Solovay and Tennenbaum (never published) which produced a model where no ST exists. The consistency of SH is usually derived from the one of Martin’s axiom. While this approach totally makes sense, we believe that our exposition may result pleasant especially to students who saw only the application of forcing to CH, because of the good intuition that trees tend to offer.

Notation

We list here some elementary definitions with the only purpose to fix non-standard notation.

Let $(X, <)$ be a partially ordered set. If $A \subseteq X$, then

$$\downarrow A := \{y \in X \mid y < a \text{ for some } a \in A\}$$

and

$$\Downarrow A := A \cup \downarrow A = \{y \in X \mid y \leq a \text{ for some } a \in A\}.$$

If $x \in X$ then $\downarrow x := \downarrow\{x\}$ and $\Downarrow x := \Downarrow\{x\}$. Similarly for $\uparrow A, \Uparrow A, \uparrow x, \Uparrow x$. If we want to avoid confusion about the ordered set we are considering, we may denote $\downarrow x$ by $X \restriction x$.

Let $f: A \rightarrow B$ and let S be a set. We denote $\{f(s) \mid s \in S\}$ by $f[S]$ or $f''S$. Moreover, $\text{ran}(f)$ means $f[A]$.

Ord is the class of ordinals. Unless otherwise stated, small Greek letters always refer to ordinals.

If $(X, <)$ is a well-order, $\text{type}(X, <)$ is its order type.

If A, B are sets, AB denotes the set of the functions with domain A and codomain B . If λ, κ are cardinals, κ^λ is $|\kappa^\lambda|$.

Let α be a limit ordinal. We say that a sequence $\langle \alpha_\xi \mid \xi < \beta \rangle$, with β limit ordinal, is *cofinal in α* if it's strictly increasing (i.e. $\xi < \delta \Rightarrow \alpha_\xi < \alpha_\delta$) and $\sup_{\xi < \beta} \alpha_\xi = \alpha$.

Let (X, \triangleleft) be a linearly ordered set. The *lexicographic order* on ${}^\omega X$ is defined by

$$f <_{\text{lex}} g \Leftrightarrow \exists n \in \omega [f(n) \triangleleft g(n) \text{ and } \forall i < n (f(i) = g(i))].$$

Preliminaries

In this work, we assume that the reader is familiar with concepts and results which are usually covered in graduate set theory courses, in particular: ordinals, transfinite induction and recursion, the constructible hierarchy and forcing. Such topics are extensively described in, e.g., [8] and [14]. For convenience, in this chapter we will very briefly summarize some basic results about (relative) constructibility and forcing.

Constructibility

If M is a set, we say that $Z \subseteq M$ is *definable over* (M, \in) if $Z \subseteq M$ and for some formula $\varphi(v_0, \dots, v_n)$ and some $x_1, \dots, x_n \in M$ we have $Z = \{z \in M \mid (M, \in) \models \varphi(z, x_1, \dots, x_n)\}$. Then we let

$$\text{def}(M) := \{Z \subseteq M : Z \text{ is definable over } (M, \in)\},$$

The *constructible hierarchy* $\langle L_\alpha \mid \alpha \in \text{Ord} \rangle$ is defined by recursion:

$$\begin{aligned} L_0 &= \emptyset; \\ L_{\alpha+1} &= \text{def}(L_\alpha); \\ L_\eta &= \bigcup_{\beta < \eta} L_\beta \quad \text{if } \eta \text{ is a limit ordinal} \end{aligned}$$

The *constructible universe* is the class $L := \bigcup_{\alpha \in \text{Ord}} L_\alpha$. L is a model of $\text{ZFC} + \text{GCH} + V = L$. L can be well-ordered (we denote the canonical well-order by $<_L$).

We now generalize this definition. Let A be a set. Then we let

$$\text{def}_A(M) := \{X \subseteq M : X \text{ is definable over } (M, \in, A \cap M)\},$$

where $A \cap M$ is considered as a unary predicate and means “ $x \in A \cap M$ ”. If we use def_A in place of def in the definition above, we obtain $\langle L_\alpha[A] \mid \alpha \in \text{Ord} \rangle$, which is called the *hierarchy of sets constructible from A* (or *relative to A*). Then the *universe constructible from A* is $L[A] := \bigcup_{\alpha \in \text{Ord}} L_\alpha[A]$.

We list some well-known facts about relative constructibility.²

- i) For every set A and $\alpha \in \text{Ord}$, $L_\alpha[A] \cap \text{Ord} = \alpha$.

²For a more complete list, see [3], pp. 102 and following.

- ii) Let A be a set and let $B := A \cap L[A]$. Then $B \in L[A]$, $L[A] = L[B]$ and $(V = L[B])^{L[A]}$. In particular, by the first point $A \cap L[A] = A$ if $A \subseteq \text{Ord}$. So we always have $A \in L[A]$ and $L[A] \models V = L[A]$ in this case.
- iii) $L[A]$ is the smallest inner model of **ZF** which contains the set $A \cap L[A]$ (i.e. $L[A]$ is the smallest inner model M of **ZF** such that $A \cap M \in M$).

The following is a very important result due to Gödel and generalized for relative constructibility:

Condensation Lemma. Let A be a set and $\alpha > \omega$ a limit ordinal. Suppose $M \preceq L_\alpha[A]$ and let mos be the collapsing function of (M, \in) . Then $\text{mos}''M = L_\beta[\text{mos}''A]$ for some $\beta \leq \alpha$.

Forcing

Forcing is an extremely powerful method for producing models of set theory. The construction of this tool is rather involved, and even a general overview would require several pages. Therefore we will present here just the central results, omitting also the definitions of most objects.

Let M be a countable transitive model of **ZFC** and let $(P, \leq, 1_P)$ be a forcing poset in M , where $1_P \in P$ is a largest element of P (i.e. $\forall p \in P [p \leq 1_P]$). Let G be generic for P over M .

The Generic Model Theorem. There exists a transitive model $M[G]$ such that

- (i) $M[G] \models \text{ZFC}$;
- (ii) $M \subseteq M[G]$ and $G \in M[G]$;
- (iii) $\text{Ord}^{M[G]} = \text{Ord}^M$;
- (iv) if N is a transitive model of **ZF** such that $M \subseteq N$ and $G \in N$, then $M[G] \subseteq N$.

The model $M[G]$ is called *generic extension* of M . Every element of $M[G]$ has a *P-name* in M . The set of names is denoted by M^P . We can define in M the *forcing relation* \Vdash and the *forcing language*. \Vdash is a relation between elements of P and statements of the forcing language. The forcing language contains every P -name. If $a \in M[G]$, we usually denote with \dot{a} a name in M for a . If a belongs also to M , then it has a *canonical name* which we denote by \check{a} . Among the names, the name \dot{G} is special, because it is defined in such a way that its interpretation relative to any filter H is H itself. Moreover, $p \Vdash [\check{p} \in \dot{G}]$, for any $p \in P$. If $1_P \Vdash \sigma$, then sometimes we simply write $\Vdash \sigma$. The following is the tool which enables us to control the truth in $M[G]$:

The Forcing Theorem. Let σ be a sentence of the forcing language. Then

$$M[G] \models \sigma \text{ if and only if } \exists p \in G [p \Vdash \sigma].$$

An important property of the forcing relation is the following:

Maximal Principle. Let $\varphi(x)$ be a formula of the forcing language with no free variable other than x . If $p \Vdash \exists x \varphi(x)$, then for some $\dot{a} \in M^P$, $p \Vdash \varphi(\dot{a})$.

In applications of forcing, it is of fundamental importance to determine the way ordinals act as cardinals in the generic extension. Let $o(M)$ denote the least ordinal not in M . Suppose $(\theta \text{ is a cardinal})^M$. We say that P *preserve cardinals* $\geq \theta$ if, for any G generic for P over M and for all $\theta \leq \beta < o(M)$,

$$(\beta \text{ is a cardinal})^M \text{ iff } (\beta \text{ is a cardinal})^{M[G]}.$$

We say that P *preserve cofinalities* $\geq \theta$ if, for any G generic for P over M and for all limit $\gamma < o(M)$ such that $\text{cf}^M(\gamma) \geq \theta$,

$$\text{cf}^M(\gamma) = \text{cf}^{M[G]}(\gamma).$$

Of course preserving cofinalities is a stronger condition than preserving cardinals.

For a cardinal θ , P has the θ *chain condition* (θ -cc) if every antichain in P has size less than θ . If $\theta = \omega_1$, we say *countable chain condition* (ccc). The following is a central result:

Fact. If $(\theta \text{ is a regular cardinal})^M$ and $(P \text{ is } \theta\text{-cc})^M$, then P preserve cofinalities $\geq \theta$ and hence P preserves cardinals $\geq \theta$.

Let G be generic for P over M . It's immediate to check that if P preserves all cofinalities, then $(\aleph_\alpha)^M = (\aleph_\alpha)^{M[G]}$ for all $\alpha < o(M)$.

When the chain condition doesn't hold or holds only above a certain cardinal, there is another way to check if smaller cardinals are preserved. We say that P is λ -*closed* if whenever $\delta < \lambda$ and $\langle p_\xi : \xi < \delta \rangle$ is a (weakly) decreasing sequence in P , then there is a $q \in P$ such that $q \leq p_\xi$ for all $\xi < \delta$. If $\lambda = \omega_1$, then we say *countably closed*.

Fact. Assume that $(P \text{ is } \lambda\text{-closed})^M$ and $A, E \in M$ with $(|A| < \lambda)^M$. If $f: A \rightarrow E$ and $f \in M[G]$, then $f \in M$.

This means that if P is λ -closed and $\omega < \lambda < o(M)$, then ${}^\delta\lambda \cap M = {}^\delta\lambda \cap M[G]$ for all ordinals $\delta < \lambda$. It follows easily that $\text{cf}^M(\gamma) = \text{cf}^{M[G]}(\gamma)$ for all limit $\gamma \leq \lambda$. So, in particular, countably closed posets don't add ω -sequences (whose ranges are in M) and preserve \aleph_1 .

We now illustrate the core theorem for product forcing.

Product lemma. Let P and Q be forcing posets in M . $P \times Q$ is the poset whose order is defined by $(p_1, q_1) \leq (p_2, q_2) \Leftrightarrow p_1 \leq p_2$ and $q_1 \leq q_2$. Let $\pi_1: P \times Q \rightarrow P$ and $\pi_2: P \times Q \rightarrow Q$ be the projections. The following are equivalent:

- (i) $G \subseteq P \times Q$ is generic over M .
- (ii) $G = G_1 \times G_2$, where $G_1 = \pi_1[G]$ is P -generic over M and $G_2 = \pi_2[G]$ is Q -generic over $M[G_1]$.

Moreover, $M[G] = M[G_1][G_2]$.

A significant generalization of product forcing is iterated forcing. In this work we will use finite support iterations. The definition of this tool is quite technical, so we refer to [14] for an in-depth presentation and for the notation. Here we just mention the important theorem which makes iterated forcing a robust method:

Fact. Let P_α be the iteration with finite support of $\langle \dot{Q}_\beta : \beta < \alpha \rangle$. Suppose that for each $\beta < \alpha$, $\Vdash_\beta [\dot{Q}_\beta \text{ is ccc}]$. Then P_α is ccc.

We conclude this list of preliminaries with a quick description of a special type of names, which are perhaps less standard compared to the concepts presented so far.

Definition. Let $\tau \in M^P$ be a name. A *nice name for a subset of τ* is a name of the form

$$\bigcup \{ \{ \sigma \} \times A_\sigma : \sigma \in \text{dom}(\tau) \},$$

where each A_σ is an antichain of P in M .

Next lemma shows that we can in fact use nice names to name subsets of a given name.

Fact. If $\tau, \mu \in M^P$, then there is a nice name $\theta \in M^P$ for a subset of τ such that $1 \Vdash (\mu \subseteq \tau \Rightarrow \mu = \theta)$.

Nice names relative to ccc posets are useful because it's easy to bound their size:

Fact. Fix $\tau \in M^P$. Let $\kappa := |P|$ and $\lambda := |\text{dom}(\tau)|$. Assume that P is ccc and that κ and λ are infinite. Then the set of nice names for subsets of τ has size at most κ^λ . (all the statements here are meant to hold in M)

On a final note, we remark that our overview about forcing above takes place in the “forcing over a countable transitive model” setting. Of course everything can be adapted to the “forcing over V ” approach. In this work, sometimes we force over a countable transitive model and sometimes we force over V . We assume that the reader is familiar with both the approaches.

Chapter 1

The tree property

1.1 Basic notions about trees

In this section we define trees and several basic concepts related to them.

Definition 1.1. A *tree* is a partially ordered set $(T, <)$ such that for all $x \in T$ the set $\downarrow x$ is well-ordered by $<$. The elements of T are called *nodes*. Moreover:

- If $x \in T$, the *order* of x is $o(x) := \text{type}(\downarrow x)$.
- The α th level of T is $U_\alpha := \{x \in T \mid o(x) = \alpha\}$.
- The *height* of T is the least ordinal such that every $x \in T$ has smaller order type, i.e. $\text{height}(T) := \sup\{o(x) + 1 \mid x \in T\}$.
- A *chain* in T is a linearly ordered subset of T . A *branch* is a maximal chain. If \mathfrak{b} is a branch in T , of course we can define $\text{height}(\mathfrak{b}) := \text{type}(\mathfrak{b})$.
- A subset $S \subseteq T$ is *cofinal* in T if for all $\alpha < \text{height}(T)$ there is $x \in S$ such that $o(x) \geq \alpha$.
- An α -tree is a tree of height α , and similarly for an α -branch.
- $T|_\alpha$ is the subset of T which contains every element of order strictly less than α , i.e. $T|_\alpha := \bigcup_{\xi < \alpha} U_\xi$. Obviously $T|_\alpha$ has height α if $\alpha \leq \text{height}(T)$.
- We say that a tree $(T_2, <_2)$ is an *extension* of $(T_1, <_1)$ if $<_1 = <_2 \cap (T_1 \times T_1)$, an *end-extension* if $T_1 = T_2|_\alpha$ for some α .

Example 1.2. We consider the family of trees given by all T which satisfy the following properties: for some $\alpha < \omega_1$,

- (i) every element $t \in T$ is a function $t: \beta \rightarrow \omega$, with $\beta < \alpha$;
- (ii) T is closed under initial segments, i.e. if $t \in T$ then $t \upharpoonright \beta$ is in T as well, for any β ;

- (iii) if $t: \beta \rightarrow \omega$ is in T and $\beta + 1 < \alpha$, then $t \frown n \in T$ for all $n \in \omega$;
- (iv) if $t: \beta \rightarrow \omega$ is in T and $\beta \leq \gamma < \alpha$, then there exists $s: \gamma \rightarrow \omega$ such that $t \subseteq s$;
- (v) $T \cap {}^\beta \omega$ is countable for all $\beta < \alpha$.

Observe that T is a countable set and the β th level consists precisely of the functions in T whose length is β .

Definition 1.3. Let $\alpha \leq \omega_1$. An α -tree T is *normal* if.

- (i) T has a unique least point (which we call *root*);
- (ii) every level of T is at most countable;
- (iii) if x is not maximal in T , then there are infinitely many $y \geq x$ at level $o(x) + 1$ (we call these *immediate successors of x*);
- (iv) if $x \in T$ then there is $y > x$ at each higher level less than α ;
- (v) the order $<$ is extensional within each level U_γ such that $\gamma < \alpha$ is a limit ordinal, that is: for all $x, y \in U_\gamma$, if $\downarrow x = \downarrow y$ then $x = y$.

It is very easy to check that the trees of last example are normal. We shall use them as forcing conditions later because of these nice properties they enjoy.

We conclude this section with a trivial remark: there exist normal ω_1 -trees with uncountable branches, like the following:

Example 1.4. Consider $T := \{t \in {}^{<\omega_1}\omega \mid t \text{ has finitely-many 0's}\}$. To see that T is a normal ω_1 -tree is immediate, and $\mathfrak{b} := \{1^\alpha : \alpha < \omega_1\}$ is clearly an ω_1 -branch in T .

1.2 The tree property at ω_1

We start with an easy and well-known fact:

1.5. König's lemma. *If T is an ω -tree whose levels are all finite, then T has an ω -branch.*

Proof. Define $T' := \{x \in T \mid \uparrow x \text{ is infinite}\}$. It is immediate to construct an ω -branch in T' by recursion. Such branch is trivially an ω -branch in T . \square

Does König's lemma hold for cardinals greater than ω ? More precisely, we say that a cardinal κ has *the tree property*, in symbols $\text{TP}(\kappa)$, if the following statement is true:

If T is a κ -tree and if every level has cardinality $< \kappa$, then T has a κ -branch.

Of course $\text{TP}(\kappa)$ is false if κ is singular: if $\langle \alpha_\xi \mid \xi < \lambda \rangle$ is a cofinal sequence in κ with $\lambda < \kappa$, then take the tree given by the disjoint union of branches of length α_ξ for all $\xi < \lambda$, where elements of two different branches are incomparable.

We will show now that the tree property fails already at ω_1 . First, a general definition:

Definition 1.6. Let κ be a cardinal. An *Aronszajn κ -tree* is a κ -tree whose levels are of power less than κ but has no κ -branch.

Thus, there exists an Aronszajn κ -tree if and only if $\text{TP}(\kappa)$ is false. Aronszajn trees are named for Nachman Aronszajn, who established their existence for the ω_1 case in 1934. Actually, his proof appeared only in 1935 in Kurepa's thesis [15]. In his work, Kurepa provided also a different proof, producing the Aronszajn ω_1 -tree as a subtree of the tree given by the non-empty bounded well-ordered sets of rationals under end-extension¹. We will now present such proof.

Theorem 1.7. *There is an Aronszajn ω_1 -tree.*

Proof. We will construct the tree T in such a way that

- every $x \in T$ is a bounded and strictly increasing sequence of rational numbers;
- the order on T is defined by: $x \leq y$ iff y extends x , i.e. $x \subseteq y$;
- T is closed under initial segments.

By last condition, the α th level will consist precisely of the sequences of length α of T . Of course such a tree can't have an uncountable branch, since its union would yield a strictly increasing (and thus injective) sequence of length ω_1 into \mathbb{Q} , which is countable. Note that T must be constructed carefully: if we let any sequence be in T , then the ω th level would be uncountable already.

We construct T by induction on levels. To make sure that everything works, we will need to preserve the following properties (inductive hypotheses) at each level $\alpha < \omega_1$:

$$(1.1) \quad |U_\alpha| \leq \aleph_0;$$

$$(1.2) \quad \text{For all } \beta < \alpha, x \in U_\beta \text{ and } q > \sup x, \text{ there is } y \in U_\alpha \text{ such that } x \subseteq y \text{ and } q \geq \sup y.$$

Define $U_0 := \{\emptyset\}$. For the successor step, suppose that we have already constructed level U_α . Then we define

$$U_{\alpha+1} := \{x \frown r \mid x \in U_\alpha, r \in \mathbb{Q} \text{ with } r > \sup x\}.$$

It's easy to check that also $U_{\alpha+1}$ satisfies (1.1) and (1.2) w.r.t. $\alpha + 1$ (but note that one needs that \mathbb{Q} is dense).

For the limit step, let α be a limit ordinal and suppose we have already defined U_β for all $\beta < \alpha$.

Claim: For each $x \in T|_\alpha$ and each $q > \sup x$ there exists a strictly increasing α -sequence of rationals y such that y extends x , $q \geq \sup y$ and $y \restriction \beta \in T|_\beta$ for all $\beta < \alpha$.

Proof: Since $\alpha < \omega_1$, its cofinality is ω . Let $\langle \alpha_n \mid n \in \omega \rangle$ be cofinal in α and such that $x \in U_{\alpha_0}$. Now let $\langle q_n \mid n \in \omega \rangle$ be a strictly increasing sequence of rationals such that

¹Such tree is usually denoted by $\sigma\mathbb{Q}$.

$q_0 = \sup x$ and $\lim_n q_n \leq q$. Set $y_0 := x$. Using the inductive hypothesis (1.2) at each step, we can recursively find for each $n \geq 1$ a sequence $y_n \in U_{\alpha_n}$ which extends y_{n-1} and such that $\sup y_n \leq q_n$. By defining $y := \cup_n y_n$ we are done. \blacksquare

For all $x \in T|_\alpha$ and all $q > \sup x$ we choose an y as provided by the claim, and we define U_α as the set of all such y 's. It's clear that (1.2) holds for U_α . Because \mathbb{Q} and $T|_\alpha = \bigcup_{\beta < \alpha} U_\beta$ are countable, also (1.1) is preserved.

Of course T is an Aronszajn ω_1 -tree by construction. \square

1.3 The tree property at higher cardinals

We will now present a generalization of Theorem 1.7 due to Specker [18]:

Theorem 1.8. *Let κ be an infinite cardinal. If $\kappa^{<\kappa} = \kappa$, then there exists an Aronszajn κ^+ -tree.*

First we need two easy lemmas.

Lemma 1.9. *Let α be a limit ordinal. There exists a sequence $\langle \alpha_\xi \mid \xi < \text{cf}(\alpha) \rangle$ cofinal in α which is also continuous, i.e. $\alpha_\gamma = \sup_{\xi < \gamma} \alpha_\xi$ for all $\gamma < \text{cf}(\alpha)$ limit.*

Proof. Let $\langle \beta_\xi \mid \xi < \text{cf}(\alpha) \rangle$ be cofinal in α . Define $\langle \alpha_\xi \mid \xi < \text{cf}(\alpha) \rangle$ by

$$\alpha_\xi := \begin{cases} \beta_\xi, & \text{if } \xi \text{ successor} \\ \cup_{\eta < \xi} \beta_\eta, & \text{if } \xi \text{ limit.} \end{cases}$$

Of course this sequence is continuous and still cofinal in α . \square

Lemma 1.10. *Let κ be an infinite cardinal. Let $\mathcal{Q} := \{f \in {}^\omega \kappa \mid f(n) \neq 0 \text{ for finitely many } n \in \omega\}$. Then every $\alpha \leq \kappa$ embeds in any non-trivial open interval of \mathcal{Q} , ordered lexicographically.*

Proof. Let $f, g \in \mathcal{Q}$ be sequences with $f < g$. Let n be the least such that $f(n) < g(n)$ and let $m > n$ be such that $f(m) = 0$. For any $\alpha \leq \kappa$, define $\varphi: \alpha \rightarrow \mathcal{Q}$ given by

$$\varphi(\beta)(i) := \begin{cases} f(i) & \text{if } i \neq m; \\ \beta + 1 & \text{if } i = m. \end{cases}$$

Clearly φ is an order-embedding of α and $\text{ran } \varphi$ is contained in the interval (f, g) . \square

We can finally proceed with the

Proof of Theorem 1.8. We will adapt the proof of Theorem 1.7. Instead of \mathbb{Q} , we shall use \mathcal{Q} of Lemma 1.10. The only properties of \mathcal{Q} we will need are that $|\mathcal{Q}| = \kappa$, a well-known fact, and the statement of Lemma 1.10. Every $x \in T$ will be a bounded and strictly increasing sequence of elements of \mathcal{Q} such that $\text{length}(x) = \alpha$ for $\alpha < \kappa^+$. As before, T will be such that $o(x) = \text{length}(x)$ for all $x \in T$.

Again, we construct T by induction on levels, preserving for every $\alpha < \kappa^+$ conditions (1.1) (of course now we require $|U_\alpha| \leq \kappa$) and (1.2)², plus the following additional condition:

(1.3) If α is limit with $\text{cf}(\alpha) < \kappa$ and \mathfrak{b} is a branch in $T|\alpha$, then $\bigcup \mathfrak{b} \in U_\alpha$.

$U_0 := \{\emptyset\}$ and the successor step are just as before:

$$U_{\alpha+1} := \{x \smallfrown q \mid x \in U_\alpha, q \in \mathcal{Q} \text{ with } q > \sup x\},$$

which satisfies (1.1) and (1.2).

For U_α with α limit, we have again the claim:

Claim: For each $x \in T|\alpha$ and each $q > \sup x$ there is a strictly increasing α -sequence y in \mathcal{Q} such that y extends x , $q \geq \sup y$ and $y \restriction \beta \in T|\alpha$ for all $\beta < \alpha$.

Proof: Let $\lambda := \text{cf}(\alpha) \leq \kappa$. By Lemma 1.10 there exists $\langle q_\xi \mid 1 \leq \xi < \lambda \rangle$ strictly increasing and contained in the interval $(\sup x, q)$ of \mathcal{Q} ³. By Lemma 1.9, let $\langle \alpha_\xi \mid \xi < \lambda \rangle$ be cofinal in α , continuous and such that $x \in U_{\alpha_0}$.

As before, we want to recursively define $\langle y_\xi \mid \xi < \lambda \rangle$ such that for all $\xi < \lambda$ the following hold:

- (i) $y_\xi \in U_{\alpha_\xi}$;
- (ii) if $\eta < \xi$ then $y_\eta \subseteq y_\xi$;
- (iii) $\sup y_\xi \leq q_\xi$ (for $\xi \geq 1$).

Let $y_0 := x$. Suppose we have already defined y_ξ . Then there exists $y_{\xi+1}$ which satisfies our requests by the inductive hypothesis (1.2), just as in Theorem 1.7. The limit case is where we use the additional condition (1.3). Suppose $\xi < \lambda$ is a limit ordinal. First observe that $\text{cf}(\alpha_\xi) \leq \xi < \lambda \leq \kappa$ because we assumed $\langle \alpha_\xi \rangle_{\xi < \lambda}$ continuous. Now suppose we have already defined y_γ for every $\gamma < \xi$. Then it's clear that $y := \bigcup_{\gamma < \xi} y_\gamma$ satisfies (ii) and (iii). Condition (i) is also true, because $\sup_{\gamma < \xi} \alpha_\gamma = \alpha_\xi$ by continuity again, and thus $\langle y_\gamma \rangle_{\gamma < \xi}$ induces a branch in $T|\alpha_\xi$. So $y_\xi \in U_{\alpha_\xi}$ by hypothesis (1.3).

By defining $y := \bigcup_{\xi < \lambda} y_\xi$ we are done. ■

Now, suppose α is limit and $\text{cf}(\alpha) = \kappa$. For all $x \in T|\alpha$ and all $q > \sup x$ we choose an y as provided by the claim, and we define U_α as the set of all such y 's. It's clear that (1.1) and (1.2) hold for U_α .

The only case left is α limit with $\text{cf}(\alpha) < \kappa$. Then we define $U_\alpha := \{\bigcup \mathfrak{b} \mid \mathfrak{b} \text{ is a branch in } T|\alpha\}$, so that condition (1.3) certainly holds. By the claim, also (1.2) is true, since “ $y \restriction \beta \in T|\alpha$ for all $\beta < \alpha$ ” means precisely that $\mathfrak{b} := \{y \restriction \beta : \beta < \alpha\}$ is a branch in $T|\alpha$, so $y = \bigcup \mathfrak{b} \in U_\alpha$ by definition. Finally, observe that $T|\alpha = \bigcup_{\beta < \alpha} U_\beta$, so $|T|\alpha| \leq \kappa$. Hence

$$|U_\alpha| \leq |\{\mathfrak{b} : \mathfrak{b} \text{ is a branch in } T|\alpha\}| \leq |\alpha|^\kappa.$$

²Formally, the supremum here lives in the Dedekind completion of \mathcal{Q} .

³Actually, $\sup x$ might not be in \mathcal{Q} , but in that case we can simply take $q' \in \mathcal{Q}$ such that $\sup x < q' < q$ and consider the interval (q', q) .

But of course every branch in $T|_\alpha$ is completely determined by $\text{cf}(\alpha)$ -many entries, therefore $|U_\alpha| \leq \kappa^{\text{cf}(\alpha)}$. Since $\text{cf}(\alpha) < \kappa$ and by hypothesis $\kappa^{<\kappa} = \kappa$, we obtain that $|U_\alpha| \leq \kappa$, i.e. also condition (1.3) is satisfied.

Clearly T is an Aronszajn κ^+ -tree by construction. \square

Last theorem is totally useless for κ singular, since in that case the hypothesis is always false: if $\text{cf}(\kappa) < \kappa$ then $\kappa^{<\kappa} = \sup\{\kappa^\lambda \mid \lambda < \kappa, \lambda \text{ cardinal}\} \geq \kappa^{\text{cf}(\kappa)}$. But $\text{cf}(\kappa^{\text{cf}(\kappa)}) > \text{cf}(\kappa)$ by König's theorem, so $\kappa^{\text{cf}(\kappa)} > \kappa$ and hence $\kappa^{<\kappa} > \kappa$. Nonetheless:

Proposition 1.11. *Let κ be a regular cardinal. Suppose that GCH holds. Then $\kappa^{<\kappa} = \kappa$.*

Proof. The following is a well-known fact under GCH (see [13]):

Let $\kappa, \lambda \geq 1$ be cardinals with $\max(\kappa, \lambda)$ infinite. Then $\kappa^\lambda = \kappa$ if $\lambda < \text{cf}(\kappa)$.

So $\kappa^{<\kappa} = \sup\{\kappa^\lambda \mid \lambda < \kappa = \text{cf}(\kappa), \lambda \text{ cardinal}\} = \kappa$. \square

Hence, if we assume GCH we have that for every κ regular there exists an Aronszajn κ^+ -tree. What happens at the successor of a singular cardinal (even assuming GCH), or simply dropping GCH, is still being investigated. However, this topic is beyond the scope of this work.

Chapter 2

Suslin trees

Definition 2.1. Let (P, \leq) be a partially ordered set. An *antichain* in P is a subset $A \subseteq P$ such that any two distinct elements of A are incomparable, i.e. $x, y \in A$ and $x \neq y$ implies $x \not\leq y$ and $y \not\leq x$. A *maximal antichain* in P is an antichain which is maximal in P w.r.t. the inclusion relation between subsets of P .

Definition 2.2. A *Suslin tree* is an ω_1 -tree such that every branch is at most countable and every antichain is at most countable.

Of course any Suslin tree is an Aronszajn tree. Furthermore, observe that if an ω_1 -tree T is normal and has no uncountable antichain, then T is a Suslin tree. For, suppose by contradiction that \mathfrak{b} is an ω_1 branch. For all $x \in \mathfrak{b}$, let $f(x)$ be an arbitrary node greater than x which is not in \mathfrak{b} . Then $\{f(x) : x \in \mathfrak{b}\}$ would be an uncountable antichain.

2.1 Suslin's Hypothesis

The notion of Suslin tree was originally introduced in the attempt of improving a characterization of the real line proved by Cantor (1895):

Let R be a linearly ordered set without endpoints. Suppose it is dense, Dedekind complete and separable. Then R is order-isomorphic to the real line.

Suslin's hypothesis (SH), formulated by Suslin in 1920, states that the separability condition can be weakened to the *countable chain condition* (ccc): every collection of mutually disjoint non-empty open intervals is countable.

One can see that for a dense linear order, deleting endpoints and taking the Dedekind completion doesn't change the ccc or separability properties. Hence, Suslin's hypothesis can be stated as:

There exists no Suslin line

where a *Suslin line* is a dense linearly ordered set that satisfies the countable chain condition but is not separable.

SH is actually independent of ZFC. This important result was one of the earliest applications of forcing. But before the advent of forcing, not much progress concerning SH was made, except for a result proved by Kurepa [15] in 1935, who came up with the purely combinatorial concept of Suslin trees and reformulated SH in terms of them: SH is true if and only if there is no Suslin tree. We dedicate the last part of the section to the proof of this fact.

Lemma 2.3. *If there exists a Suslin tree then there exists a normal Suslin tree.*

Proof. Let T be a Suslin tree. We turn T into a normal Suslin tree through progressive transformations. In the end every property of Definition 1.3 will be satisfied. First, define $T_1 := \{x \in T : |\uparrow x| > \aleph_0\}$. Observe that if $x \in T_1$ and $\alpha > o(x)$, then there is $y \in T$ above x which is at level α and such that $|\uparrow y| > \aleph_0$. So T_1 satisfies condition (iv). For every set of the form $C = \downarrow y$ for some $y \in T_1$ at limit level, we add an extra node a_C in such a way that a_C is the least node above each element of C , i.e. $C < a_C$ and $a_C < x$ for all $x > C$. We call the obtained tree T_2 . Of course any new level is still countable, so T_2 satisfies (ii), (iv) and (v). Call a node *branching* if it has at least two immediate successor. Using that T_2 has no uncountable chain and satisfies (v), it's easy to see that for each $x \in T_2$, the set $\uparrow x$ contains uncountably many branching points. So $T_3 := \{\text{branching points of } T_2\}$ satisfies (ii), (iv) and (v). Also, each $x \in T_3$ is a branching point. Now let $T_4 := \{x \in T_3 : x \text{ belongs to a limit level of } T_3\}$, where the order in T_4 is just the restriction of the one in T_3 . It's easy to verify that T_4 is an ω_1 -tree which satisfies (ii), (iii), (iv), (v). Finally, to get a normal Suslin tree we just “glue” together the least nodes of T_4 . \square

Theorem 2.4. *There exists a Suslin line if and only if there exists a Suslin tree.*

Proof. Suppose we have a Suslin line S . We want to build a Suslin tree T by taking a certain family of closed nonempty intervals on S , ordered by reverse inclusion. The construction of T proceeds by induction on $\alpha < \omega_1$. Let $I_0 := [a_0, b_0]$ be arbitrary (with $a_0 < b_0$). Assume we already defined $I_\beta = [a_\beta, b_\beta]$ for all $\beta < \alpha$. The set $C := \{a_\beta : \beta < \alpha\} \cup \{b_\beta : \beta < \alpha\}$ is countable, and thus it can't be dense since S is not separable. So we choose an interval I_α which is disjoint from C . Of course $T := \{I_\alpha : \alpha < \omega_1\}$ has size \aleph_1 and is partially ordered by \supseteq . Moreover, $\alpha < \beta$ implies that either $I_\alpha \supsetneq I_\beta$ or $I_\alpha \cap I_\beta = \emptyset$. This means that $I_\alpha \supsetneq I_\beta$ implies $\alpha < \beta$, so every set of the form $\{I \in T : I \supseteq I_\alpha\}$ is well-ordered by \supseteq . Hence T is a tree.

We shall show that T has no uncountable branches and no uncountable antichains. Then of course $\text{height}(T) \leq \omega_1$; since every level is an antichain and $|T| = \aleph_1$ we obtain $\text{height}(T) = \omega_1$.

Observe that $I, J \in T$ are incomparable in T if and only if they are disjoint in S . By the ccc in S we get that antichains in T are at most countable. Finally, suppose towards a contradiction that \mathfrak{b} is a branch of height ω_1 . Then the left endpoints of the intervals $I \in \mathfrak{b}$ form a strictly increasing sequence $\{x_\alpha : \alpha < \omega_1\}$ of points of S . Clearly the intervals $(x_\alpha, x_{\alpha+1})$, $\alpha < \omega_1$ form an uncountable family of mutually disjoint open intervals in S , which contradicts the ccc hypothesis.

Let T be a Suslin tree. By last lemma we can assume w.l.o.g. that T is normal. Let $S := \{\mathfrak{b} : \mathfrak{b} \text{ is a branch in } T\}$. Each $x \in T$ has \aleph_0 -many immediate successor, thus we can assume that every set of immediate successor is equipped with a “local” order which is isomorphic to the one of the rational numbers. This enables us to define the order on S : if α is the least level where two branches $\mathfrak{a}, \mathfrak{b} \in S$ differ, then α is a successor ordinal and the relative points $a_\alpha \in \mathfrak{a}$ and $b_\alpha \in \mathfrak{b}$ are both successors of the same point at level $\alpha - 1$. We stipulate that $\mathfrak{a} < \mathfrak{b}$ iff a_α is smaller than b_α in the local order.

It’s immediate to check that S is linearly ordered and dense. Also, if $(\mathfrak{a}, \mathfrak{b})$ is an open interval in S , one easily finds some $x \in T$ such that $B_x \subseteq (\mathfrak{a}, \mathfrak{b})$, where B_x is $B_x := \{\mathfrak{c} \in S : x \in \mathfrak{c}\}$. Observe that if B_x and B_y are disjoint, then x and y are incomparable in T . Hence, for some $x, y \in T$,

$$(\mathfrak{a}, \mathfrak{b}) \cap (\mathfrak{c}, \mathfrak{d}) = \emptyset \Rightarrow B_x \cap B_y = \emptyset \Rightarrow x \perp y.$$

Since every antichain in T is at most countable, this means that S satisfies the ccc.

Finally, let F be a countable family of branches of T and let α be a countable ordinal greater than the length of any branch $\mathfrak{b} \in F$ (which obviously exists, because every branch of T is countable). Now let $x \in T$ be a node at level α . Of course $B_x \cap F = \emptyset$, and since B_x clearly contains some open interval of S , we have that F is not dense in S . Thus S is not separable and we are done. \square

2.2 Suslin trees in L

In this section we will prove that $V = L$ implies the existence of a Suslin tree. This result is due to Jensen [5], and it is significant because Jensen formulated the famous diamond principle¹ \diamond by extracting and isolating the combinatorial fact which holds in L and allows the construction of the Suslin tree. In fact, \diamond alone implies that there exists a Suslin tree. For this reason, the standard way to prove that if $V = L$ then a Suslin tree exists is to show that \diamond holds in L and then construct the Suslin tree using \diamond . In this work we decide to present the original proof and build the Suslin tree from scratch. This is a less known approach and will hopefully make clear to the reader where Jensen’s intuition about \diamond originated from. However, in Chapter 4 we will construct two particular Suslin trees using just \diamond .

Before starting with the math, it should be mentioned that Jensen’s theorem, which implies the relative consistency of $\neg\text{SH}$, was actually proved after the independence of SH had been established by means of forcing. Nonetheless, we will present the forcing approach in the next sections, as we believe that this will result in a clearer exposition from a mathematical point of view.

We start with a rather general lemma.

Lemma 2.5. *Let θ be a regular uncountable cardinal. Suppose that M is a countable structure such that $M \preceq H_\theta$. Then the following hold:*

¹The diamond principle asserts that there exists a sequence of sets $\langle S_\alpha : \alpha < \omega_1 \rangle$ with $S_\alpha \subseteq \alpha$, such that for every $X \subseteq \omega_1$, the set $\{\alpha < \omega_1 : X \cap \alpha = S_\alpha\}$ is a stationary subset of ω_1 .

1. If $a \in M$ and a is countable, then $a \subseteq M$.
2. $M \cap \omega_1 = \beta$, where β is a countable limit ordinal.
3. If $\theta > \omega_1$, then $\omega_1 \not\subseteq M$ but $\omega_1 \in M$, and “ ω_1 is the first uncountable ordinal” is true in M .
4. Let $T := \text{mos}'' M$ and let $\beta = M \cap \omega_1$ by the second point. Then $\text{mos}(\omega_1) = \beta$ and $\text{mos}(\xi) = \xi$ for all $\xi < \beta$.
5. $\beta = (\omega_1)^T$ and $T \models \text{ZFC} - \text{P}$.

Proof. Recall that H_θ is a model of $\text{ZFC} - \text{P}$. Thus ω and all $n \in \omega$ are definable in H_θ , so $\omega \in M$ and $\omega \subseteq M$. Now, if $a = \emptyset$ then trivially $a \subseteq M$. If $a \neq \emptyset$, then there exists $f: \omega \rightarrow a$ surjective. It's immediate to check that any such function belongs to H_θ . Since $M \preceq H_\theta$, at least one such f belongs to M as well. Observe that $f, n \in M$ implies $f(n) \in M$ (by $M \preceq H_\theta$ again), and since $a = \{f(n) : n \in \omega\}$ we have $a \subseteq M$, so (1) is proved.

For (2), note that by the previous point $M \cap \omega_1$ is an initial segment of ω_1 . Therefore $M \cap \omega_1 = \beta$ is an ordinal. Of course it can't be uncountable, and it is limit because if $\xi \in \beta$ then $\xi + 1 \in \beta$ by $M \preceq H_\theta$.

Point (3) is trivial by elementarity, because ω_1 is definable in H_θ as the first uncountable ordinal (and it is absolute between H_θ and V because any surjective function $f: \omega \rightarrow (\omega_1)^{H_\theta}$ in V would also belong to H_θ).

Point (4) follows immediately from the previous points, and in particular from $M \cap (\omega_1 \cup \{\omega_1\}) = \beta \cup \{\omega_1\}$.

For point (5), observe that the relation \in is extensional on H_θ and thus also on M by elementarity. So mos is an isomorphism, hence $T \simeq M \preceq H_\theta \models \text{ZFC} - \text{P} + \text{“}\omega_1 \text{ is the first uncountable cardinal”}$, and we are done. \square

The reason why last lemma will turn out to be useful is the following:

Lemma 2.6. *If $V = L$, then $L_\kappa = H_\kappa$ for all cardinals $\kappa \geq \omega$.*

Proof. See [14, p. 141]. \square

Thus we can replace H_θ with L_θ in Lemma 2.5 if $V = L$.

Lemma 2.7. *Let T be a tree and A a maximal antichain. Suppose that A is bounded in T , i.e. there is some $\alpha < \text{height}(T)$ such that $o(x) \leq \alpha$ for all $x \in A$. Then A is maximal in every end-extension of T .*

Proof. Let T' be an end-extension of T and let $\alpha < \text{height}(T)$ such that every element of A is at level $\leq \alpha$. Take $t' \in T' \setminus T$. We want to show that t' is comparable with some $a \in A$. Of course $o(t') \geq \text{height}(T) > \alpha$ because T' is an end-extension of T . Thus there exists $t < t'$ at level α of T . In turn, there exists $a \leq t$ which is an element of A . \square

Observe that any subset of a tree T is bounded if the height of T is a successor ordinal. Thus the request that A is bounded in last lemma is always satisfied in that case. But for the limit case, it's up to us to ensure that a maximal antichain doesn't "grow". Next lemma shows how to do it.

Lemma 2.8 (Sealing maximal antichains). *Let α be a countable limit ordinal, T a normal α -tree and \mathcal{A} a countable family of maximal antichains in T . Then there exists a normal end-extension T' of T of height $\alpha+1$ such that every $A \in \mathcal{A}$ is a maximal antichain in T' (and in turn in every end-extension of T' by last lemma).*

Proof. Write $\mathcal{A} = \{A_k : k \geq 1\}$. We want to show that for all $t \in T$ there exist an α -branch which hits t and every A_k . We build such branch by induction. Let $\langle \alpha_n : n \in \omega \rangle$ be cofinal in α . Suppose w.l.o.g. that $\alpha_0 = o(t)$ and let $t_0 := t$. Suppose inductively that we have defined $\langle t_i : i < n \rangle$ such that:

- the sequence is (weakly) increasing;
- $o(t_i) \geq \alpha_i$ for all $i < n$;
- $A_k \cap \downarrow \{t_0, \dots, t_{n-1}\} \neq \emptyset$ for all $1 \leq k < n$.

Since A_n is a maximal antichain, there exists $a_n \in A_n$ comparable with t_{n-1} . To define t_n , we shall distinguish three cases. If $a_n > t_{n-1}$ and $o(a_n) \geq \alpha_n$, define $t_n := a_n$. If $a_n \leq t_{n-1}$ and $o(t_{n-1}) \geq \alpha_n$, define $t_n := t_{n-1}$. If both $o(t_{n-1}), o(a_n) < \alpha_n$, then choose $t_n > \max(t_{n-1}, o(a_n))$ at level α_n (which exists by normality).

In any case, the induction hypothesis is preserved for $\langle t_i : i < n+1 \rangle$, and thus we obtain $\langle t_i : i \in \omega \rangle$ cofinal in T . By construction, such sequence induces an α -branch \mathbf{b}_t such that $t \in \mathbf{b}_t$ and $A_k \cap \mathbf{b}_t \neq \emptyset$ for all $k \in \omega$.

Finally, we define the extension $T' := T \cup \{b_t : t \in T\}$, where b_t are new elements which are put on top of the relative branch \mathbf{b}_t . It's easy to check that T' is a normal $(\alpha+1)$ -tree and an end-extension of T . Every $A \in \mathcal{A}$ is still a maximal antichain in T' , because by construction any b_t is greater than some $a \in A$. \square

Theorem 2.9 (Jensen). *If $V = L$, then there exists a Suslin tree.*

Proof. We construct the Suslin tree T by recursion on levels. As a set, T will simply be the ordinal ω_1 . Also, the construction will proceed in such a way that $T|_\alpha$ is normal, for all $\alpha < \omega_1$; T will also be normal. If we have already built the α th level U_α , for the successor level $U_{\alpha+1}$ we just put countably many nodes above each node of U_α . We decide to be more precise, even though this wouldn't be strictly necessary. Formally, we let $U_0 := \{0\}$, $U_1 := \omega \setminus \{0\}$, $U_{n+1} := \{\omega \cdot n + k : k \in \omega\}$ for $0 < n < \omega$, and $U_\alpha := \{\omega \cdot \alpha + k : k \in \omega\}$ for $\omega \leq \alpha < \omega_1$. So now we must define the ordering of T , recursively. For $U_{\alpha+1}$, we arbitrarily partition $U_{\alpha+1}$ into \aleph_0 -many countably infinite subsets. Every such set will be the set of immediate successors of one node of U_α , so that the normality is preserved. Now comes the hard part: for the limit step, we must choose which branches to extend and which to omit. The idea is to exploit Lemma 2.8 to "seal" every maximal antichain in the

partial tree $T|_\alpha$, for α limit. As one would expect, the viability of this method will rely on some specific properties of the constructible universe. The proof of some facts is rather technical, therefore we believe that it is more convenient to state them here as claims and provide their proofs at the end for the interested reader.

Claim 1: There exists a function $f: \omega_1 \rightarrow \omega_1$ such that for each $\alpha < \omega_1$,

$$(L_{f(\alpha)}, \in) \models \alpha \text{ is countable.}$$

Furthermore, f is definable L_{ω_2} (taking f as the $<_L$ -least such function).

Now we can define the limit levels: we choose such a function f and for every limit ordinal α we use Lemma 2.8 to seal all maximal antichains of $T|_\alpha$ which are elements of $L_{f(\alpha)}$. This is clearly possible because $L_{f(\alpha)}$ is countable. One technical detail: note that we can assume that the elements of the α th level of the $(\alpha + 1)$ -tree produced by Lemma 2.8 are precisely the ones of $U_\alpha := \{\omega \cdot \alpha + k : k \in \omega\}$. This is because $T|_\alpha$ is countable and normal, whereby the set $\{b_t : t \in T|_\alpha\}$ in the proof of 2.8 must have size precisely \aleph_0 , like U_α , which is the only thing we needed to check.

Finally, $T := \cup_{\gamma < \omega_1} U_\gamma$, and the order of T is the one we just defined by recursion. It's clear that T is an ω_1 -tree, so it remains to show that T has no uncountable antichain. Suppose to the contrary that this is not true. Let A be the $<_L$ -least uncountable maximal antichain of T .

Claim 2: T and A belong to L_{ω_2} , in which they satisfy the same definition as in V .

So we consider the structure

$$\mathcal{M} := (L_{\omega_2}, \in, \omega_1, T, A).$$

By Löwenheim-Skolem Theorem, there exists a countable elementary substructure $\mathcal{N} \preceq \mathcal{M}$ such that $\omega_1, T, A \in \mathcal{N}$. Such structure has the form

$$\mathcal{N} = (N, \in, \omega_1, T, A).$$

By Gödel Condensation Lemma and Lemma 2.5 (and using that $V = L$) it follows that its transitive realization $\text{mos}''\mathcal{N}$ has the form

$$\mathcal{N}' = (L_\beta, \in, \alpha, T \cap \alpha, A \cap \alpha),$$

where $\alpha = \omega_1 \cap N = (\omega_1)^{\mathcal{N}'}$ and β is a countable ordinal.

Observe that since $\mathcal{N} \preceq \mathcal{M}$ and $\text{mos}: \mathcal{N} \rightarrow \mathcal{N}'$ is an isomorphism, we have $T \cap \alpha = T^{\mathcal{N}'}$. In particular, $\mathcal{N}' \models \text{height}(T \cap \alpha) = \alpha$. By absoluteness, $\text{height}(T \cap \alpha) = \alpha$ also in V . Looking back at the definition of the levels of T , this clearly implies that $T \cap \alpha = T|_\alpha$. Similarly, $\mathcal{N}' \models \text{"}A \cap \alpha \text{ is a maximal antichain of } T \cap \alpha\text{"}$, hence by absoluteness $A \cap \alpha$ is a maximal antichain of $T|_\alpha$ in V as well.

Now, of course $\mathcal{N}' \models \text{"}\alpha \text{ is uncountable"}$, whereas $L_{f(\alpha)} \models \text{"}\alpha \text{ is countable"}$. Thus $\beta < f(\alpha)$, which implies $A \cap \alpha \in L_{f(\alpha)}$. This means that $A \cap \alpha$ was sealed at the α th step of the recursion, i.e. it's still maximal in T . Hence $A = A \cap \alpha$, so A is countable. \square

We now prove the two claims. Since these are the first results of this kind in our thesis, we shall be a bit pedantic.

Proof of Claim 1. Let α be a countable ordinal. Of course in V there exists $g: \omega \rightarrow \alpha$ surjective. It is immediate to check that $|\text{trcl}(g)| < \omega_1$, i.e. $g \in H_{\omega_1}$. Since $V = L$ implies $L_{\omega_1} = H_{\omega_1}$, this means that $g \in L_{\omega_1}$, and an easy absoluteness argument shows that $L_{\omega_1} \models g: \omega \xrightarrow{\text{onto}} \alpha$. By an easy application of the Reflection Theorem, there exists $\beta < \omega_1$ such that the formula is true in L_β .² Thus we proved that the statement

$$(\sigma) \quad \forall \alpha < \omega_1 \exists \beta < \omega_1 [(L_\beta, \in) \models \alpha \text{ is countable}].$$

holds in V . Now recall that the relation “ \models ” and the function $\gamma \mapsto L_\gamma$ are absolute for transitive models of $\text{ZF} - \text{P}$. Hence every notion used in σ is absolute for L_{ω_2} , which then satisfies σ . Thus in L_{ω_2} there exists a function $f: \omega_1 \rightarrow \omega_1$ such that for each $\alpha < \omega_1$,

$$(L_{f(\alpha)}, \in) \models \alpha \text{ is countable}.$$

We can suppose w.l.o.g. that f is the $<_L$ -least such function, so that f is absolute for L_{ω_2} . ■

Proof of Claim 2. If we show that $T \in L_{\omega_2}$ and it’s absolute, then $A \in L_{\omega_2}$ and its absoluteness follow immediately: “being a maximal antichain” is absolute for L_{ω_2} (easy to verify) and $A \subseteq T \in L_{\omega_2}$ ($= H_{\omega_2}$ since $V = L$) implies $A \in L_{\omega_2}$.

Now, to prove the statement about T we must show that the recursion step $\alpha \mapsto U_\alpha$ is absolute for L_{ω_2} . The case when α is a successor ordinal is easy. For the limit case, observe that U_α is essentially the α th level of the tree $H(T|_\alpha, \mathcal{S}_\alpha)$, where

$$\mathcal{S}_\alpha := \{A : (A \text{ is a maximal antichain of } T|_\alpha)^{L_{f(\alpha)}}\}$$

and H is a function which takes the tree $T|_\alpha$ and the countable family \mathcal{S}_α and returns an $(\alpha + 1)$ -tree as in the statement of Lemma 2.8. It’s easy to check that H is absolute for L_{ω_2} . Furthermore, by the absoluteness of the notions involved (in particular of \models and $\gamma \mapsto L_\gamma$), we obtain that the formula

$$(A \text{ is a maximal antichain of } T|_\alpha)^{L_{f(\alpha)}}$$

is absolute between L_{ω_2} and V . Therefore $(\mathcal{S}_\alpha)^{L_{\omega_2}} = \mathcal{S}_\alpha$, which in turn implies that $(H(T|_\alpha, \mathcal{S}_\alpha))^{L_{\omega_2}} = H(T|_\alpha, \mathcal{S}_\alpha)$. Hence the induction step in the construction of T is absolute, and by the absoluteness of functions defined by recursion we are done. ■

²Of course g belongs to L_β for some $\beta < \omega_1$, but we used the Reflection Theorem to make sure that such L_β indeed sees g as a surjection $\omega \rightarrow \alpha$ (L_β need not be a model of all $\text{ZFC} - \text{P}$). This could actually be checked by an easy absoluteness argument. We will omit such details hereafter.

2.3 Forcing and Suslin trees

2.3.1 A forcing model where there is a Suslin tree

Proving the consistency of $\neg\text{SH}$ was the first breakthrough about Suslin's hypothesis since its characterization in terms of Suslin trees by Kurepa, and possibly since its very formulation. This result was achieved independently by Tennenbaum and Jech. While Jech was the first to write a publication in 1967, a letter by Solovay to Kanamori [10] witnesses that Tennenbaum already proved the result in 1963 (just after Cohen's development of forcing), although he published it [20] in 1968. To do that, he used a forcing poset consisting of finite trees ordered by extension. Such poset is ccc, and probably this is the reason why he came up with it: Cohen had proved that ccc orders preserve cardinals. On the other hand, Jech's forcing conditions are countable approximations to a Suslin tree, ordered by end-extension. The peculiarity of this poset is that cardinals are preserved because it is countably closed, not because it's ccc. Our choice is to present Jech's proof, because the core argument (i.e. the way of extending the tree at limit stages to seal the antichain) is essentially the same we've used for Jensen's Theorem 2.9.

The following is an easy and well-known property of forcing. We include it here to reference it in following proofs and make the exposition clearer. For a proof see [14, p. 284].

Fact 2.10. *Let M be a countable transitive model for ZFC and P a poset. Let $\pi \in M^P$ be a name and $r \in P$. If $r \Vdash \exists x \in \pi \varphi(x)$, then there is a $q \leq r$ and a $\sigma \in \text{dom}(\pi)$ such that $q \Vdash [\sigma \in \pi \wedge \varphi(\sigma)]$.*

In particular, suppose that π is a canonical name $\check{a} \in M^P$. Then $\sigma = \check{b}$ for some $b \in M$, and we obtain that:

If $r \Vdash \exists x \in \check{a} \varphi(x)$, then there are some $q \leq r$ and some $b \in a$ such that $q \Vdash \varphi(\check{b})$.

Theorem 2.11 (Jech). *There is a generic extension of V in which there exists a Suslin tree.*

Proof. We let the set of forcing conditions P be the family of trees described in Example 1.2. We define the order on P by:

$$T_1 \leq T_2 \text{ if and only if } T_1 \text{ is an end-extension of } T_2.$$

Let G be a generic filter on P and let $\mathcal{T} := \bigcup\{T : T \in G\}$. We shall prove that $V[G] \models$ “ \mathcal{T} is a normal Suslin tree”.

First, it's immediate to show by induction that any conditions $T_1, T_2 \in P$ are either comparable or incompatible. Hence G consists of comparable conditions, which means that \mathcal{T} is coherent union of trees in P . It follows immediately that \mathcal{T} is a normal tree with $\text{height}(\mathcal{T}) \leq \omega_1$.

Similarly, if $(T_n)_{n < \omega}$ is a decreasing sequence in P , then $\bigcup_{n < \omega} T_n$ is still in P and is a lower bound for the sequence. So P is countably closed and thus ω_1 is preserved in $V[G]$.

Now we want to show that the height of \mathcal{T} is precisely ω_1 . We must check that for all $\alpha < \omega_1$ there is $T \in G$ such that $\text{height}(T) \geq \alpha$. Using that G is generic, if we show that

$$D_\alpha := \{T \in P : \text{height}(T) \geq \alpha\}$$

is dense for all $\alpha < \omega_1$, we are done. Let $T_0 \in P$. We want to find an end-extension of T_0 which is in P and has height $\geq \alpha$. Actually, it suffices to extend it by one more level, because then iterating the argument and taking unions at limit steps we prove the general case.

If $\text{height}(T_0)$ is a successor ordinal, the extension is trivially (and uniquely) obtained. If $\text{height}(T_0)$ is limit, we simply apply Lemma 2.8 with $\mathcal{A} := \emptyset$ ³.

So it's left to show that \mathcal{T} has no uncountable antichains. The idea is to exploit again Lemma 2.8, but here we will need a weaker version than the one we used to construct the Suslin tree in L . In fact, we will need to “seal” only one maximal antichain at a time.

Suppose that, in $V[G]$, A is an antichain of \mathcal{T} . Without loss of generality we can assume that A is maximal. So there is $T \in G$ such that

$$T \Vdash \dot{A} \text{ is a maximal antichain in } \dot{\mathcal{T}}.$$

We claim that the following set is dense below T :

$$D := \{T' \leq T \mid V \text{ models that there is a bounded maximal antichain } A' \text{ in } T' \text{ such that } T' \Vdash A' \subseteq \dot{A}\}.$$

If this is true, then some $T' \in D$ belongs to G , which means that in $V[G]$ there exists a bounded maximal antichain A' of T' such that $A' \subseteq A$. Since $T \in G$, we have that \mathcal{T} is an end-extension of T' , and hence by Lemma 2.7 A' is maximal in \mathcal{T} . Therefore $A = A'$, which is countable, as wanted.

To show that D is dense below T , take $T_0 \leq T$. We shall construct $T' \leq T_0$ such that $T' \in D$. Since $T_0 \Vdash T_0 \in \dot{G}$, it follows that $T_0 \Vdash \text{“}\dot{\mathcal{T}} \text{ end-extends } T_0\text{”}$. Using also that $T_0 \Vdash \text{“}\dot{A} \text{ is a maximal antichain in } \dot{\mathcal{T}}\text{”}$, we obtain

$$\begin{aligned} T_0 \Vdash & \text{ for any } s \in T_0 \text{ there exist } T'_0 \leq T_0 \text{ end-extended by } \dot{\mathcal{T}} \\ & \text{ and } t_s \in T'_0 \text{ such that (i) } s \text{ and } t_s \text{ are comparable and (ii) } t_s \in \dot{A}. \end{aligned}$$

By (many applications of) Fact 2.10, we have that, in V ,

$$\begin{aligned} \text{for any } s \in T_0 \text{ there exist } T'_0 \leq T_0 \text{ and } t_s \in T'_0 \text{ such that} \\ \text{(i) } s \text{ and } t_s \text{ are comparable and (ii) } T'_0 \Vdash t_s \in \dot{A}. \end{aligned}$$

Working in V and repeating this argument, we can define a sequence of end-extensions of T_0 which recursively satisfy (i) and (ii) for all $s \in T_0$ and which end-extend the previous

³To be formally precise, the b_t 's from the proof of Lemma 2.8 must be defined here as $b_t := \bigcup \mathfrak{b}_t$, so that the extension is really an element of P .

ones. Being T_0 countable, also the sequence is countable. Since P is countably closed, we obtain $T_1 \in P$ such that $T_1 \leq T_0$ and

$$\forall s \in T_0 \exists t_s \in T_1 [s \text{ and } t_s \text{ are comparable and } T_1 \Vdash t_s \in \dot{A}].$$

Recursively using this argument, we define a sequence $T_0 \geq T_1 \geq \dots \geq T_n \geq \dots$ such that, for each $n < \omega$, $T_{n+1} \leq T_n$ and

$$(2.1) \quad \forall s \in T_n \exists t_s \in T_{n+1} [s \text{ and } t_s \text{ are comparable and } T_{n+1} \Vdash t_s \in \dot{A}].$$

Now we let $T_\infty := \bigcup_{n < \omega} T_n \in P$ and $A' := \{t_s : s \in T_\infty\}$. By (2.1), A' is a maximal antichain in T_∞ and $T_\infty \Vdash A' \subseteq \dot{A}$. By Lemma 2.8 we get an end-extension T' of T_∞ such that A' is a bounded maximal antichain in T' . Of course $T' \Vdash A' \subseteq \dot{A}$ and hence $T' \in D$. \square

2.3.2 A forcing model where there are no Suslin trees

After establishing the relative consistency of $\neg\text{SH}$, Tennenbaum tried to prove the consistency of SH . He realized that if T is a Suslin tree, then using T itself as the forcing poset (with the reverse order) has the following two properties:

- The fact that every antichain in T is countable means precisely that T is a ccc forcing poset (seen upside down). Thus cardinals are preserved.
- A generic filter on T is an uncountable branch.

Therefore, forcing with T “kills” T , meaning that in the generic extension T is not Suslin anymore (and of course once T is killed it stays dead in further generic extensions). The problem is that new Suslin trees might have sprung up in the generic extension. Hence, some kind of *iteration* was needed. But trying to iterate the forcing by simply taking products in the ground model has the disadvantage of possibly loosing the ccc-ness of the forcing⁴. This issue was eventually solved thanks to the collaboration with Solovay in 1964-65, which led to the development of a groundbreaking way of iterating forcing, a technique that is still among the main tools for modern set theory. The key idea was to consider forcing notions which occur in an intermediate generic extension, rather than in the ground model. In 1971, Solovay and Tennenbaum published [17] the proof of consistency of SH , together with a presentation of their iterated forcing methods; in the same year, Tarski defined their work on Suslin’s hypothesis as the most important in set theory since Cohen’s work.

⁴This can happen already with a two-step iteration. For example, note that forcing with $T \times T$ is the same as forcing with T twice by the Product Lemma. Since in the first step of the iteration we obtain an uncountable branch of T in a generic extension, obviously T is not ccc anymore in that extension (assuming that T is normal). This means that cardinals *may* collapse with this forcing. In [2, p. 38], a similar example is given where cardinals *must* collapse.

Actually, in the final publication, the consistency of SH was derived as an immediate consequence of other results concerning a new statement: Martin's Axiom⁵ (MA). In 1967, Martin extracted the statement of MA from the proof by Solovay and Tennenbaum, observed that its consistency could be proved by the same techniques used for SH. MA is a focal “axiom”, which is stronger than SH and has a wide range of consequences. For this reason, the consistency of SH is usually presented as a corollary of the consistency of MA (even in the original work by Solovay and Tennenbaum). Following the spirit of this work, we will prove the consistency of SH directly, by adapting the proof for the consistency of MA.

On a final note, this proof should be read after the one of Theorem 3.12, which kills Kurepa trees by means of a more “rudimentary” way for iterating forcing, i.e. product forcing. While this is not strictly necessary, we believe that the easier setting of Theorem 3.12 will offer a better intuition to proceed with this section.

Lemma 2.12. *Let P be a forcing poset. Suppose that $\tau \in V^P$ is such that $1_P \Vdash [|\tau| \leq \check{\lambda}]$ for some cardinal λ . Then every $\sigma \in V^P$ which is an element of τ (i.e. $p \Vdash [\sigma \in \tau]$ for some $p \in P$) can be represented by a function from an antichain in P into λ . In particular, if P is ccc, $|P| \leq \aleph_2$ and $\lambda < \omega_2$, then such functions are at most $\aleph_2^{\aleph_0}$ -many.*

Proof. Formally, we want to show the following:

For all $\sigma \in V^P$, if $p \Vdash [\sigma \in \tau]$ for some $p \in P$, then we can define a function $\varphi_\sigma: A \rightarrow \lambda$ such that A is an antichain of P and the assignment $\sigma \mapsto \varphi_\sigma$ is defined in such a way that $\varphi_\sigma = \varphi_{\sigma'}$ implies $p \Vdash \sigma = \sigma'$.

Since $1_P \Vdash |\tau| \leq \check{\lambda}$, for some $f \in V^P$ we have $1_P \Vdash [f: \tau \xrightarrow{1-1} \check{\lambda}]$. Suppose that $p \Vdash \sigma \in \tau$. Then the set

$$D_\sigma := \{r : \text{for some } \alpha_r < \lambda, r \Vdash f(\sigma) = \check{\alpha}_r\}$$

is predense under p . Take a maximal antichain A_σ in $\downarrow D_\sigma$ (and thus a maximal antichain in $P \restriction p$) and define the function

$$\begin{aligned} \varphi_\sigma: A_\sigma &\rightarrow \lambda \\ r &\mapsto \alpha_r \end{aligned}$$

(the function is well-defined because for each $r \in D$ there is obviously only one corresponding α_r).

Now suppose that $q \Vdash \sigma' \in \tau$. We want to show that $\varphi_\sigma = \varphi_{\sigma'}$ implies $p \Vdash \sigma = \sigma'$. Since $\text{dom}(\varphi_\sigma) = \text{dom}(\varphi_{\sigma'})$, we have that the same set (we call it A) is a maximal antichain under p and q . This clearly means that, for any generic filter G , $p \in G \Leftrightarrow q \in G$. Take any generic G such that $p \in G$. So there is some $r \in G \cap A$, and hence $f(\sigma_G) = \varphi_\sigma(r)$. Since $q \in G$, the same argument works for σ' , so that $f(\sigma'_G) = \varphi_{\sigma'}(r)$, whereby $f(\sigma_G) = f(\sigma'_G)$. Because f is injective, $V[G] \models \sigma_G = \sigma'_G$, and by the arbitrariness of G we obtain $p \Vdash \sigma = \sigma'$. \square

⁵Martin's Axiom asserts: If $(P, <)$ is a ccc poset, $\kappa < 2^{\aleph_0}$ is an infinite cardinal and \mathcal{D} is a collection of at most κ dense subsets of P , then there exists a \mathcal{D} -generic filter on P . It's immediate to check that SH follows from Martin's Axiom plus $\neg\text{CH}$.

Theorem 2.13 (Solovay-Tennenbaum). *There is a generic extension of V where there exist no Suslin trees.*

Proof. Suppose without loss of generality that $2^{\aleph_1} = \aleph_2$ in V . We will construct a finite support iteration P of length ω_2 given by a sequence $\langle \dot{Q}_\alpha : \alpha < \omega_2 \rangle$ such that

- (i) $\Vdash_\alpha [\dot{Q}_\alpha \text{ is a ccc forcing poset}]$, for each $\alpha < \omega_2$;
- (ii) $|P_\alpha| \leq \aleph_2$ for each $\alpha \leq \omega_2$.

The sequence will be defined in such a way that every Suslin tree will be killed.

Note that each P_α for $\alpha \leq \omega_2$ is ccc by the result about iterated forcing mentioned in the preliminaries.

Before proceeding with the construction of the sequence, observe that:

Claim: If we can ensure that $\Vdash_\alpha |\dot{Q}_\alpha| \leq \check{\omega}_1$ for each $\alpha < \omega_2$, then (ii) holds.

Proof: We show this by induction on α . If α is a limit ordinal and $|P_\beta| \leq \aleph_2$ for all $\beta < \alpha$, then $|P_\alpha| \leq \aleph_2$ because the elements of P_α are α -sequences with finite support. For the successor step, assume that $|P_\alpha| \leq \aleph_2$. Also, $\Vdash_\alpha |\dot{Q}_\alpha| \leq \check{\omega}_1$ by hypothesis. Using Lemma 2.12, we may assume w.l.o.g. that

$$|\{q \in V^{P_\alpha} : p \Vdash [q \in \dot{Q}_\alpha] \text{ for some } p \in P_\alpha\}| \leq \aleph_2^{\aleph_0},$$

which is \aleph_2 by the assumption $2^{\aleph_1} = \aleph_2$. It follows immediately that $P_{\alpha+1} = P_\alpha * \dot{Q}_\alpha$ has size $\leq \aleph_2$. ■

We define \dot{Q}_α by induction on $\alpha < \omega_2$. Fix a bijection $\pi: \omega_2 \rightarrow \omega_2 \times \omega_2$ such that if $\pi(\alpha) = (\beta, \gamma)$ then $\beta \leq \alpha$ (it's easy to find such a function). Fix $\alpha < \omega_2$ and suppose that we have already defined $\langle \dot{Q}_\beta : \beta < \alpha \rangle$. Let $\beta < \alpha$ be arbitrary. We want to count the nice names for subsets of $(\omega_1 \times \omega_1) \in V^{P_\beta}$. Of course $|\text{dom}((\omega_1 \times \omega_1))| = \aleph_1$. Since P_β is ccc and has size at most \aleph_2 , there are no more than $\aleph_2^{\aleph_1} = \aleph_2$ nice names for subsets of $(\omega_1 \times \omega_1)$ in P_β (cf. preliminaries). This is true for any $\beta < \alpha$. Now let $\pi(\alpha) = (\beta, \gamma)$. So we finally define \dot{Q}_α :

$$\dot{Q}_\alpha := \begin{cases} \begin{cases} \text{the } \gamma\text{th nice name } \tau \in P_\beta \\ \text{for a subset of } (\omega_1 \times \omega_1) \end{cases} & \text{if } \Vdash_\alpha [\tau \text{ is a ccc forcing poset}]; \\ \text{a } P_\beta\text{-name for the order on } \{0\} & \text{otherwise} \end{cases}$$

(observe that the condition makes sense, because $\beta \leq \alpha$ and thus τ can be seen as a P_α -name as well).

Of course then (i) holds. It remains to check that $\Vdash_\alpha |\dot{Q}_\alpha| \leq \check{\omega}_1$. But this is clearly true, since $\Vdash_\alpha \dot{Q}_\alpha \subseteq (\omega_1 \times \omega_1)$.

Finally, let P be the finite support iteration of $\langle \dot{Q}_\alpha : \alpha < \omega_2 \rangle$. Let G be a generic filter on P . We claim that $V[G]$ models that there are no Suslin trees.

Let (T, \sqsubset) be an ω_1 -tree in $V[G]$ with only countable antichains. We can suppose w.l.o.g. that $T \subseteq \omega_1$. We claim that $V[G] \models T$ is not Suslin. For simplicity, assume that

T “grows downwards” (formally we just take the inverse order), so that (T, \sqsupseteq) is a ccc forcing poset. Also, to make the notation more intuitive, we write T in place of \sqsupseteq . Let \dot{T} be a nice name for a subset of $(\omega_1 \times \omega_1)$ such that $\dot{T}_G = T$. Observe that this is possible because $V[G] \models T \subseteq \omega_1 \times \omega_1$, and $(\omega_1 \times \omega_1)_G = \omega_1 \times \omega_1 = \omega_1^{V[G]} \times \omega_1^{V[G]}$ because P is ccc.

Claim: We can assume w.l.o.g. that $\Vdash_P [\dot{T} \text{ is a ccc forcing poset}]$.

Proof: Let $\varphi(x)$ be the formula which asserts that “ x is a ccc forcing poset”. We have that $V[G] \models \varphi(T)$, so there are $\dot{T} \in V^P$ and $p \in G$ such that $\dot{T}_G = T$ and $p \Vdash_P \varphi(\dot{T})$. We want p to be 1_P and \dot{T} to be a nice name. This need not be true in general, but we shall find a name for T which satisfies such requirements. Observe that

$$1_P \Vdash_P \exists y [\varphi(y) \wedge [\varphi(\dot{T}) \Rightarrow y = \dot{T}]],$$

because we can let y be any appropriate ccc forcing order (e.g. the inverse of the usual order on ω_1 , restricted to the domain of T) when $\neg\varphi(\dot{T})$. Using the Maximal Principle (cf. Preliminaries), we can fix a name $\tau \in V^P$ such that

$$1_P \Vdash_P \varphi(\tau) \wedge [\varphi(\dot{T}) \Rightarrow \tau = \dot{T}].$$

Let σ be a nice name (cf. Preliminaries) for a subset of $(\omega_1 \times \omega_1)$ such that

$$1_P \Vdash \tau \subseteq (\omega_1 \times \omega_1) \Rightarrow \tau = \sigma.$$

It’s straightforward to check that σ enjoys the required features. ■

To make sure that our construction works, we will need that \dot{T} was “hit” at some step of the recursion. This will follow by the next two claims.

Claim: There is $\beta < \omega_2$ such that (w.l.o.g.) $\dot{T} \in V^{P_\beta}$.

Proof: This is shown by observing that, being \dot{T} a nice name and P a ccc poset, the elements of P mentioned in \dot{T} are $< \omega_2$ -many. A precise argument would be almost identical to the one in the proof of Theorem 3.12.

The “w.l.o.g.” is just because P_β is not formally a subset of P , but there is a complete embedding from the former into the latter. ■

So we produced a \dot{T} which is a nice name in V^{P_β} for T and is such that $\Vdash_P [\dot{T} \text{ is a ccc forcing poset}]$. It is the γ th one, for some $\gamma < \omega_2$. Let α be such that $\pi(\alpha) = (\beta, \gamma)$.

Claim: \dot{T} is a nice P_α -name as well, and $\Vdash_\alpha [\dot{T} \text{ is a ccc forcing poset}]$.

Proof: To verify that \dot{T} is nice also in V^{P_α} is easy.

Let $\varphi(x)$ be “ x is a ccc forcing poset” as before. Assume towards a contradiction that $\nVdash_\alpha \varphi(\dot{T})$, so that there is $p \in P_\alpha$ with $p \Vdash_\alpha \neg\varphi(\dot{T})$. Let G be generic for P and such that $p \in G$. Using well-known facts about iterated forcing⁶, it follows that $G_\alpha := G \restriction P_\alpha$ is generic for P_α , $V[G_\alpha] \subseteq V[G]$ and $\dot{T}_G = \dot{T}_{G_\alpha} \in V[G_\alpha]$. But of course $V[G] \models \varphi(\dot{T}_G)$, whereas $V[G_\alpha] \models \neg\varphi(\dot{T}_G)$. An easy absoluteness argument shows that this is impossible (one uses that P and P_α are ccc, so that $\omega_1^{V[G]} = \omega_1^{V[G_\alpha]}$). ■

⁶See also [14, Lemma IV.4.2 and Lemma IV.4.4, pp. 270-71].

So, by definition of \dot{Q}_α , we have

$$\dot{Q}_\alpha = \dot{T}.$$

Furthermore, if we let $G_\alpha := G \restriction P_\alpha$, we have

$$(\dot{Q}_\alpha)_{G_\alpha} = \dot{T}_{G_\alpha} = \dot{T}_G = T.$$

Recall that $P_{\alpha+1} = P_\alpha * \dot{Q}_\alpha$. By the theory of iterated forcing, in $V[G_{\alpha+1}]$ there is a generic filter H on $(\dot{Q}_\alpha)_{G_\alpha} = T$ over $V[G_\alpha]$. It's immediate to see that H must be an ω_1 -branch in T . Of course such branch belongs to $V[G]$ as well, therefore T is not a Suslin tree in $V[G]$. \square

One can see that the generic extension constructed in last theorem satisfies $\text{ZFC} + \text{SH} + \neg\text{CH}$. In [2], Devlin shows how the failure of CH is inherently related to the core idea of Solovay-Tennenbaum's argument: to kill the *Suslinity* of a Suslin tree T by adding a branch to it, which is achieved by forcing with T itself. A remarkable result by Jensen shows that it's possible to obtain a model for $\text{ZFC} + \text{SH} + \text{CH}$ by using a different approach. We refer the interested reader to [2].

Chapter 3

Kurepa trees

3.1 Kurepa's Hypothesis

Definition 3.1. A *Kurepa tree* is an ω_1 -tree which has only levels of countable size and at least \aleph_2 -many uncountable branches.

Definition 3.2. A *Kurepa family* is $\mathcal{F} \subseteq \mathcal{P}(\omega_1)$ such that $|\mathcal{F}| \geq \aleph_2$ and the set $\{X \cap \alpha : X \in \mathcal{F}\}$ is countable for all $\alpha < \omega_1$.

In 1942, Kurepa formulated *Kurepa's Hypothesis*, which asserts that there exists a Kurepa tree. He conjectured that such hypothesis doesn't hold, which is false, since it is independent of ZFC, as we will show in a while. First, as one might expect:

Proposition 3.3. *There exists a Kurepa tree if and only if there exists a Kurepa family.*

Proof. Suppose we have a Kurepa tree T . T has size \aleph_1 , so we can assume w.l.o.g. that $T = \omega_1$. We can also assume that $\alpha <_T \beta$ implies $\alpha < \beta$, because a simple re-ordering defined by induction on levels gives such a tree. By defining $\mathcal{F} := \{\mathfrak{b} : \mathfrak{b} \text{ is an } \omega_1\text{-branch of } T\}$ we clearly obtain a Kurepa family.

Let \mathcal{F} be a Kurepa family. For any $X \in \mathcal{F}$, define $f_X : \omega_1 \rightarrow \mathcal{P}(\omega_1)$ given by $f_X(\alpha) := X \cap \alpha$. Now let $U_\alpha := \{f_X \restriction \alpha : X \in \mathcal{F}\}$ for all $\alpha < \omega_1$ and $T := \bigcup_{\alpha < \omega_1} U_\alpha$. Endowing T with the " \subseteq " order we get a tree whose α th level is U_α , thus countable. Also, every f_X corresponds to an ω_1 -branch in T . Since there are $\geq \aleph_2$ -many f_X 's, T is a Kurepa tree. \square

In this chapter we will present the main consistency results about Kurepa trees, organized symmetrically to the ones about Suslin trees: first we will construct a Kurepa tree (actually, a Kurepa family) in L , and then we will establish the independence of Kurepa's Hypothesis solely via forcing (but for the consistency of the negation of Kurepa's Hypothesis we will need to assume the existence of an inaccessible cardinal).

3.2 Kurepa families in L

The existence of a Kurepa family in L was shown by Solovay around 1971, after the independence of Kurepa's hypothesis had already been proved. He was inspired by Jensen's Suslin tree construction in L . Indeed, the strategy of the proof is the same, even though some details are more involved. Similarly to what happened with \diamond , a combinatorial principle was extracted from this proof: diamond plus¹ (\diamond^+). \diamond^+ holds in L and implies that there exists a Kurepa tree. It also implies \diamond , but the reverse implication is not true.

We proceed now with Solovay's proof.

Remark. Observe that by the first point of Lemma 2.5, it follows immediately that if $M \preceq H_{\omega_1}$ with M countable, then M is transitive.

Theorem 3.4 (Solovay). *If $V = L$ then there exists a Kurepa family.*

Proof. The general structure of this proof will be very similar to the one of Theorem 2.9. Therefore we shall skip some details which were already discussed in the construction of the Suslin tree.

Assume $V = L$. We define the function $f: \omega_1 \rightarrow \omega_1$ by

$$f(\alpha) := \text{the least } \gamma \text{ such that } \alpha \in L_\gamma \preceq (L_{\omega_1}, \in).$$

The function f is well-defined: of course there is a smallest elementary substructure (M, \in) of (L_{ω_1}, \in) such that $\alpha \in M$. M is clearly countable, thus by last remark M is transitive² and using Gödel Condensation Lemma we obtain that $M = L_\gamma$ for some $\gamma < \omega_1$. Now consider the following family:

$$\mathcal{F} := \{X \subseteq \omega_1 \mid X \cap \alpha \in L_{f(\alpha)} \text{ for all } \alpha < \omega_1\}.$$

Of course $\{X \cap \alpha : X \in \mathcal{F}\}$ is countable for any α , because $L_{f(\alpha)}$ is. If we show that $|\mathcal{F}| \geq \aleph_2$ then \mathcal{F} is a Kurepa family, as wanted.

Assume towards a contradiction that $|\mathcal{F}| \leq \aleph_1$. Then there is an enumeration $C = \langle X_\xi : \xi < \omega_1 \rangle$ of \mathcal{F} . Suppose w.l.o.g. that C is the $<_L$ -least such enumeration.

Claim 1: C belongs to L_{ω_2} , in which it satisfies the same definition as in V .

Proof: The fact that $C \in L_{\omega_2}$ is immediate using that $L_{\omega_2} = H_{\omega_2}$. Recalling that the function $\theta \mapsto L_\theta$ and the relation " \preceq " are absolute (for L_{ω_2}), it's easy to check that f and, in turn, \mathcal{F} are absolute for L_{ω_2} . Thus $L_{\omega_2} \models C: \omega_1 \xrightarrow{\text{onto}} \mathcal{F}$, and we conclude by the absoluteness of $<_L$. \blacksquare

Now we recursively define a chain of elementary substructures of (L_{ω_2}, \in) , which will look like this:

$$N_0 \preceq N_1 \preceq \cdots \preceq N_\nu \preceq \cdots \preceq (L_{\omega_2}, \in)$$

¹ \diamond^+ says that there is a sequence $\langle \mathcal{A}_\alpha : \alpha < \omega_1 \rangle$, with \mathcal{A}_α a countable subset of $\mathcal{P}(\alpha)$, such that the following is true: $\forall A \subseteq \omega_1 \exists C \subseteq \omega_1 [C \text{ is club and } \forall \alpha \in C [A \cap \alpha \in \mathcal{A}_\alpha \text{ and } C \cap \alpha \in \mathcal{A}_\alpha]]$.

²Here we are using that $V = L$ to ensure that $H_{\omega_1} = L_{\omega_1}$.

for $\nu < \omega_1$. The chain is defined as follows:

$$(3.1) \quad \begin{aligned} N_0 &\text{ is the smallest elementary substructure of } L_{\omega_2}; \\ N_{\nu+1} &\text{ is the smallest } N \preceq L_{\omega_2} \text{ such that } N_\nu \subseteq N \text{ and } N_\nu \in N; \\ \text{for } \eta \text{ limit, } N_\eta &:= \bigcup_{\xi < \eta} N_\xi. \end{aligned}$$

Observe that:

- every N_ν is countable;
- there always exists N as in the successor step, because $N_\nu \in L_{\omega_2}$ since N_ν is a countable subset of L_{ω_2} (and $L_{\omega_2} = H_{\omega_2}$);
- if $\nu < \mu$ then $N_\nu \preceq N_\mu$ because $N_\nu \subseteq N_\mu$ and both are elementary substructures of L_{ω_2} ;
- for each $\nu < \omega_1$, by Lemma 2.5 we have $\omega_1 \cap N_\nu = \alpha_\nu$ for some $\alpha_\nu < \omega_1$.

The sequence $\langle \alpha_\nu : \nu < \omega_1 \rangle$ is continuous and strictly increasing. Continuity is straightforward:

$$\omega_1 \cap N_\eta = \omega_1 \cap \bigcup_{\xi < \eta} N_\xi = \bigcup_{\xi < \eta} \omega_1 \cap N_\xi = \sup_{\xi < \eta} \alpha_\xi.$$

It's trivial that the sequence is increasing, but the fact that it is *strictly* increasing is a bit more delicate. Recall that $N_\nu \in N_{\nu+1} \preceq L_{\omega_2}$. By Lemma 2.5, also $\omega_1 \in N_{\nu+1}$. So $N_\nu \cap \omega_1 \in N_{\nu+1}$ by elementarity, i.e. $\alpha_\nu \in N_{\nu+1}$. But of course $\alpha_{\nu+1} = N_{\nu+1} \cap \omega_1 \notin N_{\nu+1}$, therefore $\alpha_\nu \neq \alpha_{\nu+1}$, and we are done.

Let $X := \{\alpha_\nu : \alpha_\nu \notin X_\nu\}$. Note that X is well-defined because $\langle \alpha_\nu : \nu < \omega_1 \rangle$ is strictly increasing and thus injective. Obviously $X \neq X_\xi$ for all $\xi < \omega_1$. We claim that $X \in \mathcal{F}$, which contradicts the fact that C enumerates all the elements of \mathcal{F} .

So we want to show that $X \cap \alpha \in L_{f(\alpha)}$ for all $\alpha < \omega_1$. We prove this by induction on α . The case where α is not a limit point of the sequence $\langle \alpha_\nu \rangle_{\nu < \omega_1}$ is easy. For, let α_ν be the greatest $\alpha_\nu < \alpha$. Then clearly we have either $X \cap \alpha = X \cap \alpha_\nu$ or $X \cap \alpha = (X \cap \alpha_\nu) \cup \{\alpha_\nu\}$. In either case $X \cap \alpha \in L_{f(\alpha)}$, because $\alpha_\nu \in L_{f(\alpha_\nu)}$ and $X \cap \alpha_\nu \in L_{f(\alpha_\nu)} \subseteq L_{f(\alpha)}$ (the “ \subseteq ” is by the induction hypothesis and the “ \subseteq ” is because f is trivially an increasing function)³. It is left to show that $X \cap \alpha_\eta \in L_{f(\alpha_\eta)}$ if η is a limit ordinal.

Claim 2:

- (i) $\langle \alpha_\nu : \nu < \eta \rangle \in L_{f(\alpha_\eta)}$;
- (ii) $\langle X_\xi \cap \alpha_\eta : \xi < \alpha_\eta \rangle \in L_{f(\alpha_\eta)}$.

³Also, note that $X \cap \alpha_\nu \in L_{f(\alpha)}$ and $\alpha_\nu \in L_{f(\alpha)}$ implies $(X \cap \alpha_\nu) \cup \{\alpha_\nu\} \in L_{f(\alpha)}$ because $L_{f(\alpha)}$ is a transitive model of $\text{ZFC} - \text{P}$ by definition of f .

Proof: Let's prove (ii) first. For each $\nu < \omega_1$, let mos_ν be the Mostowski collapsing function of N_ν . For every $\nu < \omega_1$, $\text{mos}_\nu'' N_\nu = L_{\delta(\nu)}$ for some $\delta(\nu) < \omega_1$ by Gödel Condensation Lemma. Since $\omega_1 \cap N_\nu = \alpha_\nu$, by Lemma 2.5 we have $\text{mos}_\nu(\omega_1) = \alpha_\nu$. By Claim 1, C is a definable element of L_{ω_2} , hence $C \in N_\nu$ for all $\nu < \omega_1$ by elementarity. Using together that $N_\nu \preceq L_{\omega_2}$, mos_ν is an isomorphism, Lemma 2.5 and standard absoluteness results, it's not hard to verify that $\text{mos}_\nu(C) = \{X_\xi \cap \alpha_\nu : \xi < \alpha_\nu\}$.

Of course α_η is uncountable in $L_{\delta(\eta)}$, whereas it is countable in $L_{f(\alpha_\eta)}$. This means that $\delta(\eta) < f(\alpha_\eta)$, thus $\text{mos}_\nu(C) \in L_{\delta(\eta)} \subseteq L_{f(\alpha_\eta)}$ and (ii) is proved.

The proof of the first point is more involved. Of course $L_{f(\alpha_\eta)}$ is a model of $\text{ZFC} - \text{P}$, and we just showed that $\delta(\eta) < f(\alpha_\eta)$, whereby $L_{\delta(\eta)} \in L_{f(\alpha_\eta)}$. So, as before, we can construct inside $L_{f(\alpha_\eta)}$ a chain $\{N'_\nu\}_{\nu < \eta}$ of elementary substructures of $L_{\delta(\eta)}$: N'_0 is the smallest elementary substructure of $L_{\delta(\eta)}$; $N'_{\nu+1}$ is the smallest $N \preceq L_{\delta(\eta)}$ such that $N'_\nu \cup \{N'_\nu\} \subseteq N$; $N'_\gamma := \bigcup_{\xi < \gamma} N'_\xi$ for γ limit. By induction one can show that for all $\nu < \eta$, N'_ν is isomorphic to N_ν . Then N'_ν and N_ν must have the same transitive collapse $L_{\delta(\nu)}$, hence (also using absoluteness of mos) we get

$$\langle L_{\delta(\nu)} : \nu < \eta \rangle = \langle \text{mos}'' N'_\nu : \nu < \eta \rangle \in L_{f(\alpha_\eta)}.$$

In turn, we obtain

$$\langle \alpha_\nu : \nu < \eta \rangle = \langle (\omega_1)^{L_{\delta(\nu)}} : \nu < \eta \rangle \in L_{f(\alpha_\eta)},$$

as wanted. ■

Observe that

$$X \cap \alpha_\eta = \{\alpha_\nu \mid \nu < \eta \text{ and } \alpha_\nu \notin X_\nu \cap \alpha_\eta\}.$$

By last claim, this is a (absolute) definition with parameters in $L_{f(\alpha_\eta)}$. Since $L_{f(\alpha_\eta)}$ is a model of $\text{ZFC} - \text{P}$, we obtain $X \cap \alpha_\eta \in L_{f(\alpha_\eta)}$. □

3.3 Forcing and Kurepa trees

3.3.1 A forcing model where there is a Kurepa tree

In this section we will show the relative consistency of Kurepa's hypothesis. The result was developed by D. Stewart [19] in his masters' thesis (1966).

First, we recall the Delta System Lemma in its general form. The reader will have certainly seen the weaker version, which is often used to show that a certain poset is ccc. For a proof, see [11].

3.5. Delta System Lemma. *Let $\kappa \geq \omega$ a cardinal. Let $\theta > \kappa$ be regular and such that $\forall \alpha < \theta [|\alpha^{<\kappa}| < \theta]$. Suppose \mathcal{A} is a family of sets such that $|\mathcal{A}| \geq \theta$ and $\forall x \in \mathcal{A} [|x| < \kappa]$. Then there is $\mathcal{B} \subseteq \mathcal{A}$ such that $|\mathcal{B}| = \theta$ and \mathcal{B} is a Δ -system, i.e. there exists a set r such that $\forall x, y \in \mathcal{B} [x \neq y \Rightarrow x \cap y = r]$.*

Since the statement $\forall \alpha < \omega_2 [|\alpha^{<\omega_1}| < \omega_2]$ is true under CH, we obtain:

Corollary 3.6. *Assume CH. If \mathcal{A} is a family of countable sets which has size \aleph_2 , then there exists $\mathcal{B} \subseteq \mathcal{A}$ such that $|\mathcal{B}| = \aleph_2$ and \mathcal{B} is a Δ -system.*

Theorem 3.7 (Stewart). *There is a generic extension of V in which there exists a Kurepa tree.*

Proof. We assume that CH holds in V . The forcing poset is the set P which consists of all the pairs (T, f) where T is a tree from Example 1.2 and $f: S \rightarrow \mathfrak{B}_T$, with $S \subseteq \omega_2$ countable and $\mathfrak{B}_T := \{\mathfrak{b} : \mathfrak{b} \text{ is a cofinal branch in } T\}$. We stipulate that $(T_1, f_1) \leq (T_2, f_2)$ iff T_1 end-extends T_2 and f_1 *refines* f_2 , that is

$$\text{dom}(f_1) \supseteq \text{dom}(f_2) \quad \text{and} \quad f_1(\alpha) \supseteq f_2(\alpha) \text{ for all } \alpha \in \text{dom}(f_2).$$

Let G be P -generic. To lighten the notation, set $G_{\text{tr}} := \{T : (T, f) \in G \text{ for some } f\}$ and $G_{\text{fun}} := \{f : (T, f) \in G \text{ for some } T\}$. Now define

$$\mathcal{T} := \bigcup G_{\text{tr}}$$

and

$$\begin{aligned} \mathcal{F} &:= \{(\alpha, \mathfrak{b}) : \alpha \in \text{dom}(f) \text{ for some } f \in G_{\text{fun}} \text{ and} \\ &\quad \mathfrak{b} = \bigcup \{f(\alpha) : f \in G_{\text{fun}} \text{ is such that } \alpha \in \text{dom}(f)\}\}. \end{aligned}$$

Claim 1:

1. $V[G] \models \mathcal{T}$ is a tree with $\text{height}(\mathcal{T}) = \omega_1$ and only countable levels.
2. $V[G] \models \mathcal{F}: \omega_2 \rightarrow \mathfrak{B}_{\mathcal{T}}$ and \mathcal{F} is injective.

Proof: The fact that \mathcal{T} is a tree with height at most ω_1 and only countable levels is trivial. Showing that $\text{height}(\mathcal{T})$ is precisely ω_1 is a straightforward generalization of the argument we used for the same claim in the proof of Theorem 2.11 (here we must deal also with the f 's, but it's very easy to figure the way out).

For the second point, first observe that \mathcal{F} is easily seen to be a function since G is a filter. The fact that $\text{ran}(\mathcal{F}) \subseteq \mathfrak{B}_{\mathcal{T}}$ is trivial. It remains to check that $\text{dom}(\mathcal{F}) = \omega_2$ and \mathcal{F} is injective. To see this, we will show that

$$D_\alpha := \{(T, f) : \alpha \in \text{dom}(f)\} \quad \text{and} \quad E_{\alpha, \beta} := \{(T, f) : \alpha, \beta \in \text{dom}(f) \text{ and } f(\alpha) \neq f(\beta)\}$$

are dense for each $\alpha, \beta < \omega_2$ with $\alpha \neq \beta$. Obviously it suffices to show that $E_{\alpha, \beta}$ is dense for each $\alpha, \beta < \omega_2$ distinct. Let $(T, f) \in P$. We want to find $(T', f') \in E_{\alpha, \beta}$ which extends (T, f) .

CASE A: If $\text{height}(T)$ is a successor ordinal, simply extend T the usual way to a tree T' which has one more level and refine f to f' in the obvious way. In particular, if $\alpha, \beta \in \text{dom}(f)$ and $f(\alpha) = f(\beta)$, then assign $f'(\alpha)$ and $f'(\beta)$ to two distinct branches of T' that extend $f(\alpha)$, which is possible because every cofinal branch of T splits into two – actually infinitely-many – cofinal branches of T' . Using the last observation, it's immediate to extend (T, f) even in case one or both of α, β are not in $\text{dom}(f)$.

CASE B: If $\text{height}(T)$ is limit, then by our usual construction (see Lemma 2.8) we extend

T to T' which has one more level, additionally making sure that each branch in $\text{ran}(f)$ gets extended (this is possible because $\text{ran}(f)$ is countable). Then we refine f in the obvious way to a function f' with the same domain. So we obtain an extension (T', f') where $\text{height}(T')$ is a successor ordinal. We now apply CASE A to (T', f') and obtain $(T'', f'') \in E_{\alpha, \beta}$ which extends (T, f) .

Using that G is generic, it follows immediately that $\text{dom}(\mathcal{F}) = \omega_2$ and \mathcal{F} is injective. \blacksquare

To conclude that \mathcal{T} is a Kurepa tree in $V[G]$, it remains to prove that $(\omega_1)^{V[G]} = \omega_1$ and $(\omega_2)^{V[G]} = \omega_2$. The absoluteness of ω_1 follows by the fact that P is clearly countably closed. If we show that any antichain in P has size at most \aleph_1 then we can conclude that also ω_2 is preserved.

Claim 2: Let W be an antichain in P . Then $|W| \leq \aleph_1$.

Proof: Suppose towards a contradiction that $|W| = \aleph_2$. For any subset of P we use the subscripts tr and fun to indicate the relative projections, as before.

First observe that $|P_{\text{tr}}| \leq \aleph_1$. For, any tree of Example 1.2 is a countable subset of ${}^{<\omega_1}\omega$, which has power 2^{\aleph_0} , which equals \aleph_1 by CH. So $P_{\text{tr}} \subseteq [{}^{<\omega_1}\omega]^{<\omega_1}$, which has size at most \aleph_1 by CH again.

So $|W_{\text{tr}}| \leq |P_{\text{tr}}| \leq \aleph_1$, hence by $|W| \leq |W_{\text{tr}} \times W_{\text{fun}}| = \max(|W_{\text{tr}}|, |W_{\text{fun}}|)$ we get $|W_{\text{fun}}| = \aleph_2$. Observe that

$$W_{\text{fun}} \subseteq \bigcup_{\substack{T \in W_{\text{tr}} \\ X \in [\mathfrak{B}_T]^{<\omega_1}}} W_{\text{fun}}^{T, X},$$

where $W_{\text{fun}}^{T, X}$ is the set of functions in W_{fun} which are relative to the tree T (i.e. $\text{ran}(f) \subseteq \mathfrak{B}_T$) and whose range is X (which then is a countable collection of cofinal branches in T). Since any T is countable and $\mathfrak{B}_T \subseteq \mathcal{P}(T)$, using CH we get $|\mathfrak{B}_T| \leq \aleph_1$. So the set of indices we run the union over has cardinality at most \aleph_1 , thus by regularity of ω_2 there exist $T \in W_{\text{tr}}$ and $X \in [\mathfrak{B}_T]^{<\omega_1}$ such that $|W_{\text{fun}}^{T, X}| = \aleph_2$. Now:

$$W_{\text{fun}}^{T, X} \subseteq \bigcup_{\{\text{dom}(f) : f \in W_{\text{fun}}^{T, X}\}} {}^{\text{dom}(f)}X.$$

Because every X and every $\text{dom}(f)$ is countable, $|{}^{\text{dom}(f)}X| \leq \aleph_1$ by CH, whereby

$$|\{\text{dom}(f) : f \in W_{\text{fun}}^{T, X}\}| = \aleph_2.$$

By Corollary 3.6 there exists $Z \subseteq \{\text{dom}(f) : f \in W_{\text{fun}}^{T, X}\}$ such that $|Z| = \aleph_2$ and there is a root S such that $\text{dom}(f_1), \text{dom}(f_2) \in Z$ with $\text{dom}(f_1) \neq \text{dom}(f_2)$ implies $\text{dom}(f_1) \cap \text{dom}(f_2) = S$. The set $\{g \mid g : S \rightarrow X\}$ has size \aleph_1 , so there exist $f_1, f_2 \in W_{\text{fun}}^{T, X}$ distinct such that $f_1 \restriction S = f_2 \restriction S$. Also, by an easy regularity argument similar to the one above, we can assume w.l.o.g. that $\text{dom}(f_1) \neq \text{dom}(f_2)$, so that $\text{dom}(f_1) \cap \text{dom}(f_2) = S$. But then $f_1 \cup f_2$ is still a function and refines both f_1 and f_2 . So $(T, f_1 \cup f_2)$ extends (T, f_1) and (T, f_2) , which contradicts W being an antichain. \blacksquare

So \mathcal{T} is a Kurepa tree in $V[G]$, and the theorem is proved. \square

3.3.2 A forcing model where there are no Kurepa trees

In 1971, Silver produced a forcing model where no Kurepa tree exists. To do this he assumed the existence of an inaccessible cardinal in the ground model and considered the generic extension given by the Levy collapse which makes it ω_2 . The inaccessible cardinal assumption cannot be dropped: it's not hard to see that if there are no Kurepa trees, then ω_2 is inaccessible in L . Thus the existence of inaccessible cardinals is equiconsistent with the failure of Kurepa's hypothesis.

We start with an easy fact:

Lemma 3.8. *Let M be a countable transitive model for ZFC, κ a strongly inaccessible cardinal in M , P a forcing poset. Let G be generic for P over M . If $(|P| < \kappa)^M$, then κ is strongly inaccessible in $M[G]$.*

Proof. Since $|P| < \kappa$, P trivially has the κ -cc. So κ is an uncountable regular cardinal also in $M[G]$. Now let $\theta := |P|$ and take $\lambda < \kappa$ arbitrary. By inaccessibility, we obtain that there are at most $\theta^\lambda < \kappa$ nice names for subsets of $\check{\lambda}$ (cf. Preliminaries). Since every subset of λ which belongs to $M[G]$ has a (unique) corresponding nice name, it follows easily that $(|\mathcal{P}(\lambda)| < \kappa)^{M[G]}$, whereby $(2^\lambda < \kappa)^{M[G]}$. \square

This will be the key device for killing Kurepa trees:

Lemma 3.9. *Let M be a countable transitive model for ZFC. Suppose that, in M , T is an ω_1 -tree such that any level of T is countable and P is a countably closed poset. Let G be M -generic for P . If $\dot{\mathbf{b}} \in M[G]$ is a cofinal branch in T , then $\dot{\mathbf{b}} \in M$.*

Proof. In M , define $\mathfrak{B} := \{\dot{\mathbf{b}} : \dot{\mathbf{b}} \text{ is a cofinal branch in } T\}$. Assume to the contrary that $\dot{\mathbf{b}} \in M[G] \setminus M$ is a cofinal branch in T . Then there exist a name $\dot{\mathbf{b}}$ and a $p \in P$ such that

$$p \Vdash \dot{\mathbf{b}} \notin \check{\mathfrak{B}} \text{ and } \dot{\mathbf{b}} \text{ is a cofinal branch in } T.$$

We will derive that some level of T is uncountable. To show this, first we will build three sequences $\langle p_s : s \in 2^{<\omega} \rangle$, $\langle x_s : s \in 2^{<\omega} \rangle$ and $\langle \alpha_n : n \in \omega \rangle$ such that:

1. $0 = \alpha_0 < \alpha_1 < \alpha_2 < \dots < \omega_1$;
2. $p_s \in P$ and $p_s \leq p$;
3. if $\text{length}(s) = n$ then $x_s \in U_{\alpha_n}$ (the α_n th level of T);
4. $p_s \Vdash x_s \in \dot{\mathbf{b}}$;
5. $p_{s \smallfrown 0} \leq p_s$ and $p_{s \smallfrown 1} \leq p_s$;
6. $x_{s \smallfrown 0} \neq x_{s \smallfrown 1}$.

Claim: Such sequences exist.

Then let $\gamma := \sup\{\alpha_n : n \in \omega\}$, which is countable. Using (5) and the fact that P is countably closed, for any $f \in 2^\omega$ we can find p_f such that $p_f \leq p_{f \upharpoonright n}$ for all $n \in \omega$. Obviously $p_f \leq p$, so $p_f \Vdash [\dot{\mathbf{b}} \text{ is a cofinal branch in } T]$, whereby $p_f \Vdash \exists y \in U_\gamma [y \in \dot{\mathbf{b}}]$. So by Fact 2.10 there is a $q_f \leq p_f$ and an $x_f \in U_\gamma$ such that $q_f \Vdash x_f \in \dot{\mathbf{b}}$. It follows that for each $n \in \omega$, $q_f \Vdash x_{f \upharpoonright n} \in \dot{\mathbf{b}}$ (by (4)) and $\text{height}(x_{f \upharpoonright n}) = \alpha_n < \gamma$ (by (3)), so $x_{f \upharpoonright n} < x_f$. By (6), $f \neq g$ implies $x_f \neq x_g$. So $|U_\gamma| \geq 2^{\aleph_0}$, a contradiction.

Now we shall prove the claim:

Proof: We proceed by induction on $\text{length}(s)$, where $s \in 2^{<\omega}$. First, since $p \Vdash [\dot{\mathbf{b}} \text{ is a branch in } T]$, we have that by an application of Fact 2.10 there exist $p_\emptyset \leq p$ and $x_\emptyset \in U_0$ such that $p_\emptyset \Vdash x_\emptyset \in \dot{\mathbf{b}}$.

Now suppose that we have defined α_n and all p_s, x_s for $\text{length}(s) = n$. For all $s \in 2^{<\omega}$ with $\text{length}(s) = n$, define

$$E_s := \{y \in T : \exists q \leq p_s [q \Vdash y \in \dot{\mathbf{b}}]\}$$

Of course $x_s \in E_s$. Observe that:

- (i) E_s is a *subtree* of T , i.e. $x \in E_s$ implies $\downarrow x \subseteq E_s$;
- (ii) $E_s \cap T|(\alpha_n + 1) = \downarrow x_s$;
- (iii) For all $y \in E_s$, if $\text{height}(y) < \beta < \omega_1$ then there exists $z \in E_s \cap U_\beta$.

The first point is immediate to verify, using that $p_s \leq p \Vdash [\dot{\mathbf{b}} \text{ is a branch in } T]$. As for point (ii), the \supseteq inclusion follows by the first point, and the other inclusion is an immediate consequence of $p_s \Vdash [x_s \in \dot{\mathbf{b}}]$. The last point follows easily by Fact 2.10 and the fact that a witness $q \in P$ for $y \in E_s$ is such that $q \Vdash [y \in \dot{\mathbf{b}} \text{ and } \dot{\mathbf{b}} \text{ is a cofinal branch in } T]$.

E_s cannot be a cofinal branch in T (and thus, by (iii), not even just a branch). If it were, this would mean $E_s \in \mathfrak{B}$. But $p_s \Vdash \dot{\mathbf{b}} \notin \mathfrak{B}$, whereby $p_s \Vdash \dot{\mathbf{b}} \neq E_s$, so $p_s \Vdash \dot{\mathbf{b}} \not\subseteq E_s$ (because they are both cofinal branches, and thus maximal). So there would be $y \in T \setminus E_s$ and $q \leq p_s$ such that $q \Vdash y \in \dot{\mathbf{b}}$, which contradicts the definition of E_s .

Of course there are only finitely-many E_s 's, since the binary sequences of length n are finitely many. Because the E_s are not branches, using (iii) we can find $\alpha_{n+1} > \alpha_n$ (fixed) such that for each s of length n there are $x_{s \smallfrown 0}, x_{s \smallfrown 1} \in E_s \cap U_{\alpha_{n+1}}$ with $x_{s \smallfrown 0} \neq x_{s \smallfrown 1}$. By definition of E_s we obtain also $p_{s \smallfrown 0}$ and $p_{s \smallfrown 1}$ as wanted. ■

□

Definition 3.10. Let α be an ordinal. We define $\text{Fn}_{\omega_1}(\omega_1, \alpha) := \{f \mid f: S \rightarrow \alpha \text{ and } S \subseteq \omega_1 \text{ with } |S| < \omega_1\}$.

The following corollary is not strictly necessary for our aim in this section, but it will help understanding the proof of the main theorem.

Corollary 3.11. *Let M and T be as in last lemma. In M , let $\theta \geq 2^{\aleph_1}$ and $P := \text{Fn}_{\omega_1}(\omega_1, \theta)$. If G is M -generic for P , then T is not a Kurepa tree in $M[G]$.*

Proof. Since P is countably closed, $(\omega_1)^{M[G]} = (\omega_1)^M$. Moreover, observe that $\bigcup G$ is a surjective function $(\omega_1)^{M[G]} \rightarrow \theta$, so that $M[G] \models |\theta| \leq \aleph_1$. Define \mathfrak{B} as before. Of course $|\mathfrak{B}| \leq \theta$ in M , because $\mathfrak{B} \subseteq \mathcal{P}(T)$. So, because by last lemma in $M[G]$ there are no new branches of T , we get $M[G] \models |\mathfrak{B}| \leq |\theta| \leq \aleph_1$. Thus, in $M[G]$, T has at most \aleph_1 -many uncountable branches, i.e. it's not a Kurepa tree. \square

We would like to claim that in $M[G]$ there are no Kurepa trees, but this is not quite true. While no tree in M is a Kurepa tree in $M[G]$, new Kurepa trees might have come up in the generic extension. To address this issue we will need to use product forcing, but it will be necessary to assume the existence of a strongly inaccessible cardinal.

Theorem 3.12 (Silver). *Let M be a countable transitive model for ZFC. Suppose that $(\kappa$ is strongly inaccessible) M . Then there is a generic extension $M[G]$ such that $M[G] \models \kappa = \omega_2$ and $M[G] \models$ “there are no Kurepa trees”.*

Proof. Working in M , suppose that $S \subseteq \kappa$ and define

$$P_S := \left\{ p \in \prod_{\alpha \in S} \text{Fn}_{\omega_1}(\omega_1, \alpha) : |\{\alpha : p_\alpha \neq 1\}| \leq \aleph_0 \right\}.$$

Define also $P := P_\kappa$. Observe that clearly $P \simeq P_S \times P_{\kappa \setminus S}$ for all $S \subseteq \kappa$.

Let G be P -generic over M . Since κ is strongly inaccessible, the Δ -system Lemma 3.5 applies. An argument similar to the one in Theorem 3.7 shows that P has the κ -cc. Hence cofinalities and cardinals $\geq \kappa$ are preserved. Moreover, it's easy to show that P is countably closed, whereby $(\omega_1)^{M[G]} = (\omega_1)^M$.

Claim: $(\omega_2)^{M[G]} = \kappa$.

Proof: Of course $(\omega_2)^{M[G]} \leq \kappa$ because cofinalities are preserved. For the other inequality, take $\alpha < \kappa$ arbitrary. There is an obvious embedding (defined in M) $i : (\text{Fn}_{\omega_1}(\omega_1, \alpha))^M \rightarrow P$ which is complete. It follows easily⁴ that the preimage $i^{-1}[G] \in M[G]$ and that $i^{-1}[G]$ is $(\text{Fn}_{\omega_1}(\omega_1, \alpha))^M$ -generic over M . Its union is a surjective map from $(\omega_1)^{M[G]}$ onto α , so that $(\alpha < \omega_2)^{M[G]}$. \blacksquare

Suppose towards a contradiction that in $M[G]$ there is a Kurepa tree, which we may assume w.l.o.g. to be (ω_1, \sqsubset) , where $\sqsubset \subseteq \omega_1 \times \omega_1$ is a tree order. Let $\dot{\sqsubset}$ be a nice name for a subset of $(\omega_1 \times \omega_1)$. By the result about nice names mentioned in the preliminaries, we can assume that $\dot{\sqsubset}_G = \sqsubset$. Note that this is possible also because $M[G] \models \sqsubset \subseteq \omega_1 \times \omega_1$ and $(\omega_1 \times \omega_1)_G = \omega_1 \times \omega_1 = \omega_1^{M[G]} \times \omega_1^{M[G]}$ since ω_1 is preserved. So

$$\dot{\sqsubset} = \bigcup \{ \{ \sigma \} \times A_\sigma : \sigma \in \text{dom}((\omega_1 \times \omega_1)) \},$$

where the A_σ 's are all antichains of P (in M). Now let

$$B := \bigcup \{ A_\sigma : \sigma \in \text{dom}((\omega_1 \times \omega_1)) \}.$$

⁴See [14, Lemma IV.4.2, p. 270].

Since every A_σ has size $< \kappa$ and $|\text{dom}((\omega_1 \check{\times} \omega_1))| = |\omega_1 \times \omega_1| = \aleph_1 < \kappa$, by regularity of κ we have $|B| < \kappa$. Define

$$S := \bigcup \{\text{supt}(p) : p \in B\},$$

where $\text{supt}(p) := \{\alpha : p_\alpha \neq 1\}$, which is always countable. So $|S| < \kappa$ by regularity again. Hence, M models that

$$(3.2) \quad \forall \gamma < \kappa \exists \alpha < \kappa [\alpha > \gamma \text{ and } \alpha \notin S].$$

Now define $P^- := P_S$ and $P^+ := P_{\kappa \setminus S}$, so that $P \simeq P^- \times P^+$. By the Product Lemma, we have that $G = G^- \times G^+$, where G^- is P^- generic over M and G^+ is P^+ -generic over $M[G^-]$. Also, $M[G] = M[G^-][G^+]$. Furthermore, the definition of S ensures that $\square \in M[G^-]$. Observe that P^- is countably closed, thus it doesn't add ω -sequences. Since P^+ is countably closed in M , this implies that P^+ is countably closed in $M[G^-]$ as well. Therefore we can apply Lemma 3.9 to conclude that every cofinal branch in (ω_1, \square) that lies in $M[G]$ is already an element of $M[G^-]$. In $M[G]$ (equivalently, in $M[G^-]$), let

$$\mathfrak{B} := \{\mathfrak{b} : \mathfrak{b} \text{ is a cofinal branch in } (\omega_1, \square)\}.$$

Claim : $(|P^-| < \kappa)^M$.

Proof: Observe that, for each $\alpha < \kappa$, $|\text{Fn}_{\omega_1}(\omega_1, \alpha)| \leq |\alpha^{\omega_1}| < \kappa$ by inaccessibility. Moreover, since κ is regular and $|S| < \kappa$, there is $\beta < \kappa$ greater than any element of S , so that $|\text{Fn}_{\omega_1}(\omega_1, \alpha)| < |\text{Fn}_{\omega_1}(\omega_1, \beta)|$ for all $\alpha \in S$. So $|P^-| \leq |^S(\text{Fn}_{\omega_1}(\omega_1, \beta))| < \kappa$ by inaccessibility again. ■

Using Lemma 3.8, by last claim we obtain that κ stays strongly inaccessible in $M[G^-]$. Clearly $|\mathfrak{B}| \leq 2^{\aleph_1}$ because $\mathfrak{B} \subseteq \mathcal{P}(\omega_1)$. So $|\mathfrak{B}| < \kappa$ by inaccessibility, and thus there is $|\mathfrak{B}| < \alpha < \kappa$. Of course (3.2) holds also in $M[G^-]$ by absoluteness. So we can assume that $\alpha \notin S$. This implies that $\text{Fn}_{\omega_1}(\omega_1, \alpha)$ is a factor of the product which defines P^+ . By repeating the argument in the proof of the first claim, we obtain that in $M[G]$

$$|\mathfrak{B}| \leq |\alpha| \leq \omega_1,$$

so (ω_1, \square) is not Kurepa in $M[G]$. □

Chapter 4

Automorphisms of trees

In this chapter we will present some less-known results concerning automorphisms of trees. Most of them are due to Jensen, who proved them in some lecture notes written in 1969. The reference we used is Devlin's book [2], which was published in 1974 and is based on those notes.

Definition 4.1. An *isomorphism* of two trees (T_1, \leq_1) and (T_2, \leq_2) is an isomorphism of ordered sets, i.e. a bijection $\sigma: T_1 \rightarrow T_2$ such that for all $x, y \in T_1$, $x \leq_1 y$ iff $\sigma(x) \leq_2 \sigma(y)$. An *automorphism* of a tree is an isomorphism with itself.

Definition 4.2. A tree T is *homogeneous* if for all $x, y \in T$ such that $o(x) = o(y)$, there exists an automorphism σ of T such that $\sigma(x) = y$ and $\sigma(y) = x$.

Which trees are homogeneous? First, we prove that every countable normal tree is homogeneous. To do this, we will first show that any countable normal tree is completely determined by its height, up to isomorphism. This is a very easy result, but note that it is not completely trivial as it might seem at a first look. In fact, a naive approach would generalize to normal trees of height ω_1 . But of course not all of these are isomorphic, as, for example, some of them are Aronszajn and some are not (Theorem 1.7 and Example 1.4).

Proposition 4.3. *If T_1 and T_2 are normal α -trees with $\alpha < \omega_1$, then they are isomorphic.*

Proof. This is proven by induction on α . The only non-trivial case is when α is limit. First take $\{\mathfrak{a}_n\}_{n \in \omega}$ and $\{\mathfrak{b}_m\}_{m \in \omega}$ countable collections of α -branches of T_1 and T_2 respectively, such that $T_1 = \bigcup_{n \in \omega} \mathfrak{a}_n$ and $T_2 = \bigcup_{m \in \omega} \mathfrak{b}_m$. Such collections exist for any countable normal tree, because applying Lemma 2.8 with $\mathcal{A} := \emptyset$ we get a normal end-extension with one more level (which is countable), and the family given by the sets of predecessors of every node in the new level is the wanted collection of cofinal branches.

Now we define $\sigma_n: \mathfrak{a}_n \rightarrow T_2$ by induction on $n < \omega$. We assume inductively that the σ_i 's defined previously are coherent and σ_i is always an order-isomorphism of \mathfrak{a}_i with some (unique) \mathfrak{b}_{m_i} . Now let's define σ_n . Of course we don't "touch" σ_n on previously visited

nodes; formally

$$\sigma_n \upharpoonright \left(\mathfrak{a}_n \cap \bigcup_{i < n} \mathfrak{a}_i \right) := \bigcup_{i < n} \sigma_i \upharpoonright \left(\mathfrak{a}_n \cap \bigcup_{i < n} \mathfrak{a}_i \right),$$

thus σ_n is coherent with every other σ_i .

Now let x be the least node of $\mathfrak{a}_n \setminus \bigcup_{i < n} \mathfrak{a}_i$. By normality, $o(x)$ is a successor ordinal. Let v be its immediate predecessor. Consider the immediate successors of $\sigma_n(v)$. Any such node lies on some \mathfrak{b}_m . We assign the remaining part of \mathfrak{a}_n to \mathfrak{b}_m with m minimal, in the obvious order-preserving way. Of course $\sigma_n[\mathfrak{a}_n] = \mathfrak{b}_m$, so that the induction hypothesis is preserved.

It is straightforward to verify that $\bigcup_{n \in \omega} \sigma_n$ is an isomorphism $T_1 \rightarrow T_2$. \square

Theorem 4.4. *Every countable normal tree T is homogeneous.*

Proof. We proceed by induction on $\alpha := \text{height}(T)$. For $\alpha = 0$ it's trivial. Suppose that the theorem is true for every $T|_\beta$ with $\beta < \alpha$. Take $x, y \in T$ at the same level γ . We consider two cases:

CASE A: $\gamma = \delta + 1$ for some δ . Then let x' and y' be the immediate predecessors of x and y respectively. Of course $x', y' \in T|_\gamma$. By the induction hypothesis, there is some automorphism σ' of $T|_\gamma$ which swaps x' and y' . Now, for any $z \in U_\gamma$ (the γ th level of T), consider $T_z := \{t \in T \mid t \geq z\}$. All the T_z 's are trivially countable normal trees of the same height, so they are all isomorphic by Proposition 4.3. Of course then we can define an automorphism τ of $\bigcup_{z \in U_\gamma} T_z$ which induces a permutation of $\{T_z\}_{z \in U_\gamma}$ such that:

- (i) $\tau[T_x] = T_y$ and $\tau[T_y] = T_x$;
- (ii) If $z \in U_\gamma$ and z' is its immediate predecessor, then $\tau[T_z] = T_{\tau(z)}$ where $\tau(z)$ is an immediate successor of $\sigma'(z')$.

Now define $\sigma := \sigma' \cup \tau$. Condition (ii) ensures that σ is indeed an automorphism of T , and by (i) it clearly swaps x and y .

CASE B: γ is limit. Then consider $\mathfrak{d}_0 := \downarrow x$ and $\mathfrak{d}_1 := \downarrow y$. Now repeat the proof of Proposition 4.3, setting $T_1 = T_2 = T$, $\mathfrak{a}_0 = \mathfrak{b}_1 = \mathfrak{d}_0$ and $\mathfrak{a}_1 = \mathfrak{b}_0 = \mathfrak{d}_1$. If we look closely at that proof, it's immediate to see that, in our case, σ_0 maps \mathfrak{a}_0 to \mathfrak{b}_0 and σ_1 maps \mathfrak{a}_1 to \mathfrak{b}_1 , and this is of course true also for the final isomorphism. Then the same argument gives us an automorphism σ' of $T|_\gamma$ which maps \mathfrak{d}_0 to \mathfrak{d}_1 and vice versa. Moreover, let $\{t_i : i < \omega\}$ be an enumeration of the γ th level of T , with $t_0 = x$ and $t_1 = y$. Define $\mathfrak{d}_i := \downarrow t_i$. Then the automorphism σ' acts as a permutation of $\{\mathfrak{d}_i : i < \omega\}$. If we extend σ' by mapping every t_i to the t_j such that $\sigma'[\mathfrak{d}_i] = \sigma'[\mathfrak{d}_j]$, then we obtain an automorphism σ'' of $T|_{(\gamma+1)}$ which swaps $t_0 = x$ and $t_1 = y$.

Finally, the trees of the form T_{t_i} are all isomorphic, so σ'' can be easily extended to an automorphism of T by permutating them according to σ'' as we did in CASE A. \square

At this point, it's natural to ask whether the result holds also for normal ω_1 -tree. This is however not true: assuming \diamond , we can construct both a normal Suslin tree which

is homogeneous and one which is *rigid* (i.e. it has no non-trivial automorphisms). The following two sections will respectively show how to produce them.

On a side note, we remark that ZFC is sufficient to prove the existence of both homogeneous and rigid normal ω_1 -trees. Even more: we can have both the homogeneous and the rigid trees be Aronszajn. To construct the former is rather easy. The latter is a non-trivial problem, raised by Jech in 1972 [7] and independently solved by Uri Abraham [1] and Todorcevic [21], both in 1979 (that is, after Jensen constructed the homogeneous and rigid Suslin trees using \diamond).

4.1 A homogeneous Suslin tree

First we introduce a slightly modified version of normal trees, which turns out to be more appropriate for the scope of this section.

Definition 4.5. A *binary normal tree* is a normal tree as in Definition 1.3, but we require that each node has only two immediate successors instead of infinitely many.

In the remaining part of this chapter, by normal we mean binary normal.

To show that $\diamond \Rightarrow$ there exists a homogeneous Suslin tree, we will need to adapt Lemma 2.5. The reason is that we always used 2.5 in combination with the hypothesis $V = L$, so that $L_\kappa = H_\kappa$ for all infinite cardinals. But our proof of existence of a homogeneous ST will assume only \diamond . So we need to somehow adapt 2.5 to “natively” deal with L_θ ’s. The solution is simply to relativize the whole statement to L . Together with some easy absoluteness arguments, the result is the following:

Lemma 4.6. *Let θ be a regular uncountable cardinal in L . Suppose that M is a structure such that $M \preceq L_\theta$ and $(M \text{ is countable})^L$. Then the following hold:*

1. *If $a \in M$ and $(a \text{ is countable})^L$, then $a \subseteq M$.*
2. *$M \cap \omega_1^L = \beta$, where $\beta < \omega_1^L$ and β is a limit ordinal.*
3. *If $\theta > \omega_1^L$, then $\omega_1^L \not\subseteq M$ but $\omega_1^L \in M$, and “ ω_1^L is the first uncountable ordinal” is true in M .*
4. *Let $T := \text{mos}'' M$ and let $\beta = M \cap \omega_1^L$ by the second point. Then $\text{mos}(\omega_1^L) = \beta$ and $\text{mos}(\xi) = \xi$ for all $\xi < \beta$.*
5. *$\beta = (\omega_1)^T$.*

Proof. The statement clearly follows by Lemma 2.5 plus the fact that $L \models V = L$ and $V = L$ implies $H_\kappa = L_\kappa$ for all cardinals $\kappa \geq \omega$. Nevertheless, we decide to prove the first point from scratch in order to illustrate a kind of argument which will come up often hereafter. We will refer to this strategy as “usual methods”. The main observation is essentially the following: if L believes that a certain set belongs to H_κ (which is *not* absolute), then it belongs to L_κ , which *is* absolute between V and L , so that $x \in L_\kappa$ also

in V . Let's now proceed with the proof of (1):

For simplicity, we assume that $a \subseteq \omega_1^L$ (this assumption will always be true when we will use this point; anyway, to adapt the proof to the general case is straightforward). It's easy to see that ω and all $n \in \omega$ are definable in L_θ , so $\omega \in M$ and $\omega \subseteq M$. Now, if $a = \emptyset$ then trivially $a \subseteq M$. If $a \neq \emptyset$, then in L there exists $f: \omega \rightarrow a$ surjective. It's immediate to check that any such function is L -hereditarily countable, i.e. belongs to $(H_{\omega_1})^L$. Since $V = L \Rightarrow L_{\omega_1} = H_{\omega_1}$ by Lemma 2.6, we have that $(f \in H_{\omega_1})^L \Rightarrow (f \in L_{\omega_1})^L$, i.e. $f \in L_{\omega_1^L} \subseteq L_\theta$. Since $M \preceq L_\theta$, at least one such f belongs to M as well. Observe that $f, n \in M$ implies $f(n) \in M$ (by $M \preceq L_\theta$ again), and since $a = \{f(n) : n \in \omega\}$ we have $a \subseteq M$, so (1) is proved. \square

Remark. It's easy to see that Lemma 4.6 can be generalized in the obvious way to $L[A]$, where $A \subseteq \omega_1$.

Before proceeding with the main theorem, two trivial lemmas:

Lemma 4.7. *Let $A \subseteq \text{Ord}$. Then $L_\alpha[A] = L_\alpha[A \cap \alpha]$ for every ordinal α .*

Proof. An easy induction. \square

Lemma 4.8. *Let A be a set. Then $L[A \cap \beta] \subseteq L[A \cap \gamma]$ for all $\beta, \gamma \in \text{Ord}$ with $\beta \leq \gamma$.*

Proof. Immediate by the characterization of $L[X]$ as the smallest inner model which contains X . \square

Recall that the Diamond Principle (\diamond) asserts:

There exists a sequence of sets $\langle S_\alpha : \alpha < \omega_1 \rangle$ with $S_\alpha \subseteq \alpha$, such that for every $X \subseteq \omega_1$, the set $\{\alpha < \omega_1 : X \cap \alpha = S_\alpha\}$ is a stationary subset of ω_1 .

We can now present the main result of this section:

Theorem 4.9. *If \diamond holds, then there exists a homogeneous normal Suslin tree.*

Proof. Assume \diamond and observe that $|H_{\omega_1}| = 2^{<\omega_1} = \omega_1$, because \diamond implies CH. This means that H_{ω_1} can be “coded” by some $A \subseteq \omega_1$, so that $H_{\omega_1} = L_{\omega_1}[A]$.¹ In particular, every bijection between ω and a countable ordinal belongs to $L[A]$. This clearly implies $\omega_1^{L[A]} = \omega_1$.

Now let $\langle S_\alpha \mid \alpha < \omega_1 \rangle$ be the sequence given by \diamond . We define² by recursion on $\alpha < \omega_1$ the function $\alpha \mapsto \delta_\alpha$ as follows: δ_α is the least ordinal $\delta > \alpha$ such that

- (i) $L_\delta[A] \prec H_{\omega_1}$;
- (ii) $S_\alpha, \langle \delta_\nu \mid \nu < \alpha \rangle \in L_\delta[A]$.

Set $M_\alpha := L_{\delta_\alpha}[A]$. Of course each M_α is countable.

To continue the proof, we need an important claim which we will prove later:

¹For an exhaustive treatment of this topic, see [9], section 13.

²The function is well defined by an argument very similar to the one in the proof of Theorem 3.4.

Claim 4.10. *Let T be a normal ω_1 -tree with $T \subseteq H_{\omega_1}$ and let $C \subseteq \omega_1$ be a club set of limit ordinals such that for all $\alpha \in C$ the following holds:*

If $T|_\alpha \in M_\alpha$ and x belongs to the α th level of T , then $\downarrow x$ is M_α -generic³ for $T|_\alpha$.

Then T is a Suslin tree.

We proceed with the construction of a homogeneous Suslin tree T , which we define by induction on levels U_α as usual. T will be a set of countable binary sequences s such that $o(s) = \text{length}(s)$, ordered by \subseteq . By induction, it will also follow that for every $\alpha < \omega_1$:

- (a) $T|_\alpha$ is normal;
- (b) $T|_\alpha \in M_\alpha$;
- (c) if $s, t \in T|_\alpha$ and $o(s) = o(t)$, then the set $\{\xi \mid s(\xi) \neq t(\xi)\}$ is finite.

Of course we define $U_0 := \{\emptyset\}$ and

$$(4.1) \quad U_{\alpha+1} := \{s \smallfrown i \mid s \in T_\alpha \text{ and } i \in 2\}.$$

The inductive hypotheses (a), (b) and (c) are trivially preserved here. So it is left to define U_α for α limit. Let \mathfrak{b} be the least (w.r.t. the canonical well-order of $L[A]$) M_α -generic branch of $T|_\alpha$. Such a branch exists by the Generic Filter Existence Lemma, because $(L[A]$ models that) M_α is countable. Define

$$(4.2) \quad U_\alpha := \left\{ \bigcup \mathfrak{a} \mid \mathfrak{a} \text{ is an } \alpha\text{-branch through } T|_\alpha, \text{ and } \bigcup \mathfrak{a} \text{ and } \bigcup \mathfrak{b} \text{ differ only on finitely-many entries} \right\}.$$

To preserve the normality, we need to check that U_α is countable and for every $x \in T|_\alpha$ there is $y \in U_\alpha$ such that $x < y$. The first claim is immediate: every sequence $\bigcup \mathfrak{a}$ in the definition of U_α is completely determined by $\bigcup \mathfrak{b}$, except for finitely-many entries. Since there are at most \aleph_0 -many such sequences, U_α is countable. As for the second condition, observe that if $x \in T|_\alpha$, then there is an α -branch \mathfrak{a} of $T|_\alpha$ such that $x \in \mathfrak{a}$ and, above x , \mathfrak{a} has exactly the same “shape” of \mathfrak{b} . More precisely

If $x \in T|_\alpha$, then $\{x \restriction \xi : \xi \leq o(x)\} \cup \{x \smallfrown (b_\xi \restriction [o(x) + 1, \xi)) : o(x) + 1 \leq \xi < \alpha\}$ is an α -branch in $T|_\alpha$, where b_ξ is the only node of \mathfrak{b} at level ξ .

(this is proven straightforwardly by induction showing that $x \smallfrown (b_\xi \restriction [o(x) + 1, \xi)) \in T|_\alpha$ for all $\xi \in [o(x) + 1, \alpha)$, using that $T|_\alpha$ was constructed according to (4.1) and (4.2)) Since $\bigcup \mathfrak{a}$ and $\bigcup \mathfrak{b}$ differ only on $\text{dom}(x)$, by the induction hypothesis (c) they differ only on a finite set. So $\bigcup \mathfrak{a} \in T|_\alpha$ and clearly $x < \bigcup \mathfrak{a}$. Hence (a) is preserved. Conditions (b) and (c) are also easily seen to be preserved.

Finally, define $T := \bigcup_{\alpha < \omega_1} U_\alpha$. Of course conditions (a) and (c) are still true with T in

³Of course here we mean generic w.r.t. the reverse order (but obviously $\downarrow x$ is referred to the original order).

place of $T|_\alpha$. Now, for any finite $\Omega \subseteq \omega_1$ define $\sigma_\Omega: 2^{<\omega_1} \rightarrow 2^{<\omega_1}$ as the function which swaps the entries in each point of Ω . Formally: $\text{dom}(\sigma_\Omega(s)) = \text{dom}(s)$ and

$$\sigma_\Omega(s)(\xi) := \begin{cases} s(\xi) & \text{if } \xi \notin \Omega \\ 1 - s(\xi) & \text{if } \xi \in \Omega. \end{cases}$$

It follows directly from its definition that T is closed under the map σ_Ω for all finite $\Omega \subseteq \omega_1$. Hence, if $x, y \in T$ and $\Omega := \{\xi < \omega_1 \mid x(\xi) \neq y(\xi)\}$ (which is finite), then σ_Ω is an automorphism of T such that $\sigma_\Omega(x) = y$ and $\sigma_\Omega(y) = x$. That is, T is homogeneous.

Finally, note that if α is limit, then every α -branch which is extended in the construction of T is M_α -generic for $T|_\alpha$. To see this, let \mathfrak{a} be such a branch and take \mathfrak{b} the least M_α -generic branch of $T|_\alpha$. By definition of T , there is a finite $\Omega \subseteq \omega_1$ such that $\mathfrak{b} = \sigma_\Omega[\mathfrak{a}]$. Let $D \in M_\alpha$ be a dense subset of $T|_\alpha$ and suppose towards a contradiction that $D \cap \mathfrak{a} = \emptyset$. But then $\emptyset = \sigma_\Omega[D \cap \mathfrak{a}] = \sigma_\Omega[D] \cap \mathfrak{b}$, which is impossible because $\sigma_\Omega[D]$ is dense (and belongs to M_α because Ω is definable in M_α and thus the automorphism σ_Ω belongs to M_α). Therefore Claim 4.10 applies and so T is Suslin. \square

We will now prove Claim 4.10. Some parts of the following proof are very similar to the proof of Theorem 3.4, which showed that there exists a Kurepa family if $V = L$. We will occasionally refer to that proof to justify some steps.

Proof of Claim 4.10. Let X be a maximal antichain in T . If we show that X is countable, then T is Suslin by normality. Let $B \subseteq \omega_1$ be such that $A, C, T, X \in L[B]$ (it's easy to see that such a B exists). Observe that by $\omega_1^{L[A]} = \omega_1$ and $A \in L[B]$ it follows that $\omega_1^{L[B]} = \omega_1$ as well. Define

$$N^* := L_{\omega_2^{L[B]}}[B].$$

We recursively define a chain of elementary substructures of (N^*, \in) , which will look like this:

$$N_0 \preceq N_1 \preceq \cdots \preceq N_\nu \preceq \cdots \preceq (N^*, \in)$$

for $\nu < \omega_1$. The chain is defined as follows:

$$(4.3) \quad \begin{aligned} N_0 & \text{ is the smallest } N \preceq N^* \text{ such that } B, A, C, T, X \in N; \\ N_{\nu+1} & \text{ is the smallest } N \preceq N^* \text{ such that } N_\nu \subseteq N \text{ and } N_\nu \in N; \\ N_\eta & \text{ is } \bigcup_{\xi < \eta} N_\xi \text{ for } \eta \text{ limit.} \end{aligned}$$

Observe that:

- every N_ν is countable;

- the $\nu = 0$ and the $\nu + 1$ steps are well defined (i.e. $B, A, C, T, X \in N^*$ and $N_\nu \in N^*$). This is proved as in the proof of Theorem 3.4, with the following additional observation: Since $B \in L[B]$, we can define N^* within $L[B]$ and the definition is absolute, i.e. $N^* = (L_{\omega_2}[B])^{L[B]}$. Hence we can define the sequence $\langle N_\nu \rangle_{\nu < \omega_1}$ within $L[B]$ (and it is absolute). Since $(V = L[B])^{L[B]}$, we can apply the usual methods (cf. proof of Lemma 4.6) to repeat the argument of the proof of 3.4, which relied on the assumption $V = L$.
- for each $\nu < \omega_1$, by Lemma 4.6 we have $\omega_1 \cap N_\nu = \alpha_\nu$ for some $\alpha_\nu < \omega_1$.

For every $\nu < \omega_1$, let $\overline{N}_\nu := \text{mos}_\nu \restriction N_\nu$ be the Mostowski collapse of N_ν . By Condensation Lemma, $\overline{N}_\nu = L_{\beta_\nu}[\text{mos}_\nu(B)] = L_{\beta_\nu}[B \cap \alpha_\nu]$ for some $\beta_\nu \leq \omega_2^{L[B]}$. Moreover, observe that

$$\text{mos}_\nu \restriction L_{\alpha_\nu}[B \cap \alpha_\nu] = \text{id} \restriction L_{\alpha_\nu}[B \cap \alpha_\nu].$$

This is true because $L_\gamma[B] \subseteq N_\nu$ for every $\gamma \leq \alpha_\nu$, which is proven by induction on γ . But $\text{mos}_\nu(L_{\omega_1}[B]) = L_{\text{mos}_\nu(\omega_1)}[\text{mos}_\nu(B)] = L_{\alpha_\nu}[B \cap \alpha_\nu]$ (observe that $L_{\omega_1}[B] \in N_\nu$ by elementarity because $L_{\omega_1}[B]$ is definable in N^*). So $N_\nu \cap L_{\omega_1}[B] = L_{\alpha_\nu}[B \cap \alpha_\nu]$ and hence, if $Y \in N_\nu$ and $Y \subseteq L_{\omega_1}[B]$, then $\text{mos}_\nu(Y) = Y \cap L_{\alpha_\nu}[B \cap \alpha_\nu]$.

Now observe that $C \in N^*$ by the usual methods and $(C \text{ is club in } \omega_1)^{N^*}$. Moreover, for any $\nu < \omega_1$, $C \in N_\nu$ and $C \subseteq L_{\omega_1}[B]$, so that $\text{mos}_\nu(C) = C \cap L_{\alpha_\nu}[B \cap \alpha_\nu] = C \cap \alpha_\nu$. So, by elementarity, $(C \cap \alpha_\nu \text{ is unbounded in } \alpha_\nu)^{\overline{N}_\nu}$, and of course the same holds also in V . Therefore $\alpha_\nu = \sup(C \cap \alpha_\nu) \in C$ for all $\nu < \omega_1$. Moreover, it's immediate to check that $\{\alpha_\nu \mid \nu < \omega_1\}$ is a *normal* sequence in ω_1 , i.e. continuous and strictly increasing. It is well-known that every normal function has club-many fixed points. So $\{\gamma \mid \alpha_\gamma = \gamma\}$ is a club set in ω_1 . By \diamond , the set $\{\gamma : B \cap \gamma = S_\gamma\}$ is stationary in ω_1 . So there exists an ordinal ζ such that $\alpha_\zeta = \zeta$ and $B \cap \zeta = S_\zeta$. Since $S_\zeta \in M_\zeta$, also $B \cap \zeta \in M_\zeta$.

Claim: $\beta_\zeta \in M_\zeta$.

The proof of last claim is very technical, so we will present it at the end. So $\beta_\zeta \in M_\zeta = L_{\delta_\zeta}[A]$, hence $\beta_\zeta < \delta_\zeta$. Then we obtain

$$\overline{N}_\zeta = L_{\beta_\zeta}[B \cap \zeta] \subseteq L_{\delta_\zeta}[B \cap \zeta] \subseteq M_\zeta,$$

where the last inclusion is true because in $M_\zeta = L_{\delta_\zeta}[A]$ we can re-define every $L_\gamma[B \cap \zeta]$ for $\gamma < \delta_\zeta$ (recall that $B \cap \zeta \in M_\zeta$). Observe that $T, X \in N_\zeta$ and $T, X \subseteq L_{\omega_1}[B]$, so $\text{mos}_\zeta(T) = T \restriction \zeta$ and $\text{mos}_\zeta(X) = X \cap T \restriction \zeta$. Thus

$$T \restriction \zeta \in \overline{N}_\zeta \subseteq M_\zeta \quad \text{and} \quad X \cap T \restriction \zeta \in \overline{N}_\zeta \subseteq M_\zeta.$$

Also, by elementarity $X \cap T \restriction \zeta$ is a maximal antichain in $T \restriction \zeta$. If $x \in T$ is of height ζ , then by hypothesis the branch $\downarrow x$ is M_ζ -generic for $T \restriction \zeta$. Hence it has non-empty intersection with $X \cap T \restriction \zeta$, so x lies above a point in $X \cap T \restriction \zeta$. This true for every x at level ζ , therefore $X \cap T \restriction \zeta$ is a maximal antichain in $T \restriction \zeta + 1$ as well. By Lemma 2.7 it is maximal also in T . Thus $X = X \cap T \restriction \zeta$ and so X is countable.

It remains to prove the claim:

Proof: We have to show that $\beta_\zeta \in M_\zeta$. We consider two cases.

CASE A: $\omega_1^{L[B \cap \gamma]} = \omega_1$ for some $\gamma < \omega_1$. Note that any bijection $\omega_1^{L[B \cap \gamma]} \rightarrow \omega_1$ in $L[B \cap \gamma]$ belongs to $(H_{\omega_2})^{L[B \cap \gamma]} = (L_{\omega_2}[B \cap \gamma])^{L[B \cap \gamma]} = L_{\omega_2^{L[B \cap \gamma]}}[B \cap \gamma] \subseteq L_{\omega_2^{L[B]}}[B] = N^*$.

So, if we let γ_0 be the least ordinal such that $\omega_1^{L[B \cap \gamma_0]} = \omega_1$, we have that γ_0 is definable in N^* by the usual methods. Since $N_\zeta \preceq N^*$, $\gamma_0 \in N_\zeta$ as well. Hence $\gamma_0 < \alpha_\zeta = \zeta$, so $\omega_1^{L[B \cap \zeta]} = \omega_1$ by Lemma 4.8. Since $\zeta < \omega_1$, of course (ζ is countable) $^{L[B \cap \zeta]}$. By the usual methods, an injection $\zeta \rightarrow \omega$ already lies at some countable level of the hierarchy, i.e. there is $\tau < \omega_1$ such that ζ is countable in $L_\tau[B \cap \zeta]$. Of course τ is definable in H_{ω_1} using the parameter $B \cap \zeta$. Recalling that $B \cap \zeta \in M_\zeta \prec H_{\omega_1}$ we obtain $\tau \in M_\zeta$. But $\zeta = \alpha_\zeta = \omega_1^{\overline{N}_\zeta}$, i.e. ζ is uncountable in $L_{\beta_\zeta}[B \cap \zeta]$. Thus $\beta_\zeta < \tau$, whereby $\beta_\zeta \in M_\zeta$.

CASE B: $\omega_1^{L[B \cap \gamma]} < \omega_1$ for all $\gamma < \omega_1$. First we claim that there exists no γ such that $\omega_2^{L[B \cap \gamma]} = \omega_1$. For, assume to the contrary that there is such a γ , so that $\omega_1^{L[B \cap \gamma]} < \omega_1 = \omega_1^{L[B]}$. By the usual methods, there is some $\eta < \omega_1$ such that $\omega_1^{L[B \cap \eta]}$ is countable in $L_\eta[B] = L_\eta[B \cap \eta]$, so it's clearly countable in $L[B \cap \eta]$ as well. Hence we know that

$$\omega_1^{L[B \cap \gamma]} < \omega_1^{L[B \cap \eta]} \leq \omega_1 = \omega_2^{L[B \cap \gamma]}.$$

Because $\omega_1^{L[B \cap \eta]}$ is a cardinal in $L[B \cap \eta] \supseteq L[B \cap \gamma]$, it's a cardinal also in $L[B \cap \gamma]$, whence $\omega_1^{L[B \cap \eta]} = \omega_1$, a contradiction.

Thus $\omega_2^{L[B \cap \zeta]}$ is countable, and hence is H_{ω_1} -definable in the parameter $B \cap \zeta$. So $\omega_2^{L[B \cap \zeta]} \in M_\zeta$. Moreover, observe that

$$L_{\omega_2^{L[B]}}[B] \models \text{for every ordinal } \gamma \text{ there is a surjection } f \\ \text{from the first uncountable ordinal onto } \gamma$$

(this is proven by the usual methods, observing that such f clearly exists in $L[B]$ and in particular $f \in (H_{\omega_2})^{L[B]} = L_{\omega_2^{L[B]}}[B]$).

By elementarity, the statement is true also in $\overline{N}_\zeta = L_{\beta_\zeta}[B \cap \zeta]$. This clearly means $\omega_2^{L[B \cap \zeta]} \notin L_{\beta_\zeta}[B \cap \zeta]$, whereby $\beta_\zeta \leq \omega_2^{L[B \cap \zeta]}$. Since we just proved that $\omega_2^{L[B \cap \zeta]} \in M_\zeta$, we can finally conclude that $\beta_\zeta \in M_\zeta$. ■

□

4.2 A rigid Suslin tree

We present now Jensen's construction of a rigid normal Suslin tree by \diamond . In fact, the tree constructed below is the one originally defined by Jensen in his first proof of $\neg\text{SH}$ in L .

Theorem 4.11. *If \diamond holds, then there exists a rigid normal Suslin tree.*

Proof. The proof has much in common with the proof of Theorem 4.9. Let $A \subseteq \omega_1$ and M_α for $\alpha < \omega_1$ be as before. The rigid Suslin tree will be a set of countable sequences ordered by \subseteq . We define it by induction on levels. As before, $U_0 := \{\emptyset\}$ and $U_{\alpha+1} := \{s \smallfrown i \mid s \in U_\alpha \text{ and } i \in 2\}$, so it's left to deal with limit cases. Let $\alpha < \omega_1$ and suppose inductively that $T|_\alpha \in M_\alpha$ (this condition is clearly preserved in the successor steps). As usual, we will define the α th level U_α by extending α -branches in $T|_\alpha$. But differently from the methods we've seen so far, we will get *all* the necessary branches by a single forcing over M_α . We define (in M_α) the forcing poset (P, \leq_P) as follows:

$$P := \{p \mid p: a \rightarrow T|_\alpha, \text{ for some finite set } a \subseteq \omega\}$$

and the order is given by

$$p \leq_P q \iff \text{dom}(p) \supseteq \text{dom}(q) \text{ and } \forall n \in \text{dom}(q) [p(n) \supseteq q(n)].$$

M_α is countable, so in $L[A]$ there exists an M_α -generic filter for P . We take G as the least w.r.t. the canonical order of $L[A]$. Observe that actually $M_\alpha \in M_{\alpha+1}$ and $M_{\alpha+1} \preceq H_{\omega_1}$, so that M_α is countable in $M_{\alpha+1}$ and hence $G \in M_{\alpha+1}$. This will easily ensure that the induction hypothesis $T|_{(\alpha+1)} \in M_{\alpha+1}$ is preserved, once we complete the definition of U_α . For all $n \in \omega$, define

$$\mathfrak{b}_n := \{p(n) \mid p \in G\}.$$

Claim: The following hold:

- (i) every \mathfrak{b}_n is an α -branch of $T|_\alpha$;
- (ii) $n \neq m$ implies $\mathfrak{b}_n \neq \mathfrak{b}_m$;
- (iii) every \mathfrak{b}_n is M_α -generic for $T|_\alpha$;
- (iv) if $n \neq m$, then $\mathfrak{b}_n \times \mathfrak{b}_m$ is M_α -generic for $T|_\alpha \times T|_\alpha$ with the product ordering;
- (v) $T|_\alpha = \bigcup_{n \in \omega} \mathfrak{b}_n$.

Proof: Points (i) and (iii) clearly follow from (iv). Also (ii) does: take $n \neq m$ and suppose towards a contradiction that $\mathfrak{b}_n = \mathfrak{b}_m = \mathfrak{b}$. By (iv), $\mathfrak{b}_n \times \mathfrak{b}_m$ is M_α -generic for $T|_\alpha \times T|_\alpha$. By the Product Lemma, we get that \mathfrak{b} is $M[\mathfrak{b}]$ -generic for $T|_\alpha \times T|_\alpha$, and this is not possible⁴ because clearly $\mathfrak{b} \in M[\mathfrak{b}]$. Let's prove (iv). Suppose $n \neq m$ and $D \subseteq T|_\alpha \times T|_\alpha$ is dense. The set

$$D^* := \{p \in P \mid (p(n), p(m)) \in D\}$$

is clearly dense in P . So take $p \in G \cap D^*$. Then $(p(n), p(m)) \in \mathfrak{b}_n \times \mathfrak{b}_m \cap D$ and we are done.

As for (v), let $s \in T|_\alpha$ and define

$$D_s := \{p \in P \mid p(n) \supseteq s, \text{ for some } n \in \text{dom}(p)\}.$$

⁴It is a well-known and easy fact that if M is a countable transitive model and P is an atomless separative forcing poset, then no M -generic filter for P belongs to M .

D_s is dense in P , so there is $p \in G \cap D_s$. Thus $s \subseteq p(n) \in \mathfrak{b}_n$ for some $n \in \text{dom}(p)$, so $s \in \mathfrak{b}_n$ and the claim is proved. \blacksquare

So we define $U_{\alpha+1} := \{\bigcup \mathfrak{b}_n \mid n \in \omega\}$. By point (v) of the claim, $T|(\alpha+1)$ is normal. Finally, $T := \bigcup_{\alpha < \omega_1} U_\alpha$ is a normal ω_1 -tree. Using Claim 4.10, point (iii) of the claim ensures that T is Suslin. It remains to show that T is rigid. Suppose to the contrary that σ is a non-trivial automorphism of T . Using essentially the same argument of the proof of Claim 4.10, we will find some $\zeta < \omega_1$ such that $\sigma \restriction (T|_\zeta) \in M_\zeta$. By the construction of T , this will be easily seen to imply the existence of two branches \mathfrak{b}_n and \mathfrak{b}_m such that $\mathfrak{b}_n \in M_\zeta[\mathfrak{b}_m]$, which will in turn lead to a contradiction by the Product Lemma. We now give the full details.

Let $B \subseteq \omega_1$ be such that $A, T, \sigma \in L[B]$. Set $N^* := L_{\omega_2}^{L[B]}[B]$ and define an increasing sequence $\langle N_\nu \rangle_{\nu < \omega_1}$ of submodels of N^* precisely as in (4.3). For all $\nu < \omega_1$, set $\alpha_\nu := \omega_1 \cap N_\nu$ and mos_ν the collapsing function of N_ν , whose image is $\overline{N}_\nu = L_{\beta_\nu}[B \cap \alpha_\nu]$. As in the proof of Claim 4.10, there is $\zeta = \alpha_\zeta$ such that $\overline{N}_\zeta \subseteq M_\zeta$. Therefore

$$\text{mos}_\zeta(T) = T|_\zeta \in M_\alpha \quad \text{and} \quad \text{mos}_\zeta(\sigma) = \sigma \restriction (T|_\zeta) \in M_\alpha.$$

By elementarity, $\sigma \restriction (T|_\zeta)$ is non-trivial. So there is $s \in T|_\zeta$ such that $\sigma(s) \neq s$. Take $s' \in U_\zeta$ such that $s' \supset s$. Of course $\sigma(s') \neq s'$. Of course ζ is limit, hence by the construction of T at limit levels there are m and n distinct such that $s' = \bigcup \mathfrak{b}_m$ and $\sigma(s') = \bigcup \mathfrak{b}_n$. Thus $\sigma(s') = \bigcup_{\nu < \alpha} \sigma(s' \restriction \nu)$. Of course the latter belongs to any transitive models which contains M_ζ and s' , so that $\sigma(s') \in M_\zeta[s']$. But clearly $M_\zeta[s'] = M_\zeta[\mathfrak{b}_m]$, hence $\mathfrak{b}_n \in M_\zeta[\mathfrak{b}_m]$. However, $\mathfrak{b}_m \times \mathfrak{b}_n$ is M_ζ -generic for $T|_\zeta \times T|_\zeta$ by (iv) of the claim, so \mathfrak{b}_n is $M_\zeta[\mathfrak{b}_m]$ -generic for $T|_\alpha$ by the Product Lemma. Thus $\mathfrak{b}_n \notin M_\alpha[\mathfrak{b}_m]$, contradiction. \square

Bibliography

- [1] U. Abraham, *Construction of a rigid Aronszajn tree*, Proc. Amer. Math. Soc., 77(1):136–137, 1979.
- [2] K. Devlin and H. Johnsbråten, *The Souslin Problem*, Lecture Notes in Mathematics (405), Springer, 1974
- [3] K. Devlin, *Constructibility*, Springer-Verlag, 1984.
- [4] T. Jech, *Non-provability of Souslin's Hypothesis*, Commentationes mathematicae Universitatis Carolinae 8, 1967.
- [5] R. Jensen, *Souslin's Hypothesis is incompatible with $V = L$* , Notices of the American Mathematical Society 16, 1968.
- [6] T. Jech, *Trees*, The Journal of Symbolic Logic, Volume 36, Number 1, March 1971.
- [7] T. Jech, *Automorphisms of ω_1 -trees*, Trans. Amer. Math. Soc., 173:57–70, 1972.
- [8] Thomas J. Jech, *Set Theory*, The third millennium edition, Springer-Verlag, 2003.
- [9] A. Kanamori, *The Higher Infinite*, Springer, 2003.
- [10] A. Kanamori, *Historical remarks on Suslin's Problem*, Set Theory, Arithmetic and Foundations of Mathematics: Theorems, Philosophies, Lecture Notes in Logic, volume 36, 1-12. Association for Symbolic Logic, 2011.
- [11] K. Kunen, *Set Theory*, North-Holland Pub. Co., 1980.
- [12] K. Kunen and J. Vaughan, *Handbook of Set-theoretic Topology*, North-Holland, 1984.
- [13] K. Kunen, *The Foundations of Mathematics*, College Publications, 2009.
- [14] K. Kunen, *Set Theory*, College Publications, 2013.
- [15] G. Kurepa, *Ensembles ordonnés et ramifiés*, Publ. math. Univ. Belgrade, 1935.
- [16] J. Silver, *The independence of Kurepa's conjecture and two-cardinal conjectures in model theory*, Axiomatic Set Theory, Proc. Sympos. Pure Math., XIII, Providence, R.I.: Amer. Math. Soc., pp. 383–390, 1971.

- [17] R. Solovay and S. Tennenbaum, *Iterated Cohen extensions and Souslin's problem*, Annals of Mathematics (2) 94, 201–245, 1971.
- [18] E. Specker, *Sur un problème de Sikorski*, Colloquium Mathematicum, vol. 2 (1949), pp. 9–12.
- [19] D. Stewart, (title unknown), Master's thesis, University of Bristol, 1966.
- [20] S. Tennenbaum, *Souslin's Problem*, Proceedings of the National Academy of Sciences of the United States of America 59, 60–63, 1968.
- [21] S. Todorčević, *Rigid Aronszajn trees*, Publ. Inst. Math. (Beograd), 27(41):259–265, 1979.