PEG Powder Prep (2 Bottles)

Colonoscopy prep · Kurt Vernon, MD, PA

Buy at the Pharmacy

- ✓ PEG powder prep kit with 2 bottles and flavor packets. We send the prescription to your pharmacy. Tell the pharmacist that coupon savings information is attached in the "Note to Pharmacist."

Before Your Prep Day

1. 2 weeks before: if you take a weekly GLP-1 shot (such as semaglutide or dulaglutide) **only for weight loss**, stop it now.
2. 1 week before: if you take a weekly GLP-1 shot **for diabetes**, stop it now. If you take a daily diabetes medicine of this kind (such as liraglutide, exenatide, oral semaglutide or dapagliflozin), take half your dose the day before your procedure.
3. 7 days before: stop phentermine.
4. 5 days before: stop aspirin 325 mg. Keep taking aspirin 81 mg.
5. If you take a blood thinner such as warfarin, enoxaparin, heparin, rivaroxaban or clopidogrel, call the office for instructions.
6. For pain or a headache before your procedure, acetaminophen is fine.

The Day Before

1. Clear liquids only, all day. No solid food. Drink plenty of clear liquids so you stay hydrated.
2. If you have diabetes, take half your usual dose of diabetes medicine today and check your blood sugar during the day. If you take insulin, bring it to your appointment.
3. **4:00 pm, first dose:** open one flavor-enhancing packet and pour it into one bottle. Fill the bottle with lukewarm water to the fill line, cap it and shake gently until all the powder dissolves. (For the best taste, refrigerate it for an hour first.) Drink it, then drink another 16 ounces of water. Keep drinking clear liquids.
4. **10:00 pm, second dose:** do the same with the second packet and bottle, then drink another 16 ounces of water. Keep drinking clear liquids until midnight.
5. Nothing by mouth after midnight, including gum, hard candy or lozenges. No smoking, chewing tobacco, e-cigarettes or vaping.

The Day of Your Colonoscopy

1. Nothing to eat or drink. Eating or drinking anything may get your procedure canceled.
2. The only medicines you may take this morning are pain, anti-anxiety, seizure, heart or blood pressure medicines, with a sip of water. Hold all other medicines until after your procedure, unless the office tells you otherwise.
3. If you use a prescription inhaler, bring it with you.
4. Someone must come with you and drive you home, because you will be sedated. Plan for someone to stay with you the rest of the day, since you may be drowsy or dizzy.
5. Leave jewelry and valuables at home. We can't be responsible for them.
6. Your colonoscopy is at our Fuquay-Varina Endoscopy Center.

Clear Liquids You Can Have

No milk or milk products (including non-dairy creamer), and nothing red, blue or purple.

- Water
- Soft drinks, including colas
- Coffee or tea, without milk, cream or creamer
- Sports drinks (green or yellow)
- Gelatin (green or yellow)
- Ice pops (green or yellow)
- Hard candy
- Chicken or beef broth or bouillon (no vegetables)
- Apple juice, white grape juice or other strained fruit juice

---

You must finish the whole prep. A completely clean colon lets Dr. Vernon see everything, including polyps.

Can't make it? Call at least 48 hours ahead to reschedule. There is a $75 fee for a no-show, or for canceling with less than 48 hours' notice.

Questions about your prep? Call Fuquay-Varina at (919) 577-0085 or Dunn at (910) 891-5808, Monday – Thursday, 8:30 am – 5:00 pm.

