

Website: thermofisher.com Customer Service (US): 1 800 955 6288 ext. 1 Technical Support (US): 1 800 955 6288 ext. 441 thermofisher.com/contactus



Product data sheet

Tau Monoclonal Antibody (T14)

Catalog Number 13-1400

Details	
Size	100 μg
Host/Isotope	Mouse / IgG1, kappa
Class	Monoclonal
Туре	Antibody
Clone	T14
Immunogen	Bovine Tau.
Conjugate	Unconjugated
Form	Liquid
Purification	purified
Storage buffer	PBS
Contains	0.1% sodium azide
Storage Conditions	-20°C

Species Reactivity	
Species reactivity	Human
Published species	Mouse, Human, Not Applicable
Tested Applications	Dilution *
ELISA (ELISA)	0.1-0.5 μg/mL
Immunofluorescence (IF)	5-10 μg/mL
Immunohistochemistry (IHC)	5-10 μg/mL
Immunoprecipitation (IP)	2-5 μg
Western Blot (WB)	1:250
Published Applications	
Immunofluorescence (IF)	See 1 publications below
Western Blot (WB)	See 5 publications below
Immunohistochemistry (IHC)	See 3 publications below

^{*} Suggested working dilutions are given as a guide only. It is recommended that the user titrate the product for use in their own experiment using appropriate negative and positive controls.

Background/Target Information

Tau is a neuronal microtubule-associated protein found predominantly on axons. The function of Tau is to promote tubulin polymerization and stabilize microtubules. The C-terminus binds axonal microtubules while the N- terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by TAU/MAPT localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton while the longer isoforms may preferentially play a role in its stabilization. In its hyper-phosphorylated form, Tau is the major component of paired helical filaments (PHF), the building block of neurofibrillary lesions in Alzheimer's diseases (AD) brain. Hyper-phosphorylation impairs the microtubule binding function of Tau, resulting in the destabilization of microtubules in AD brains, ultimately leading to the degeneration of the affected neurons. Numerous serine/threonine kinases phosphorylate Tau, including GSK-3beta, protein kinase A (PKA), cyclindependent kinase 5 (cdk5) and casein kinase II. Hyper-phosphorylated Tau is found in neurofibrillary lesions in a range and other central nervous system disorders such as Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy.

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization.


For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization.

Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any product will conform to such model or sample furnished to Buyer is merely illustrative of the general type and quality of goods and does not represent that any product will conform to such model or sample furnished to Buyer is merely illustrative of the product by each of the product by an advantage of the product and the product of the product by a supplied to the

NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD IS LIMITED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCTS) AT SELLER'S SOLE OPTION. THERE IS NO BUILGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MALEURE, (II) MISUSE, FAULT OR MERCHANT OF FOR BY BUYER, (III) USE OF THE PRODUCTS IN A MANNEE POR WHICH THEY WERE NOT DESIGNED, OR (IV) IMPROPER STORAGE AND HANDLING OF THE PRODUCTS. Unless otherwise expressely stated on the Product or in the documentation accompanying the Product, the Product of in the object of the Product of Indian Companying the Product of Indian C




Product Images For Tau Monoclonal Antibody (T14)



Tau Antibody (13-1400) in IHC

Immunohistochemical staining of Alzheimer's disease brain tissue using Ms anti-Tau (Product # 13-1400).



Tau Antibody (13-1400) in WB

Western blot analysis of Tau was performed by loading 20 µg of SHSY-5Y (lane1), Hep G2 (lane2), COLO205 (lane3), MCF7 (lane4) and Jurkat (lane5) cell lysates using Novex® NuPAGE® 4-12 % Bis-Tris gel (Product # NP0322BOX), XCell SureLock™ Electrophoresis System (Product # El0002), Novex® Sharp Pre-Stained Protein Standard (LC5800), and iBlot® 2 Dry Blotting System (lB21001). Proteins were transferred to a nitrocellulose membrane and blocked with 5 % skim milk at 4°C overnight. Tau was detected at ~ 78 kDa using Tau Mouse Monoclonal Antibody (Product # 13-1400) at 2-3 µg /mL in 5 % skim milk for 3 hours at room temperature on a rocking platform. Goat Anti-Mouse - HRP Secondary Antibody (Product # 62-6520) at 1:4000 dilution was used and chemiluminescent detection was performed using Novex® ECL Chemiluminescent Substrate Reagent Kit (WP20005).

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization.

Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package inserts ("Documentation"). No claim of suitability for use in applications regulated by PDA is made. The warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample furnished to Buyer merely illustrative of the general type and quality of goods and does not represent that any Poucit will conform to such model or some

IO OTHER WARRANTIES, EVPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD S
SUMMED TO REPAIR, REPLACEMENT OF OR REFUND FOR THE NON-CONFORMING PRODUCTS) AT SILE AND EXPLORED. FOR THE NON-CONFORMING PRODUCTS IS AT SILE AND EXPLORED. FOR THE NON-CONFORMING PRODUCTS IS AT SILE AND EXPLORED. FOR THE NON-CONFORMING PRODUCTS AT SILE AND EXPLORED. FOR THE N



1 Immunoflugrassanas Daf	
1 Immunofluorescence Ref	
Species / Dilution	Summary
Human / Not Cited	Annals of neurology (Feb 1998; 43: 259) "Cognitive, neuroimaging, and pathological studies in a patient with Pick's disease." Author(s):Lieberman AP,Trojanowski JQ,Lee VM,Balin BJ,Ding XS,Greenberg J,Morrison D,Reivich M,Grossman M PubMed Article URL:http://dx.doi.org/10.1002/ana.410430218
5 Western Blot References	
Species / Dilution	Summary
Mouse / Not Cited	The Journal of biological chemistry (Jul 1999; 274: 21464) "Characterization of mouse dishevelled (DvI) proteins in Wnt/Wingless signaling pathway." Author(s):Lee JS,Ishimoto A,Yanagawa S PubMed Article URL:http://dx.doi.org/null
Human / Not Cited	Annals of neurology (Feb 1998; 43: 259) "Cognitive, neuroimaging, and pathological studies in a patient with Pick's disease." Author(s):Lieberman AP,Trojanowski JQ,Lee VM,Balin BJ,Ding XS,Greenberg J,Morrison D,Reivich M,Grossman M PubMed Article URL:http://dx.doi.org/10.1002/ana.410430218
Human / Not Cited	Journal of neuropathology and experimental neurology (Jun 1998; 57: 588) "The neuropathology of a chromosome 17-linked autosomal dominant parkinsonism and dementia ("pallido-ponto nigral degeneration")." Author(s):Reed LA,Schmidt ML,Wszolek ZK,Balin BJ,Soontornniyomkij V,Lee VM,Trojanowski JQ,Schelper RL PubMed Article URL:http://dx.doi.org/null
	13-1400 was used in western blot to suggest Wnt-activation occurs prior to 3 months of age in the JNPL3 mouse model of frontotemporal dementia
Not Applicable / Not Cited	Neurobiology of aging (Jan 2009; 30: 14) "Wnt-pathway activation during the early stage of neurodegeneration in FTDP-17 mice." Author(s):Wiedau-Pazos M,Wong E,Solomon E,Alarcon M,Geschwind DH PubMed Article URL:http://dx.doi.org/10.1016/j.neurobiolaging.2007.05.015
Mouse / Not Cited	The Journal of neuroscience: the official journal of the Society for Neuroscience (Nov 2005; 25: 10637) "Age-dependent neurofibrillary tangle formation, neuron loss, and memory impairment in a mouse model of human tauopathy (P301L)."
Human / Not Cited	Author(s):Ramsden M,Kotilinek L,Forster C,Paulson J,McGowan E,SantaCruz K,Guimaraes A,Yue M,Lewis J,Carlson G, Hutton M,Ashe KH PubMed Article URL:http://dx.doi.org/10.1523/JNEUROSCI.3279-05.2005
3 Immunohistochemistry R	eferences
Species / Dilution	Summary
Mouse / Not Cited	Science (New York, N.Y.) (Aug 2001; 293: 1491) "Formation of neurofibrillary tangles in P301I tau transgenic mice induced by Abeta 42 fibrils." Author(s):Götz J,Chen F,van Dorpe J,Nitsch RM PubMed Article URL:http://dx.doi.org/10.1126/science.1062097
Human / Not Cited	Annals of neurology (Feb 1998; 43: 259) "Cognitive, neuroimaging, and pathological studies in a patient with Pick's disease." Author(s):Lieberman AP,Trojanowski JQ,Lee VM,Balin BJ,Ding XS,Greenberg J,Morrison D,Reivich M,Grossman M PubMed Article URL:http://dx.doi.org/10.1002/ana.410430218
Mouse / Not Cited Human / Not Cited	The Journal of neuroscience: the official journal of the Society for Neuroscience (Nov 2005; 25: 10637) "Age-dependent neurofibrillary tangle formation, neuron loss, and memory impairment in a mouse model of human tauopathy (P301L)." Author(s):Ramsden M,Kotilinek L,Forster C,Paulson J,McGowan E,SantaCruz K,Guimaraes A,Yue M,Lewis J,Carlson G, Hutton M,Ashe KH

For Research Use Only. Not for use in diagnostic procedures. Not for resale without express authorization.

Products are warranted to operate or perform substantially in conformance with published Product specifications in effect at the time of sale, as set forth in the Production documentation, specifications and/or accompanying package insents ("Documentation"). No claim of suitability for use in applications regulated by FDA is made. The warranty provided herein is valid only when used by properly trained individuals. Unless otherwise stated in the Documentation, this warranty is limited to one year from date of shipment when the Product is subjected to normal, proper and intended usage. This warranty does not extend to anyone other than the Buyer. Any model or sample. It is metry illustrative of the general type and quality of goods and does not represent that any Product will confront to sample.

NO OTHER WARRANTIES, EXPRESS OR IMPLED, ARE GRANTED INCLUDING WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR NON INFRINGEMENT. BUYER'S EXCLUSIVE REMEDY FOR NON-CONFORMING PRODUCTS DURING THE WARRANTY PERIOD ISLIMITED TO REPAIR, REPLACE MENT OF OR REFUND FOR THE NON-CONFORMING PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGATION TO REPAIR, REPLACE OR REFUND FOR PRODUCTS AS THE RESULT OF (I) ACCIDENT, DISASTER OR EVENT OF FORCE MAJEURE, (II) MISUSE, FAULT OR NEGLIGATION OF OR Y BUYER, (II) USE OF THE PRODUCTS IN AN ANNAIRE FOR WHITCH THEY WERE NOT DESIGNED, ON IMPROPER'S STORAGE AND HANDING OF THE PRODUCTS. Unless otherwise expressely stated on the Product or in the documentation accompanying the Product, the Product is intended for research only and is not to be used for any other purpose, including without limitation, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses, or any type of consumption by or application to human or animals.

