

CHAPTER 19

Spontaneous fMRI activity during resting wakefulness and sleep

Jeff Duyn*

Section for Advanced MRI, LFMI, NINDS, National Institutes of Health, Bethesda, MD, USA

Abstract: Functional magnetic resonance imaging (fMRI) studies performed during both waking rest and sleep show that the brain is continually active in distinct patterns that appear to reflect its underlying functional connectivity. In this review, potential sources that contribute to spontaneous fMRI activity will be discussed.

Keywords: sleep; brain function; fmri; spontaneous activity; connectivity; fluctuations.

Introduction

Much of the success of blood oxygen level dependent (BOLD) functional magnetic resonance imaging (fMRI) over the past two decades in studying the brain's functional architecture can be attributed to its ability to map activity changes in response to carefully crafted behavioral tasks. Nevertheless, it is becoming increasingly clear that fMRI may have important applications to the study of spontaneous brain activity as well (Biswal et al., 1995; Raichle, 2009). Similar to electrical and optical signals recorded from the brain (Arieli et al., 1995; Kenet et al., 2003; Shoham and Grinvald, 2001; Tsodyks et al.,

1999), fMRI signals show a rich spatiotemporal structure even during periods of apparent mental or behavioral inactivity, including sleep and anesthetized conditions. This structure appears, at least in part, to reflect the brain's underlying functional architecture (Biswal et al., 1995; Damoiseaux et al., 2006; Hampson et al., 2002; Salvador et al., 2005). fMRI of spontaneous activity may therefore aid in the mapping of the brain's major communication pathways, despite its temporal resolution being far too poor to cover the breadth of the spectrum of neural communications. Nevertheless, the origin and role of spontaneous activity are still poorly understood. In the following I will discuss some of the characteristics of spontaneous fMRI signal fluctuations and speculate on their origin.

*Corresponding author.

Tel.: +1-301-594-7305; Fax: +1-301-480-1981

E-mail: jhd@helix.nih.gov

What is spontaneous fMRI activity?

The BOLD fMRI signal derives from repetitive magnetic resonance imaging (MRI) scans that are sensitized to changes in blood oxygenation (Ogawa et al., 1990). In most brain regions, changes in neuronal activity evoked by sensory or behavioral tasks lead to disproportionately large blood flow increases, thought to overcompensate for the metabolic demands supporting the increased glutamate cycling at synapses of pyramidal cells (Logothetis, 2002). The resulting increases in blood oxygenation lead to small (about 1%) fMRI signal increases, the dynamics of which can be sampled at a seconds-resolution time scale.

Interestingly, in absence of behavioral and sensory tasks or overt activity, similarly sized signal fMRI fluctuations can be observed (Biswal et al., 1995); these fluctuations are correlated between regions that have an apparent functional relationship (Biswal et al., 1995; Krienen and Buckner, 2009) and generally occur in multiple, spatially independent, and reproducible patterns that resemble many of the patterns observed during task activation studies (Smith et al., 2009). Thus, the study of spontaneous fMRI activity may provide information about the brain's architecture and function.

Despite the intriguing characteristics of spontaneous fMRI signal fluctuations, and their rapidly increasing application to neuroscientific questions, their origin and role are still poorly understood. Importantly, BOLD fMRI signal fluctuations may contain both neuronal and non-neuronal contributions, and it is important separate these. For example, head motion and MRI system instabilities may lead to fMRI signal fluctuations (see below) that may confound any neuronal sources. In this regard, it is important to differentiate between spontaneous fMRI *fluctuations* and spontaneous fMRI *activity*, the latter presumably representing neurometabolic contributions only. Another often overlooked fact is that spontaneous and task-evoked fMRI activity are not mutually exclusive (Bianciardi et al.,

2009; de Zwart et al., 2008; Fox et al., 2007). An indication of this oversight is the use of the term “resting state activity,” which incorrectly suggests that spontaneous activity ceases in the presence of overt tasks. As of yet, the interaction of these two types of activity is not well understood. Summarized, it appears safe to say that fruitful application to neuroscience will require further understanding of origin and role of spontaneous and sophisticated techniques to remove potential neuronal from non-neuronal sources.

Nonneuronal contributions to the BOLD fMRI signal

As indicated in the previous section, the BOLD fMRI signal is measured as a small deviation of the baseline or mean signal. For this reason, it is relatively sensitive to subtle fluctuations in the baseline signal due to nonneuronal sources such as thermal noise, instrumental drift, head motion, and physiological variations (Bianciardi et al., 2009). These need to be dealt with to properly interpret the effects of spontaneous neural activity.

Thermal noise originates from resistive electrical effects in brain tissue and the MRI detector. It can be minimized by appropriate choice of MRI acquisition parameters (e.g., spatial resolution) and the use of advanced technologies such as high magnetic field and array detectors (de Zwart et al., 2004; Triantafyllou et al., 2005). Unfortunately, the amplitudes of all other noise sources roughly scale with signal amplitude, and some even have BOLD like contrast behavior (Hyde et al., 2001) making it difficult to reduce their contribution relative to the (neuronal) signal of interest.

A rather effective way to separate noise sources is the use of postprocessing and linear model regressors. For example, head motion can be compensated for by image alignment across the time-series scans; subsequently the alignment parameters can be used as variance regressors. Similarly, regressors can be constructed for the effects of cardiac and respiratory cycles (and their

rates) on the fMRI signal, all of which can be substantial (Birn et al., 2006; Chang and Glover, 2009; Fukunaga et al., 2006; Glover et al., 2000; Shmueli et al., 2007). Another important source of fMRI signal variability is low-frequency drift, which is a slow, widespread temporal variation in the baseline signal and may originate from subject motion, (neuro-) physiological sources, and instrumental instabilities (Smith et al., 1999). Although low frequency drifts are generally removed using high-pass Fourier filtering, time-domain polynomial fitting or wavelet analysis may be preferable as these avoid time-series truncation effects on the Fourier spectrum. A difficulty common to all methods that tackle low frequency drift is the potential removal of slowly varying neuronal effects. This is especially true for conditions where there is a potential comodulation of neuronal and physiological signals (Golanov et al., 1994; Musizza et al., 2007).

Alternative other methods for the removal of nonneuronal contributions include independent component analysis (based on spatial distribution of correlation patterns; Beckmann and Smith, 2004) and the use of regressors derived from reference regions assumed to not to reflect neuronal activity (e.g., Cerebro-Spinal Fluid (CSF) or white matter). The effectiveness of the latter may be compromised by venous drainage from neuronally active regions, potentially injecting a neuronal contribution into the reference signal (Bianciardi et al., 2011). In summary, as the separation of neuronal and nonneuronal sources has only been partly resolved, caution needs to be exercised with the interpretation of spontaneous fMRI studies.

Neuronal correlates of the BOLD fMRI signal

Although the origin of spontaneous fMRI signal fluctuations is still poorly understood, there is converging evidence that these fluctuations, at least in part, represent underlying neuronal activity. One category of evidence is the similarity between the fMRI findings and electrophysiological recordings.

For example, spontaneous fluctuations in synaptic membrane potentials, measured with voltage-sensitive dyes in animal models, have been shown to contain a spatially localized and functionally specific component (Arieli et al., 1995; Shoham and Grinvald, 2001). In humans, electrocortical recordings have confirmed this functionally specific correlation (He et al., 2008; Nir et al., 2006), and further have shown low-frequency variations with spatiotemporal properties similar to the fMRI data (He et al., 2008). Lastly, noninvasive macroscopic electrical recordings with magnetoencephalography (MEG) have shown long range correlation in specific networks similar to that seen in fMRI (de Pasquale et al., 2010; Liu et al., 2010).

Another piece of evidence for a neuronal contribution to the spontaneous fMRI signal fluctuations comes from an fMRI study recording BOLD and perfusion signals simultaneously (Fukunaga et al., 2008), both of which have been shown to fluctuate spontaneously (Biswal et al., 1997). During task-evoked activity, it has been demonstrated that their relative fluctuation level depends on the metabolic contribution to the underlying process (Davis et al., 1998; Hoge et al., 1999). This metabolic contribution is thought to be primarily associated with the adenosinetriphosphate (ATP) turnover underlying glutamate cycling at neuronal synapses (Attwell and Iadecola, 2002). Applying this approach to the study of spontaneous activity, Fukunaga et al. (2008) found that the metabolic contribution to spontaneous activity is similar to that during task evoked activity. The tentative interpretation is that similar neuronal events are responsible for both task evoked and spontaneous activity.

Finally, multimodal recordings in humans such as positron emission tomography (PET)-EEG (electroencephalography) and fMRI-EEG are also suggestive of a neuroelectrical contribution to spontaneous fMRI signal fluctuations. For example, during awake rest, substantial negative correlation with EEG band-limited power in the alpha range has been observed over extended

brain regions (Goldman et al., 2002; Laufs et al., 2003a, 2006; Moosmann et al., 2003; Sadato et al., 1998), and correlation with other EEG bands and features have been found as well (Britz et al., 2010; de Munck et al., 2009; Laufs et al., 2003b; Mantini et al., 2007; Musso et al., 2010). During sleep and anesthesia, significant correlation with EEG delta and subdelta, and with spindle activity has been found (Dang-Vu et al., 2005, 2008; Schabus et al., 2007). Finally, studies recording intracortical electrical signals concurrently with fMRI in macaque have found substantial intermodal correlation in various frequency ranges, including the gamma band (Scholvinck et al., 2010; Shmuel and Leopold, 2008).

Spontaneous fMRI activity during sleep

To the extent that spontaneous fMRI fluctuations represent neuroelectrical processes, how do they relate to brain function? Do they simply represent ongoing and uncontrolled conscious mentation? This is probably at least partly true for many of the fMRI studies on spontaneous activity presented in the literature, as they often used poorly controlled behavioral conditions that may have included periods of “random episodic silent thinking” or “REST” (Kanwisher, 2001), mental imagery (Kosslyn et al., 1999; Wheeler et al., 2000), monitoring of body posture and the environment, and fluctuations in vigilance, arousal, and attention. Although initial reports of a dependence on behavioral state appeared to confirm a primary contribution of uncontrolled mental activity (Fox et al., 2005; Fransson, 2005; McKiernan et al., 2003), this has been recently put into doubt by studies showing at least maintained (Boly et al., 2008, 2009; Fukunaga et al., 2006, 2008; Horovitz et al., 2008; Kiviniemi et al., 2000; Larson-Prior et al., 2009; Vincent et al., 2007) or only partly reduced (Horovitz et al., 2009; Lu et al., 2007; Martuzzi et al., 2010; Peltier et al., 2005) levels of spontaneous fMRI activity during conditions of reduced consciousness. Notably, in

sleep, which is a natural condition of reduced consciousness, spontaneous activity is largely maintained (Fukunaga et al., 2006; Horovitz et al., 2008, 2009; Larson-Prior et al., 2009). The figure illustrates this point: the widespread involvement of the brain in spontaneous activity observed during the waking condition is also observed during states of light and deep sleep (Fig. 1).

In contrast, baseline metabolic activity may be reduced during conditions of reduced consciousness in most brain regions, and most notably in the thalamus and frontal cortex (Alkire et al., 1999; Braun et al., 1997; Buchsbaum et al., 2001; Maquet et al., 1990). However, this does not necessarily contradict the fMRI findings, which report on *fluctuations in* rather than *mean levels of* baseline metabolic levels underlying synaptic activity. The continuation of spontaneous fMRI activity during sleep and anesthesia suggests that this activity may not simply represent mentation and sensory processing, but at least in part subserves more basic brain functions, possibly including (synaptic) homeostasis and memory consolidation (Fukunaga et al., 2005, 2006). Conversely, the study of spontaneous activity could elucidate the role of sleep for brain function, which remains a fascinating and poorly understood issue.

Origin and role of spontaneous fMRI activity

Despite the increasing evidence of a neuronal contribution to spontaneous fMRI signals, the suggestions of their behavioral relevance (Sadaghiani et al., 2010), and the rapidly increasing application of spontaneous fMRI to the study of brain functional connectivity in health and disease, the origin of role of this neuronal activity is still poorly understood. One question that is relevant to the functional connectivity is whether spontaneous activity represents corticocortical communication or rather the effect of a common driving source. For example, the apparent corticocortical correlations may result from




Fig. 1. *Intersubject consistency of ICA-derived spatial components of spontaneous activity across sleep stages.* Left half of figure: different colors indicate the individual consistent components (colors were randomly assigned). Right half of figure: color scale indicates probability of spatial overlap of components across subjects. Across both wake and sleep, much of the brain involved in spontaneous activity. From: Masaki Fukunaga, Silvina Horovitz, Walter Carr, Dante Picchioni, Jacco de Zwart, Tom Balkin, Allen Braun, Jeff Duyn, presented at the Organization for Human Brain Mapping 2007 conference.

changes in excitability or homeostatic processes orchestrated from brainstem and midbrain regions; such processes could, for example, be effectuated by cholinergic brain stem neurons (Mena-Segovia et al., 2008a,b), neurons in the rostral ventrolateral medulla (Golanov et al., 1994), or neurons in the reticular complex of the thalamus, amongst others. Arguing against this

“common source” hypothesis are recent studies of spontaneous activity with voltage sensitive dyes and fMRI, which found much reduced interhemispheric corticocortical correlation of spontaneous activity when integrity of the corpus callosum, which form the main body of interhemispheric fibers, was compromised (Johnston et al., 2008; Mohajerani et al., 2010; Quigley et al., 2003).

A caveat however with these studies is that the integrity of the “common source” may rely on an intact corpus callosum, as may be the case for the thalamus (Raos and Bentivoglio, 1993).

One potential role for spontaneous brain activity may be a homeostatic one in support of synaptic downscaling and/or consolidation (Fukunaga et al., 2005, 2006; Miall and Robertson, 2006). A number of electrophysiological studies has reported on hippocampal–cortical communications preferentially active during “off-line” waking periods and sleep, that may support a hebbian adjustment of synaptic strengths for memory consolidation (Diba and Buzsaki, 2007; Foster and Wilson, 2006; Hahn et al., 2006; Isomura et al., 2006; Ji and Wilson, 2007; Sirota et al., 2003). Notably, during sleep, this hippocampal–cortical dialogue is primarily played out during so-called cortical “Up” states, during which the firing rates of large neuronal groups synchronously increase; additionally, the repetitive cycling between “Up” and “Down” states, observed in electrical recordings as a neocortical slow oscillation (Steriade et al., 1993), has been suggested to facilitate sleep dependent synaptic plasticity (Brazier, 1949; Luczak et al., 2007; Massimini et al., 2004; Petersen et al., 2003; Tononi and Cirelli, 2003). Furthermore, learning has been demonstrated to affect the pattern of slow oscillations during subsequent sleep (Huber et al., 2004; Molle et al., 2004, 2009). Importantly, preliminary evidence from studies of deep sleep in humans and anesthesia in animals (the latter also show a repetitive cycling of “Up” and “Down” states) suggest that slow oscillations may have a substantial fMRI correlate (Dang-Vu et al., 2008; Liu et al., 2011), although their scalp distribution may not possess the spatial specificity exhibited by the fMRI activity (Dang-Vu et al., 2008; Murphy et al., 2009).

On the other hand, both the slow oscillation and the alternation between “Up” and “Down” states are virtually absent during the waking state, during which a substantial level of spontaneous fMRI persists. These hypothesized sources

therefore do not fully explain the characteristics of spontaneous fMRI activity, and it is possible that some of the activity relates to “off-line” learning during the waking state without the need for facilitatory slow oscillations. In fact, there have been a number of recent reports showing increased spontaneous activity in specific functional networks during off-line periods following learning tasks, including motor learning (Albert et al., 2009), visual processing (Stevens et al., 2010), and visual perceptual learning (Lewis et al., 2009).

Use of spontaneous fMRI to study brain connectivity

Concurrently with the attempts to establish the origin and role of spontaneous activity, an increasing number of researchers are starting to analyze spontaneous fMRI activity to improve the understanding of brain connectivity. Recent work has shown that the major correlation patterns in spontaneous activity show a striking resemblance with predictions derived from structural connectivity modeling based on diffusion tensor imaging (Honey et al., 2009). The major network patterns of spontaneous activity may therefore reflect communication between cortical hubs that subserve, in a highly efficient manner, the elaborate connectivity between the local processing modules of the human brain (Buckner et al., 2009). Spontaneous fMRI activity may therefore provide complementary information to diffusion tensor imaging (DTI) in studying brain connectivity, and provide functional information that is difficult to obtain from electrophysiological studies, which are either too spatially constrained (in the case of intracortical electrodes) or have poor spatial specificity (in the case of EEG and MEG).

One potentially important clinical application that is developing rapidly is the study of mental and neurological disorders (Fox and Greicius, 2010). Striking examples are altered network activity (and possibly connectivity) in diseases of

consciousness (Boly et al., 2008, 2009) and in dementia, including Alzheimer's disease (Zhang et al., 2010; Zhou et al., 2010). What is currently not well understood is what the causal relationship is between the fMRI findings and the disease process, their neurobiological origin, and their specificity to the disease. Future mechanistic studies on animal models may shed a light on these issues.

Abbreviations

ATP	adenosinetriphosphate
DTI	diffusion tensor imaging
EEG	electroencephalography
fMRI	functional magnetic resonance imaging
MEG	magnetoencephalography
MRI	magnetic resonance imaging
PET	positron emission tomography

References

- Albert, N. B., Robertson, E. M., & Miall, R. C. (2009). The resting human brain and motor learning. *Current Biology*, 19, 1023–1027.
- Alkire, M. T., Pomfrett, C. J., Haier, R. J., Gianzero, M. V., Chan, C. M., Jacobsen, B. P., et al. (1999). Functional brain imaging during anesthesia in humans: Effects of halothane on global and regional cerebral glucose metabolism. *Anesthesiology*, 90, 701–709.
- Arieli, A., Shoham, D., Hildesheim, R., & Grinvald, A. (1995). Coherent spatiotemporal patterns of ongoing activity revealed by real-time optical imaging coupled with single-unit recording in the cat visual cortex. *Journal of Neurophysiology*, 73, 2072–2093.
- Attwell, D., & Iadecola, C. (2002). The neural basis of functional brain imaging signals. *Trends in Neurosciences*, 25, 621–625.
- Beckmann, C. F., & Smith, S. M. (2004). Probabilistic independent component analysis for functional magnetic resonance imaging. *IEEE Transactions on Medical Imaging*, 23, 137–152.
- Bianciardi, M., Fukunaga, M., van Gelderen, P., de Zwart, J. A., & Duyn, J. H. (2011). Negative BOLD-fMRI signals in large cerebral veins. *Journal of Cerebral Blood Flow & Metabolism*, 31, 401–412.
- Bianciardi, M., Fukunaga, M., van Gelderen, P., Horovitz, S. G., de Zwart, J. A., & Duyn, J. H. (2009). Modulation of spontaneous fMRI activity in human visual cortex by behavioral state. *NeuroImage*, 45, 160–168.
- Bianciardi, M., Fukunaga, M., van Gelderen, P., Horovitz, S. G., de Zwart, J. A., Shmueli, K., et al. (2009). Sources of functional magnetic resonance imaging signal fluctuations in the human brain at rest: A 7T study. *Magnetic Resonance Imaging*, 27, 1019–1029.
- Birn, R. M., Diamond, J. B., Smith, M. A., & Bandettini, P. A. (2006). Separating respiratory-variation-related fluctuations from neuronal-activity-related fluctuations in fMRI. *NeuroImage*, 31, 1536–1548.
- Biswal, B. B., Van Kylen, J., & Hyde, J. S. (1997). Simultaneous assessment of flow and BOLD signals in resting-state functional connectivity maps. *NMR in Biomedicine*, 10, 165–170.
- Biswal, B., Yetkin, F. Z., Haughton, V. M., & Hyde, J. S. (1995). Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magnetic Resonance in Medicine*, 34, 537–541.
- Boly, M., Phillips, C., Tshibanda, L., Vanhaudenhuyse, A., Schabus, M., Dang-Vu, T. T., et al. (2008). Intrinsic brain activity in altered states of consciousness: How conscious is the default mode of brain function? *Annals of the New York Academy of Sciences*, 1129, 119–129.
- Boly, M., Tshibanda, L., Vanhaudenhuyse, A., Noirhomme, Q., Schnakers, C., Ledoux, D., et al. (2009). Functional connectivity in the default network during resting state is preserved in a vegetative but not in a brain dead patient. *Human Brain Mapping*, 30, 2393–2400.
- Braun, A. R., Balkin, T. J., Wesenten, N. J., Carson, R. E., Varga, M., Baldwin, P., et al. (1997). Regional cerebral blood flow throughout the sleep-wake cycle. An H2(15)O PET study. *Brain*, 120(Pt 7), 1173–1197.
- Brazier, M. A. (1949). The electrical fields at the surface of the head during sleep. *Electroencephalography and Clinical Neurophysiology*, 1, 195–204.
- Britz, J., Van De Ville, D., & Michel, C. M. (2010). BOLD correlates of EEG topography reveal rapid resting-state network dynamics. *NeuroImage*, 52, 1162–1170.
- Buchsbaum, M. S., Hazlett, E. A., Wu, J., & Bunney, W. E. Jr. (2001). Positron emission tomography with deoxyglucose-F18 imaging of sleep. *Neuropsychopharmacology*, 25, S50–S56.
- Buckner, R. L., Sepulcre, J., Talukdar, T., Krienen, F. M., Liu, H., Hedden, T., et al. (2009). Cortical hubs revealed by intrinsic functional connectivity: Mapping, assessment of stability, and relation to Alzheimer's disease. *The Journal of Neuroscience*, 29, 1860–1873.
- Chang, C., & Glover, G. H. (2009). Relationship between respiration, end-tidal CO₂, and BOLD signals in resting-state fMRI. *NeuroImage*, 47, 1381–1393.
- Damoiseaux, J. S., Rombouts, S. A., Barkhof, F., Scheltens, P., Stam, C. J., Smith, S. M., et al. (2006). Consistent resting-state networks across healthy subjects. *Proceedings of the*

- National Academy of Sciences of the United States of America*, 103, 13848–13853.
- Dang-Vu, T. T., Desseilles, M., Laureys, S., Degueldre, C., Perrin, F., Phillips, C., et al. (2005). Cerebral correlates of delta waves during non-REM sleep revisited. *NeuroImage*, 28, 14–21.
- Dang-Vu, T. T., Schabus, M., Desseilles, M., Albouy, G., Boly, M., Darsaud, A., et al. (2008). Spontaneous neural activity during human slow wave sleep. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 15160–15165.
- Davis, T. L., Kwong, K. K., Weisskoff, R. M., & Rosen, B. R. (1998). Calibrated functional MRI: Mapping the dynamics of oxidative metabolism. *Proceedings of the National Academy of Sciences of the United States of America*, 95, 1834–1839.
- de Munck, J. C., Goncalves, S. I., Mammoliti, R., Heethaar, R. M., & Lopes da Silva, F. H. (2009). Interactions between different EEG frequency bands and their effect on alpha-fMRI correlations. *NeuroImage*, 47, 69–76.
- de Pasquale, F., Della Penna, S., Snyder, A. Z., Lewis, C., Mantini, D., Marzetti, L., et al. (2010). Temporal dynamics of spontaneous MEG activity in brain networks. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 6040–6045.
- de Zwart, J. A., Gelderen, P., Fukunaga, M., & Duyn, J. H. (2008). Reducing correlated noise in fMRI data. *Magnetic Resonance in Medicine*, 59, 939–945.
- de Zwart, J. A., Ledden, P. J., van Gelderen, P., Bodurka, J., Chu, R., & Duyn, J. H. (2004). Signal-to-noise ratio and parallel imaging performance of a 16-channel receive-only brain coil array at 3.0 Tesla. *Magnetic Resonance in Medicine*, 51, 22–26.
- Diba, K., & Buzsaki, G. (2007). Forward and reverse hippocampal place-cell sequences during ripples. *Nature Neuroscience*, 10, 1241–1242.
- Foster, D. J., & Wilson, M. A. (2006). Reverse replay of behavioural sequences in hippocampal place cells during the awake state. *Nature*, 440, 680–683.
- Fox, M. D., & Greicius, M. (2010). Clinical applications of resting state functional connectivity. *Frontiers in Systems Neuroscience*, 4, 19.
- Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E. (2005). The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 9673–9678.
- Fox, M. D., Snyder, A. Z., Vincent, J. L., & Raichle, M. E. (2007). Intrinsic fluctuations within cortical systems account for intertrial variability in human behavior. *Neuron*, 56, 171–184.
- Fransson, P. (2005). Spontaneous low-frequency BOLD signal fluctuations: An fMRI investigation of the resting-state default mode of brain function hypothesis. *Human Brain Mapping*, 26, 15–29.
- Fukunaga, M., Horovitz, S. G., van Gelderen, P., de Zwart, J. A., & Duyn, J. H. (2005). Brain metabolic activity in absence of stimuli. *Proceedings of VIIIth International Conference on Quantification of Brain Function with PET*, 26, S63 June 7–11.
- Fukunaga, M., Horovitz, S. G., van Gelderen, P., de Zwart, J. A., & Duyn, J. H. (2008). Brain metabolic activity in absence of stimuli. *Journal of Cerebral Blood Flow & Metabolism*, 28, 1377–1387.
- Fukunaga, M., Horovitz, S. G., van Gelderen, P., de Zwart, J. A., Jansma, J. M., Ikonomidou, V. N., et al. (2006). Large-amplitude, spatially correlated fluctuations in BOLD fMRI signals during extended rest and early sleep stages. *Magnetic Resonance Imaging*, 24, 979–992.
- Glover, G. H., Li, T. Q., & Ress, D. (2000). Image-based method for retrospective correction of physiological motion effects in fMRI: RETROICOR. *Magnetic Resonance in Medicine*, 44, 162–167.
- Golanov, E. V., Yamamoto, S., & Reis, D. J. (1994). Spontaneous waves of cerebral blood flow associated with a pattern of electrocortical activity. *The American Journal of Physiology*, 266, R204–214.
- Goldman, R. I., Stern, J. M., Engel, J., Jr. & Cohen, M. S. (2002). Simultaneous EEG and fMRI of the alpha rhythm. *Neuroreport*, 13, 2487–2492.
- Hahn, T. T., Sakmann, B., & Mehta, M. R. (2006). Phase-locking of hippocampal interneurons' membrane potential to neocortical up-down states. *Nature Neuroscience*, 9, 1359–1361.
- Hampson, M., Peterson, B. S., Skudlarski, P., Gatenby, J. C., & Gore, J. C. (2002). Detection of functional connectivity using temporal correlations in MR images. *Human Brain Mapping*, 15, 247–262.
- He, B. J., Snyder, A. Z., Zempel, J. M., Smyth, M. D., & Raichle, M. E. (2008). Electrophysiological correlates of the brain's intrinsic large-scale functional architecture. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 16039–16044.
- Hoge, R. D., Atkinson, J., Gill, B., Crelier, G. R., Marrett, S., & Pike, G. B. (1999). Linear coupling between cerebral blood flow and oxygen consumption in activated human cortex. *Proceedings of the National Academy of Sciences of the United States of America*, 96, 9403–9408.
- Honey, C. J., Sporns, O., Cammoun, L., Gigandet, X., Thiran, J. P., Meuli, R., et al. (2009). Predicting human resting-state functional connectivity from structural connectivity. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 2035–2040.
- Horovitz, S. G., Braun, A. R., Carr, W. S., Picchioni, D., Balkin, T. J., Fukunaga, M., et al. (2009). Decoupling of the brain's default mode network during deep sleep.

- Proceedings of the National Academy of Sciences of the United States of America*, 106, 11376–11381.
- Horovitz, S. G., Fukunaga, M., de Zwart, J. A., van Gelderen, P., Fulton, S. C., Balkin, T. J., et al. (2008). Low frequency BOLD fluctuations during resting wakefulness and light sleep: A simultaneous EEG-fMRI study. *Human Brain Mapping*, 29, 671–682.
- Huber, R., Ghilardi, M. F., Massimini, M., & Tononi, G. (2004). Local sleep and learning. *Nature*, 430, 78–81.
- Hyde, J. S., Biswal, B. B., & Jesmanowicz, A. (2001). High-resolution fMRI using multislice partial k-space GR-EPI with cubic voxels. *Magnetic Resonance in Medicine*, 46, 114–125.
- Isomura, Y., Sirota, A., Ozen, S., Montgomery, S., Mizuseki, K., Henze, D. A., et al. (2006). Integration and segregation of activity in entorhinal-hippocampal subregions by neocortical slow oscillations. *Neuron*, 52, 871–882.
- Ji, D., & Wilson, M. A. (2007). Coordinated memory replay in the visual cortex and hippocampus during sleep. *Nature Neuroscience*, 10, 100–107.
- Johnston, J. M., Vaishnavi, S. N., Smyth, M. D., Zhang, D., He, B. J., Zempel, J. M., et al. (2008). Loss of resting interhemispheric functional connectivity after complete section of the corpus callosum. *The Journal of Neuroscience*, 28, 6453–6458.
- Kanwisher, N. (2001). Neural events and perceptual awareness. *Cognition*, 79, 89–113.
- Kenet, T., Bibitchkov, D., Tsodyks, M., Grinvald, A., & Arieli, A. (2003). Spontaneously emerging cortical representations of visual attributes. *Nature*, 425, 954–956.
- Kiviniemi, V., Jauhainen, V., Tervonen, O., Paakko, E., Oikarinen, J., Vainionpaa, V., et al. (2000). Slow vasomotor fluctuation in fMRI of anesthetized child brain. *Magnetic Resonance in Medicine*, 44, 373–378.
- Kosslyn, S. M., Pascual-Leone, A., Felician, O., Camposano, S., Keenan, J. P., Thompson, W. L., et al. (1999). The role of area 17 in visual imagery: Convergent evidence from PET and rTMS. *Science*, 284, 167–170.
- Krienen, F. M., & Buckner, R. L. (2009). Segregated frontocerebellar circuits revealed by intrinsic functional connectivity. *Cerebral Cortex*, 19, 2485–2497.
- Larson-Prior, L. J., Zempel, J. M., Nolan, T. S., Prior, F. W., Snyder, A. Z., & Raichle, M. E. (2009). Cortical network functional connectivity in the descent to sleep. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 4489–4494.
- Laufs, H., Holt, J. L., Elfont, R., Krams, M., Paul, J. S., Krakow, K., et al. (2006). Where the BOLD signal goes when alpha EEG leaves. *NeuroImage*, 31, 1408–1418.
- Laufs, H., Kleinschmidt, A., Beyerle, A., Eger, E., Salek-Haddadi, A., Preibisch, C., et al. (2003a). EEG-correlated fMRI of human alpha activity. *NeuroImage*, 19, 1463–1476.
- Laufs, H., Krakow, K., Sterzer, P., Eger, E., Beyerle, A., Salek-Haddadi, A., et al. (2003b). Electroencephalographic signatures of attentional and cognitive default modes in spontaneous brain activity fluctuations at rest. *Proceedings of the National Academy of Sciences of the United States of America*, 100, 11053–11058.
- Lewis, C. M., Baldassarre, A., Committeri, G., Romani, G. L., & Corbetta, M. (2009). Learning sculpts the spontaneous activity of the resting human brain. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 17558–17563.
- Liu, Z., Fukunaga, M., de Zwart, J. A., & Duyn, J. H. (2010). Large-scale spontaneous fluctuations and correlations in brain electrical activity observed with magnetoencephalography. *NeuroImage*, 51, 102–111.
- Liu, X., Zhu, X. H., Zhang, Y., & Chen, W. (2011). Neural Origin of spontaneous hemodynamic fluctuations in rats under burst-suppression anesthesia condition. *Cerebral Cortex*, 21, 374–384.
- Logothetis, N. K. (2002). The neural basis of the blood-oxygen-level-dependent functional magnetic resonance imaging signal. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 357, 1003–1037.
- Lu, H., Zuo, Y., Gu, H., Waltz, J. A., Zhan, W., Scholl, C. A., et al. (2007). Synchronized delta oscillations correlate with the resting-state functional MRI signal. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 18265–18269.
- Luczak, A., Bartho, P., Marguet, S. L., Buzsaki, G., & Harris, K. D. (2007). Sequential structure of neocortical spontaneous activity in vivo. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 347–352.
- Mantini, D., Perrucci, M. G., Del Gratta, C., Romani, G. L., & Corbetta, M. (2007). Electrophysiological signatures of resting state networks in the human brain. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 13170–13175.
- Maquet, P., Dive, D., Salmon, E., Sadzot, B., Franco, G., Poirrier, R., et al. (1990). Cerebral glucose utilization during sleep-wake cycle in man determined by positron emission tomography and [18 F]2-fluoro-2-deoxy-D-glucose method. *Brain Research*, 513, 136–143.
- Martuzzi, R., Ramani, R., Qiu, M., Rajeevan, N., & Constable, R. T. (2010). Functional connectivity and alterations in baseline brain state in humans. *NeuroImage*, 49, 823–834.
- Massimini, M., Huber, R., Ferrarelli, F., Hill, S., & Tononi, G. (2004). The sleep slow oscillation as a traveling wave. *The Journal of Neuroscience*, 24, 6862–6870.
- McKiernan, K. A., Kaufman, J. N., Kucera-Thompson, J., & Binder, J. R. (2003). A parametric manipulation of factors affecting task-induced deactivation in functional neuroimaging. *Journal of Cognitive Neuroscience*, 15, 394–408.
- Mena-Segovia, J., Sims, H. M., Magill, P. J., & Bolam, J. P. (2008a). Cholinergic brainstem neurons modulate cortical

- gamma activity during slow oscillations. *The Journal of Physiology*, 586, 2947–2960.
- Mena-Segovia, J., Winn, P., & Bolam, J. P. (2008b). Cholinergic modulation of midbrain dopaminergic systems. *Brain Research Reviews*, 58, 265–271.
- Miall, R. C., & Robertson, E. M. (2006). Functional imaging: Is the resting brain resting? *Current Biology*, 16, R998–1000.
- Mohajerani, M. H., McVea, D. A., Fingas, M., & Murphy, T. H. (2010). Mirrored bilateral slow-wave cortical activity within local circuits revealed by fast bihemispheric voltage-sensitive dye imaging in anesthetized and awake mice. *The Journal of Neuroscience*, 30, 3745–3751.
- Molle, M., Eschenko, O., Gais, S., Sara, S. J., & Born, J. (2009). The influence of learning on sleep slow oscillations and associated spindles and ripples in humans and rats. *The European Journal of Neuroscience*, 29, 1071–1081.
- Molle, M., Marshall, L., Gais, S., & Born, J. (2004). Learning increases human electroencephalographic coherence during subsequent slow sleep oscillations. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 13963–13968.
- Moosmann, M., Ritter, P., Krastel, I., Brink, A., Thees, S., Blankenburg, F., et al. (2003). Correlates of alpha rhythm in functional magnetic resonance imaging and near infrared spectroscopy. *NeuroImage*, 20, 145–158.
- Murphy, M., Riedner, B. A., Huber, R., Massimini, M., Ferrarelli, F., & Tononi, G. (2009). Source modeling sleep slow waves. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 1608–1613.
- Musizza, B., Stefanovska, A., McClintock, P. V., Palus, M., Petrovic, J., Ribaric, S., et al. (2007). Interactions between cardiac, respiratory and EEG-delta oscillations in rats during anaesthesia. *The Journal of Physiology*, 580, 315–326.
- Musso, F., Brinkmeyer, J., Mobascher, A., Warbrick, T., & Winterer, G. (2010). Spontaneous brain activity and EEG microstates. A novel EEG/fMRI analysis approach to explore resting-state networks. *NeuroImage*, 52, 1149–1161.
- Nir, Y., Hasson, U., Levy, I., Yeshurun, Y., & Malach, R. (2006). Widespread functional connectivity and fMRI fluctuations in human visual cortex in the absence of visual stimulation. *NeuroImage*, 30, 1313–1324.
- Ogawa, S., Lee, T. M., Kay, A. R., & Tank, D. W. (1990). Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proceedings of the National Academy of Sciences of the United States of America*, 87, 9868–9872.
- Peltier, S. J., Kerssens, C., Hamann, S. B., Sebel, P. S., Byas-Smith, M., & Hu, X. (2005). Functional connectivity changes with concentration of sevoflurane anesthesia. *Neuroreport*, 16, 285–288.
- Petersen, C. C., Hahn, T. T., Mehta, M., Grinvald, A., & Sakmann, B. (2003). Interaction of sensory responses with spontaneous depolarization in layer 2/3 barrel cortex. *Proceedings of the National Academy of Sciences of the United States of America*, 100, 13638–13643.
- Quigley, M., Cordes, D., Turski, P., Moritz, C., Haughton, V., Seth, R., et al. (2003). Role of the corpus callosum in functional connectivity. *AJNR. American Journal of Neuroradiology*, 24, 208–212.
- Raichle, M. E. (2009). A paradigm shift in functional brain imaging. *The Journal of Neuroscience*, 29, 12729–12734.
- Raos, V., & Bentivoglio, M. (1993). Crosstalk between the two sides of the thalamus through the reticular nucleus: A retrograde and anterograde tracing study in the rat. *The Journal of Comparative Neurology*, 332, 145–154.
- Sadaghiani, S., Hesselmann, G., Friston, K. J., & Kleinschmidt, A. (2010). The relation of ongoing brain activity, evoked neural responses, and cognition. *Frontiers in Systems Neuroscience*, 4, 20.
- Sadato, N., Nakamura, S., Ohashi, T., Nishina, E., Fuwamoto, Y., Waki, A., et al. (1998). Neural networks for generation and suppression of alpha rhythm: A PET study. *Neuroreport*, 9, 893–897.
- Salvador, R., Suckling, J., Coleman, M. R., Pickard, J. D., Menon, D., & Bullmore, E. (2005). Neurophysiological architecture of functional magnetic resonance images of human brain. *Cerebral Cortex*, 15, 1332–1342.
- Schabus, M., Dang-Vu, T. T., Albouy, G., Baletau, E., Boly, M., Carrier, J., et al. (2007). Hemodynamic cerebral correlates of sleep spindles during human non-rapid eye movement sleep. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 13164–13169.
- Scholvinck, M. L., Maier, A., Ye, F. Q., Duyn, J. H., & Leopold, D. A. (2010). Neural basis of global resting-state fMRI activity. *Proceedings of the National Academy of Sciences of the United States of America*, 107, 10238–10243.
- Shmuel, A., & Leopold, D. A. (2008). Neuronal correlates of spontaneous fluctuations in fMRI signals in monkey visual cortex: Implications for functional connectivity at rest. *Human Brain Mapping*, 29, 751–761.
- Shmueli, K., van Gelderen, P., de Zwart, J. A., Horovitz, S. G., Fukunaga, M., Jansma, J. M., et al. (2007). Low-frequency fluctuations in the cardiac rate as a source of variance in the resting-state fMRI BOLD signal. *NeuroImage*, 38, 306–320.
- Shoham, D., & Grinvald, A. (2001). The cortical representation of the hand in macaque and human area S-I: High resolution optical imaging. *The Journal of Neuroscience*, 21, 6820–6835.
- Sirota, A., Csicsvari, J., Buhl, D., & Buzsaki, G. (2003). Communication between neocortex and hippocampus during sleep in rodents. *Proceedings of the National Academy of Sciences of the United States of America*, 100, 2065–2069.
- Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., et al. (2009). Correspondence of the brain's functional architecture during activation and rest.

- Proceedings of the National Academy of Sciences of the United States of America*, 106, 13040–13045.
- Smith, A. M., Lewis, B. K., Ruttimann, U. E., Ye, F. Q., Sinnwell, T. M., Yang, Y., et al. (1999). Investigation of low frequency drift in fMRI signal. *NeuroImage*, 9, 526–533.
- Steriade, M., Nunez, A., & Amzica, F. (1993). Intracellular analysis of relations between the slow (<1 Hz) neocortical oscillation and other sleep rhythms of the electroencephalogram. *The Journal of Neuroscience*, 13, 3266–3283.
- Stevens, W. D., Buckner, R. L., & Schacter, D. L. (2010). Correlated low-frequency BOLD fluctuations in the resting human brain are modulated by recent experience in category-preferential visual regions. *Cerebral Cortex*, 20, 1997–2006.
- Tononi, G., & Cirelli, C. (2003). Sleep and synaptic homeostasis: A hypothesis. *Brain Research Bulletin*, 62, 143–150.
- Triantafyllou, C., Hoge, R. D., Krueger, G., Wiggins, C. J., Potthast, A., Wiggins, G. C., et al. (2005). Comparison of physiological noise at 1.5 T, 3 T and 7 T and optimization of fMRI acquisition parameters. *NeuroImage*, 26, 243–250.
- Tsodyks, M., Kenet, T., Grinvald, A., & Arieli, A. (1999). Linking spontaneous activity of single cortical neurons and the underlying functional architecture. *Science*, 286, 1943–1946.
- Vincent, J. L., Patel, G. H., Fox, M. D., Snyder, A. Z., Baker, J. T., Van Essen, D. C., et al. (2007). Intrinsic functional architecture in the anaesthetized monkey brain. *Nature*, 447, 83–86.
- Wheeler, M. E., Petersen, S. E., & Buckner, R. L. (2000). Memory's echo: Vivid remembering reactivates sensory-specific cortex. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 11125–11129.
- Zhang, H. Y., Wang, S. J., Liu, B., Ma, Z. L., Yang, M., Zhang, Z. J., et al. (2010). Resting brain connectivity: Changes during the progress of Alzheimer disease. *Radiology*, 256, 598–606.
- Zhou, J., Greicius, M. D., Gennatas, E. D., Growdon, M. E., Jang, J. Y., Rabinovici, G. D., et al. (2010). Divergent network connectivity changes in behavioural variant frontotemporal dementia and Alzheimer's disease. *Brain*, 133, 1352–1367.