

Definicja 1 Niech $\alpha_1, \alpha_2, \dots, \alpha_k \in T_p^*M \in \Lambda'(M)$, wówczas $\alpha_1 \wedge \alpha_2 \wedge \dots \wedge \alpha_k \in \Lambda^k(M)$ i dla $v_1, v_2, \dots, v_k \in T_p^*M$,

$$\langle \alpha_1 \wedge \alpha_2 \wedge \dots \wedge \alpha_k; v_1, v_2, \dots, v_k \rangle \stackrel{def}{=} \begin{bmatrix} \alpha_1(v_1)\alpha_2(v_1) \dots \alpha_k(v_1) \\ \vdots \\ \alpha_1(v_k)\alpha_2(v_k) \dots \alpha_k(v_k) \end{bmatrix}.$$

Uwagi do operatora d ($dd = 0$):

Niech $M = \mathbb{R}^3, f : \mathbb{R}^3 \rightarrow \mathbb{R}^1 \in \Lambda^0(M)$

$$\begin{aligned} df &= \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy + \frac{\partial f}{\partial z} dz \\ ddf &= d\left(\frac{\partial f}{\partial x}\right) \wedge dx + d\left(\frac{\partial f}{\partial y}\right) \wedge dy + d\left(\frac{\partial f}{\partial z}\right) \wedge dz = \\ &= \left(\frac{\partial^2 f}{\partial x^2} dx + \frac{\partial^2 f}{\partial y \partial x} dy + \frac{\partial^2 f}{\partial z \partial x} dz\right) \wedge dx + \left(\frac{\partial^2 f}{\partial x \partial z} dx + \frac{\partial^2 f}{\partial y \partial z} dy\right) \wedge dy \\ &\quad + \left(\frac{\partial^2 f}{\partial x \partial z} dx + \frac{\partial^2 f}{\partial y \partial z} dz\right) \wedge dz \\ &= \left(\frac{\partial^2 f}{\partial y \partial x} - \frac{\partial^2 f}{\partial x \partial y}\right) dy \wedge dx + \left(\frac{\partial^2 f}{\partial z \partial y} - \frac{\partial^2 f}{\partial y \partial z}\right) dz \wedge dy + \\ &= \left(\frac{\partial^2 f}{\partial z \partial x} - \frac{\partial^2 f}{\partial x \partial z}\right) dz \wedge dx = 0. \end{aligned}$$

Niech $\alpha = A_x dx + A_y dy + A_z dz$

$$\begin{aligned} d\alpha &= \left(\frac{\partial A_x}{\partial y} - \frac{\partial A_y}{\partial x}\right) dy \wedge dx + \left(\frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z}\right) dz \wedge dy + \\ &\quad + \left(\frac{\partial A_z}{\partial x} - \frac{\partial A_x}{\partial z}\right) dz \wedge dx \\ d d\alpha &= \left(\pm \left(\frac{\partial^2 A_x}{\partial z \partial y} - \frac{\partial^2 A_x}{\partial z \partial x}\right) \pm \left(\frac{\partial^2 A_z}{\partial x \partial y} - \frac{\partial^2 A_y}{\partial x \partial z}\right) \pm \left(\frac{\partial^2 A_z}{\partial y \partial x} - \frac{\partial^2 A_x}{\partial y \partial z}\right)\right) dx \wedge dy \wedge dz \\ &= 0. \end{aligned}$$

$$\beta = A_x dy \wedge dz + A_y dx \wedge dz + A_z dy \wedge dx$$

$$d\beta = 0$$

$$dd\beta = 0.$$

Niech $M = \mathbb{R}^4$, $A = \phi dt + A_x dx + A_y dy + A_z dz$.

$$dA = \underbrace{\left(\frac{\partial \phi}{\partial x} - \frac{\partial A_x}{\partial t} \right)}_{E_x} dx \wedge dt + \underbrace{\left(\frac{\partial \phi}{\partial y} - \frac{\partial A_y}{\partial t} \right)}_{E_y} dy \wedge dt + \underbrace{\left(\frac{\partial \phi}{\partial z} - \frac{\partial A_z}{\partial t} \right)}_{E_z} dz \wedge dt +$$

$$\underbrace{\left(\frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right)}_{B_z} dx \wedge dy + \underbrace{\left(\frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right)}_{B_x} dy \wedge dz + \underbrace{\left(\frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right)}_{B_y} dz \wedge dx$$

$$ddA = 0.$$

niech $dA = F$

$$dF = 0.$$

Pytanie: niech M - rozmaitość wymiaru 3 (bo mamy bijekcję między $\theta \in M$ i \mathbb{R}^3). Czy istnieje $\Lambda^4(M)$?

niech $M = \mathbb{R}^3$

$\Lambda^0(M)$	$f : \mathbb{R}^3 \rightarrow M$	$\dim \Lambda^1(M) = 3$
$\Lambda^1(M)$	$\alpha = A_x dx + A_y dy + A_z dz$	$\Lambda^1(\eta) = \langle dx, dy, dz \rangle$
$\Lambda^2(M)$	$\beta = A_z dx \wedge dy + A_y dz \wedge dx + A_x dy \wedge dz$	$\Lambda^2(M) = \langle dx \wedge dy, dz \wedge dx, dy \wedge dz \rangle$
$\dim(\Lambda^2(M)) = 3$		
$\Lambda^3(\eta)$	$\gamma = f dx \wedge dy \wedge dz$	$\Lambda^3(M) = \langle dx \wedge dy \wedge dz \rangle$
$\dim(\Lambda^3(M)) = 1.$		

Niech $M = \mathbb{R}^4$.

$\Lambda^0(M)$	$f(t, x, y, z) \rightarrow \mathbb{R}$	$\dim \Lambda^0(M) = 1$
$\Lambda^1(M)$	$\alpha = A_t dt + A_x dx + A_y dy + A_z dz$	$\dim \Lambda^1(M) = 4$
$\Lambda^2(M)$	$\beta = A_1 dt \wedge dx + A_2 dt \wedge dy + A_3 dt \wedge dz + B_1 dy \wedge dx + B_2 dz \wedge dx + C_1 dz \wedge dy$	$\dim \Lambda^2(M) = 6$
$\Lambda^3(M) :$	$\gamma = C_1 dy \wedge dt \wedge dx + C_2 dz \wedge dt \wedge dx + D_1 dz \wedge dt \wedge dy + E_1 dx \wedge dy \wedge dz$	$\dim \Lambda^3(M) = 4$
$\Lambda^4(M)$	$\delta = g dt \wedge dx \wedge dy \wedge dz$	$\dim \Lambda^4(M) = 1.$

0.1 Pchnięcia i cofnięcia

Niech M, N - rozmaitości $\dim M = n, \dim N = k$ i niech $h : M \rightarrow N$. (h nie musi być bijekcją !!!)



Niech $p \in M$. Pchnięciem punktu p w odwzorowaniu h nazywamy punkt $h_*(p) \stackrel{\text{def}}{=} h(p)$


Przykład 1 Niech $M = \mathbb{R}^2$, $N = \mathbb{R}$, $h(x, y) = x + y, h : \mathbb{R}^2 \rightarrow \mathbb{R}$.

$$p = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, h_*(p) = 3$$

$$M = \mathbb{R}^1, N = \mathbb{R}^3, h(t) = \begin{bmatrix} \cos t \\ \sin t \\ t \end{bmatrix}, p = \frac{\pi}{2}.$$

$$h_x(\frac{\pi}{2}) = \begin{bmatrix} \cos \frac{\pi}{2} \\ \sin \frac{\pi}{2} \\ \frac{\pi}{2} \end{bmatrix}$$





Niech $\sigma(t)$ - krzywa na M . Pchnięciem krzywej σ w odwzorowaniu h nazywamy krzywą $h_*(\sigma(t)) \stackrel{\text{def}}{=} h(\sigma(t))$

Niech $f : N \rightarrow \mathbb{R}^2$. Cofnięciem funkcji f w odwzorowaniu h nazywamy funkcję

$$h^* f(p) = f(h(p)).$$

Przykład 2 $M = \mathbb{R}^2, N = \mathbb{R}, f : N \rightarrow \mathbb{R}^2, f(t) = \begin{bmatrix} 2t \\ t \end{bmatrix}, h(x, y) = x + y$.

$$h^* f(x, y) = f(h(x, y)) = \begin{bmatrix} 2(x + y) \\ x + y \end{bmatrix}.$$

Pchnięciem wektora V w odwzorowaniu h nazywamy wektor

$$h_* V = [h(\sigma)], h_* v \in T_{h(p)} N.$$

Przykład 3 Niech $M = \mathbb{R}^2, N = \mathbb{R}, h(x, y) = x + 2y, v = 2\frac{\partial}{\partial x} + 3\frac{\partial}{\partial y}$. Co to jest $h_* v$?
 $p = (1, 2) = (\varphi^1(p), \varphi^2(p))$

$$\sigma(t) : \frac{d}{dt}(\varphi(\sigma(t)))|_{t=0}$$

$$\varphi(\sigma(t)) = \begin{bmatrix} 2t+1 \\ 3t+2 \end{bmatrix}$$

$$h[\sigma(t)] = 2t+1+2(3t+2)$$

$$h[\sigma(t)] = 8t+5$$

$$[h[\sigma(t)]] = 8 \frac{\partial}{\partial t} \in t_s N.$$

$$\dim M = n, \varphi(\sigma(t)) = (\varphi^1(\sigma(t)), \varphi^2(\sigma(t)), \dots, \varphi^n(\sigma(t))), v \in T_p M.$$

$$v = \frac{\partial \varphi^1(\sigma(t))}{\partial t} \Big|_{t=0} \frac{\partial}{\partial x^1} + \frac{\partial \varphi^2(\sigma(t))}{\partial t} \Big|_{t=0} \frac{\partial}{\partial x^2} \cdots \frac{\partial \varphi^n(\sigma(t))}{\partial t} \Big|_{t=0} \frac{\partial}{\partial x^n}.$$


$$\frac{d(\varphi \circ h(\sigma(t)))}{dt} \Big|_{t=0} = \frac{d}{dt} (\psi \circ h \circ \varphi^{-1} \sigma)_{t=0} = \frac{d}{dt} (\tilde{h} \circ \tilde{\sigma}(t)).$$

$$= \frac{d}{dt} \tilde{h}(\tilde{\sigma}_1(t), \tilde{\sigma}_2(t), \dots, \tilde{\sigma}^n(t))_{t=0} = \tilde{h}'_{\tilde{\sigma}(0)} \frac{d\tilde{\sigma}}{dt} \Big|_{t=0} = \tilde{h}' \cdot v.$$

$$\text{Czyli ostatecznie } v = \begin{bmatrix} 2 \\ 3 \end{bmatrix}_{\frac{\partial}{\partial x}, \frac{\partial}{\partial y}}, \tilde{h}(x, y) = x + 2y \rightarrow \tilde{h}(x, y) = [1, 2].$$

$$h_* v = [1, 2] \begin{bmatrix} 2 \\ 3 \end{bmatrix} = 2 \cdot 1 + 6 = 8 \frac{\partial}{\partial t}.$$

Niech $\alpha \in \Lambda^1(?)$ - pytanie: czy formy się pcha, czy cofa?



Rysunek 1: $\tilde{h} = \psi h \varphi^{-1}$