

Stability of Minimal Surfaces

A Study on Scherk Surface

K. Oskar Negron

Department of Economics, CSU Fullerton

August 26, 2016

Outline

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction

basics

First
Fundamental
Form

Complex
Analysis

Basics

Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second
Variation

Theorem
S.V. for Scherk
surface

1 Introduction

2 Complex Analysis

3 Scherk surface

4 Second Variation

Area

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction
basics
First
Fundamental
Form

Complex
Analysis
Basics
Weierstrass-
Enneper
Representation

Scherk surface
Parametrization
Plots

Second
Variation
Theorem
S.V. for Scherk
surface

Definition

The Area $A(R)$ of the part $\sigma(R)$ of the surface patch $\sigma : U \rightarrow \mathbb{R}^3$ corresponding to a region $R \subset U$ is

$$\begin{aligned} A(R) &= \int_R \|\sigma_u \wedge \sigma_v\| du dv \\ &= \int_R \sqrt{(\sigma_u \cdot \sigma_u)(\sigma_v \cdot \sigma_v) - (\sigma_u \cdot \sigma_v)^2} du dv \\ &= \int_R \sqrt{EG - F^2} du dv. \end{aligned}$$

Weingarten map

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction

basics

First
Fundamental
Form

Complex
Analysis

Basics

Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second

Variation

Theorem

S.V. for Scherk
surface

Definition

Let σ be a regular curve, i.e. $\sigma_u \wedge \sigma_v$ is non-vanishing. The Weingarten map $S = S(u, v)$ is a linear map of the tangent space T_p into itself defined as follows: if $a = a_1\sigma_u + a_2\sigma_v$ then,

$$Sa = -a_1N_u - a_2N_v.$$

S is also known as the shape operator.

Fundamental Forms

Definition

We can define the following three symmetric bilinear forms:

$$I(v, w) = v \cdot w$$

$$II(v, w) = S v \cdot w$$

$$III(v, w) = S v \cdot S w.$$

We will use the following notation, when appropriate, for the coefficients of the fundamental forms:

$$g_{ij} = \sigma_i \cdot \sigma_j$$

$$b_{ij} = -N_i \cdot \sigma_j$$

$$c_{ij} = N_i \cdot N_j.$$

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction
basics
First
Fundamental
Form

Complex
Analysis

Basics
Weierstrass-
Enneper
Representation

Scherk surface
Parametrization
Plots

Second
Variation
Theorem
S.V. for Scherk
surface

Notation

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction

basics

First
Fundamental
Form

Complex
Analysis

Basics
Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second
Variation

Theorem
S.V. for Scherk
surface

$$g_{ij} = (g_{ij}) = \begin{bmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{bmatrix} = \begin{bmatrix} E & F \\ F & G \end{bmatrix}$$

$$b_{ij} = (b_{ij}) = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} L & M \\ M & N \end{bmatrix}$$

$$W = \sqrt{g} = \sqrt{EG - F^2} = \sqrt{|g_{ij}|}$$

Curvature

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction

basics

First
Fundamental
Form

Complex
Analysis

Basics

Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second
Variation

Theorem
S.V. for Scherk
surface

The Gaussian curvature K is given by

$$K = k_1 k_2 = |b_i^j| = |g^{jk} b_{ki}|$$

and the mean curvature H is given by

$$2H = k_1 + k_2 = b_{ij} g^{ij}.$$

First Variation

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction
basics
First
Fundamental
Form

Complex
Analysis

Basics
Weierstrass-
Enneper
Representation

Scherk surface
Parametrization
Plots

Second
Variation
Theorem
S.V. for Scherk
surface

Theorem

The surface $\sigma : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$ is a critical point of the area function A if and only if its mean curvature H is identically zero.

Proof outline. Choose a bounded domain $D \subset U$ and a differentiable function $\phi : \bar{D} \rightarrow \mathbb{R}$ where $\bar{D} = D \cup \partial D$ and $(u, v) \in D$. Define

$$\tilde{\sigma} = \sigma + t\phi\mathbf{N}$$

for some fixed $t \in (-\epsilon, \epsilon)$. Let \tilde{g} and g be the metric tensor associated with $\tilde{\sigma}$ and σ , respectively.

First Variation Proof

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction

basics

First
Fundamental
Form

Complex
Analysis

Basics

Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second
Variation

Theorem
S.V. for Scherk
surface

$$A(\tilde{\sigma}) = \int_{\bar{D}} \sqrt{\tilde{g}_{11}\tilde{g}_{22} - \tilde{g}_{12}^2} \, du \, dv.$$

where

$$\begin{aligned}\tilde{g}_{11} &= \tilde{\sigma}_u \cdot \tilde{\sigma}_u \\ &= (\sigma_u + t\phi_u \mathbf{N} + t\phi \mathbf{N}_u)^2 \\ &= \sigma_u^2 + 2t\phi \langle \mathbf{N}_u | \sigma_u \rangle + O(t^2) \\ &= g_{11} + 2t\phi b_{11} + O(t^2),\end{aligned}$$

$$\tilde{g}_{12} = g_{12} + t\phi(\langle \mathbf{N}_u | \sigma_v \rangle + \langle \mathbf{N}_v | \sigma_u \rangle) + O(t^2).$$

and

$$\tilde{g}_{22} = g_{22} + 2t\phi b_{22} + O(t^2),$$

First Variation Proof

Stability of
Minimal
Surfaces

K. Oskar
Negron

$$\tilde{g}_{11}\tilde{g}_{22} - \tilde{g}_{12}^2 = g_{11}g_{22} - g_{12}^2 + 4t\phi W^2 H + O(t^2).$$

The are function is then,

$$A(\tilde{\sigma}) = \int_{\bar{D}} \sqrt{W^2(1 + 4t\phi H)} \, du \, dv + O(t^2)$$

and using the expansion $\sqrt{1+x} = 1 + \frac{x}{2} + O(x^2)$, then,

$$\begin{aligned} A(\tilde{\sigma}) &= \int_{\bar{D}} W(1 + 2t\phi H) \, du \, dv + O(t^2) \\ &= \int_{\bar{D}} W \, du \, dv + 2t \int_{\bar{D}} HW\phi \, du \, dv + O(t^2) \end{aligned}$$

First Variation Proof

Taking the derivative with respect to t ,

$$D[A(\sigma)](\phi) = \frac{d}{dt} A(\tilde{\sigma}) = \int_{\bar{D}} 2\phi HW \, du \, dv.$$

If $D[A(\sigma)](\phi) = 0$ then $\int_{\bar{D}} 2\phi HW \, du \, dv = 0$. Since this is true for every ϕ , we can choose $\phi = H$ and so

$$\int_{\bar{D}} H^2 W \, du \, dv = 0.$$

However, since $H^2 \geq 0$ everywhere we must have $H = 0$.

Definition

We say σ is a minimal surface if it has zero mean curvature.

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction

basics

First
Fundamental
Form

Complex
Analysis

Basics

Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second

Variation

Theorem

S.V. for Scherk
surface

Holomorphic function

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction
basics
First
Fundamental
Form

Complex
Analysis

Basics
Weierstrass-
Enneper
Representation

Scherk surface
Parametrization
Plots

Second
Variation

Theorem
S.V. for Scherk
surface

Definition

Holomorphic function. If a complex function $f(z) = f(x + iy) = u(x, y) + iv(x, y)$ is holomorphic (or analytical), then u and v have first partial derivatives with respect to x and y , and satisfy the Cauchy-Riemann equations:

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Complex Analysis

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction

basics

First
Fundamental
Form

Complex
Analysis

Basics

Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second

Variation

Theorem

S.V. for Scherk
surface

Theorem

Let $\sigma : U \rightarrow \mathbb{R}^3$ be a conformal surface patch. Consider the complex coordinates in the plane of which U is an open subset by setting $z = u + iv$ for $(u, v) \in U$. We define $\phi(z) = \sigma_u - i\sigma_v$ and we say σ is minimal if and if the function ϕ is holomorphic on U .

Complex Analysis

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction
basics
First
Fundamental
Form

Complex
Analysis

Basics
Weierstrass-
Enneper
Representation

Scherk surface
Parametrization
Plots

Second
Variation
Theorem
S.V. for Scherk
surface

Theorem

If $\sigma : U \rightarrow \mathbb{R}^3$ is a conformal minimal surface, the vector-valued holomorphic function $\phi = (\phi_1, \phi_2, \phi_3)$, defined in the previous slide, satisfies the following:

(a.) ϕ is nowhere zero

(b.) $\sum \phi_k^2 = 0$ for $k = 1, 2, 3$.

Conversely, if U is simple-connected and if ϕ_k are holomorphic functions on U satisfying (a.) and (b.), there is a conformally parametrized minimal surface $\sigma : U \rightarrow \mathbb{R}^3$ such that ϕ satisfies $\phi(z) = \sigma_u - i\sigma_v$. Moreover, σ is uniquely determined by ϕ_k up to a translation.

Weierstrass-Enneper Representation

Theorem

Weierstrass-Enneper Representation. *Let $f(z)$ be a holomorphic function on an open set $U \subset \mathbb{C}$, not identically zero, and let $g(z)$ be a meromorphic function on U such that if $z_0 \in U$ is a m^{th} pole of g , for $m \geq 1$, then z_0 is also a zero of f of order greater than $2m$. Then,*

$$\phi = \left(\frac{1}{2}f(1 - g^2), \frac{i}{2}f(1 + g^2), fg \right)$$

and it satisfies the previous theorem. Conversely, every holomorphic function ϕ satisfying these conditions arises in this way.

Observe that if $\phi_1 - i\phi_2$ is not identically zero, we can define

$$f = \phi_1 - i\phi_2 \text{ and } g = \frac{\phi_3}{\phi_1 - i\phi_2}.$$

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction
basics
First
Fundamental
Form

Complex
Analysis
Basics
Weierstrass-
Enneper
Representation

Scherk surface
Parametrization
Plots

Second
Variation
Theorem
S.V. for Scherk
surface

Parametrization

Schrek's surface can be given by the parametric equation

$$\sigma(u, v) = \left(\arg \frac{z + i}{z - i}, \arg \frac{z + 1}{z - 1}, \log \left| \frac{z^2 + 1}{z^2 - 1} \right| \right)$$

with $z \neq \{\pm 1, \pm i\}$ and where $\arg z$ is the angle that the real axis makes with z .

We can construct a complex function $\phi_k = (\sigma^k)_u - i(\sigma^k)_v$

$$\phi_1 = -\frac{2}{1 + z^2}, \phi_2 = -\frac{2i}{1 - z^2}, \text{ and } \phi_3 = \frac{4z}{1 - z^4}$$

and from this we get the Weierstrass-Enneper data

$$f = \frac{4}{z^4 - 1} \text{ and } g = -z.$$

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction

basics

First
Fundamental
Form

Complex
Analysis

Basics

Weierstrass-
Enneper
Representation

Schrek surface

Parametrization
Plots

Second

Variation

Theorem

S.V. for Schrek
surface

Plot

Stability of Minimal Surfaces

K. Oskar
Negron

Introduction

basics
First
Fundamental
Form

Complex Analysis


Basics
Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second Variation

Theorem
S.V. for Scherk
surface



Plot

Stability of Minimal Surfaces

K. Oskar
Negron

Introduction

basics
First
Fundamental
Form

Complex Analysis


Basics
Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second Variation

Theorem
S.V. for Scherk
surface



Second Variation

Stability of
Minimal
Surfaces

K. Oskar
Negron

Introduction

basics
First
Fundamental
Form

Complex
Analysis

Basics
Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second
Variation

Theorem
S.V. for Scherk
surface

Theorem

Suppose $\sigma = \sigma(u, v)$ is regular in the closure of Ω , then

$$\begin{aligned} D^2[A(\sigma)](\phi) &= \frac{d^2}{dt^2} A(\tilde{\sigma}) \\ &= \int_{\Omega} (|\nabla_u \phi|^2 + 2K\phi^2) \sqrt{|g_{ij}|} \, du \, dv \\ &= \int_{\Omega} (|\nabla_u \phi|^2 + 2K\phi^2) \, dA \\ &= \int_{\Omega} (|\nabla \phi|^2 + 2KW\phi^2) \, du \, dv. \end{aligned}$$

Code

Stability of Minimal Surfaces

K. Oskar
Negron

Introduction

basics
First
Fundamental
Form

Complex Analysis

Basics
Weierstrass-
Enneper
Representation

Scherk surface

Parametrization
Plots

Second Variation

Theorem
S.V. for Scherk
surface

We found that $\phi = K$ destabilizes the integral,

$$\int_{\Omega} (\phi \frac{1}{\sqrt{|(g_{jk})|}} \sum_{j,k} \partial_j (g^{jk} \sqrt{|(g_{jk})|} \partial_k \phi) + 2 |g^{jk} b_{ki}| \sqrt{|(g_{ij})|} \phi^2) dudv$$

$$\approx -2.4439.$$