


Malaria

#NigeriaDHS



2024 Nigeria Demographic and Health Survey (NDHS)


- **Malaria prevention**
- Malaria in pregnancy
- Case management of fever in children



Find these data in
Chapter 12
of the Final Report

Trends in ITN Ownership

Percent of households with at least 1 insecticide-treated net (ITN)




Note: An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In the 2008 NDHS, 2010 NMIS, 2013 NDHS, and 2015 NMIS, this was known as a long-lasting insecticidal net (LLIN).

Ownership of Mosquito Nets by Residence


Percent of households with at least:

■ Nigeria ■ Urban ■ Rural



ITN Ownership by Wealth


Percent of households with at least 1 ITN



ITN Ownership by State


*Percent of households
with at least 1 ITN*

Nigeria
59%



Full Household Coverage of ITNs


Percent distribution of households



ITN Access and Use by Residence


Percent of household population with access to an ITN and percent who slept under an ITN the night before the survey

■ Access to an ITN ■ Slept under an ITN



Trends in ITN Access and Use


Percent of household population with access to an ITN and percent who slept under an ITN the night before the survey




Note: An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In the 2008 NDHS, 2010 NMIS, 2013 NDHS, and 2015 NMIS, this was known as a long-lasting insecticidal net (LLIN).

ITN Use

*Percent who slept under an ITN the night before the survey among **all households***




*Percent who slept under an ITN the night before the survey among **households with at least one ITN***



Trends in ITN Use among Children and Pregnant Women

Percent of children under age 5 and pregnant women age 15-49 who slept under an ITN the night before the survey




Note: An insecticide-treated net (ITN) is a factory-treated net that does not require any further treatment. In the 2008 NDHS, 2010 NMIS, 2013 NDHS, and 2015 NMIS, this was known as a long-lasting insecticidal net (LLIN).


ITN Use Among Children by State

Percent of children under 5 who slept under an ITN the night before the survey

Nigeria
43%




- Malaria Prevention
- **Malaria in pregnancy**
- Case management of fever in children



Find these data in
Chapter 12
Of the Final Report


Intermittent Preventive Treatment during Pregnancy (IPTp)

Percent of women age 15-49 with a live birth and/or stillbirth in the 2 years before the survey who received at least 1, 2, or 3 doses of SP/Fansidar




Trends in IPTp

Percent of women age 15-49 with a live birth and/or stillbirth in the 2 years before the survey who received at least 1, 2, or 3 doses of SP/Fansidar



- Malaria prevention
- Malaria in pregnancy
- **Case management of fever in children**




Find these data in
Chapter 12
of the Final Report

Fever in Children


*Percent of
children under 5*

*Percent of children under 5 with fever
in the 2 weeks before the survey*



Type of Antimalarial Drugs Used

Among children under age 5 with fever in the 2 weeks before the survey who took an antimalarial, percent who took:



Key Findings

- **59%** of households own at least **1 ITN**.
- **48%** of the household population has **access to an ITN**.
- **55%** of **household members**, **61%** of **children under age 5**, and **66%** of **pregnant women slept under an ITN** the night before the survey.
- **Advice or treatment** was sought for **60%** of children under age 5 with fever in the 2 weeks before the survey.