

RESEARCH ARTICLE

Open Access



What makes a successful species? Traits facilitating survival in altered tropical forests

Mareike Hirschfeld* and Mark-Oliver Rödel

Abstract

Background: Ongoing conversion, disturbance and fragmentation of tropical forests stress this ecosystem and cause the decline or disappearance of many species. Particular traits have been identified which indicate an increasing extinction risk of a species, but traits facilitating survival in altered habitats have mostly been neglected. Here we search for traits that make a species tolerant to disturbances, thus independent of pristine forests. We identify the fauna that have an increasing effect on the ecosystem and its functioning in our human-dominated landscapes.

Methods: We use a unique set of published data on the occurrences of 243 frog species in pristine and altered forests throughout the tropics. We established a forest dependency index with four levels, based on these occurrence data and applied Random Forest classification and binomial Generalized Linear Models to test whether species life history traits, ecological traits or range size influence the likelihood of a species to persist in disturbed habitats.

Results: Our results revealed that indirect developing species exhibiting a large range size and wide elevational distribution, being independent of streams, and inhabiting the leaf litter, cope best with modifications of their natural habitats.

Conclusion: The traits identified in our study will likely persist in altered tropical forest systems and are comparable to those generally recognized for a low species extinction risk. Hence our findings will help to predict future frog communities in our human-dominated world.

Keywords: Forest degradation, Frogs, Life-history traits, Adaptation, Extinction risk, Tropics

Background

The anthropogenic conversion of natural environments, in particular of forest habitats, is a major threat to tropical biodiversity [1]. Beside the intensive loss of forest cover [2], fragmentation of the pristine remnants further affects species [3] and limits their ability to move into adequate areas. Thus the ability to cope with altered landscapes is crucial for the persistence of a species, especially in the face of climate change.

Numerous empirical and comparative approaches on species response to environmental changes and studies relating species properties to their extinction risk were conducted on invertebrates e.g. [4–6] as well as vertebrates e.g. [7–10]. However, the general pattern which

leads to the persistence of some species but the decrease or loss of other species due to forest disturbances is not fully understood. In different taxonomic groups some life-history and ecological traits show parallel patterns in their response to forest alteration, e.g. small range size [8, 10, 11] or low fecundity [12, 13] that lead to higher extinction risks. Whereas other traits, like body size exhibit a fuzzy prediction of a species' risk to decline in fragmented habitats [summary in 14]. The susceptibility of species is not determined by a single trait, but by a combination of properties which lead to a species-specific extinction risk [15–17]. So far, the majority of studies have focused on species affected by environmental changes and filter for traits increasing the extinction risk. Species not responding to habitat alterations and the traits required for their persistence in disturbed landscapes are frequently neglected. However, those species remaining are of high interest as they will make up the

*Correspondence: mail@mareikepetersen.de
Department Diversity Dynamics, Museum für Naturkunde Berlin-Leibniz
Institute for Evolutionary and Biodiversity Science, Invalidenstraße 43,
10115 Berlin, Germany

majority of the fauna in our human-dominated world and thus have an increasing effect on ecosystems and their functioning [18, 19].

Frogs are strongly influenced by their environment and the degradation and conversion of natural forests is one major cause for their current global decline [20–22]. However, not all species are affected by degradation or fragmentation [23–25] and a set of life-history or ecological traits is assumed to reduce their susceptibility [8, 26].

In this study, we search for factors allowing a species to be independent of pristine areas and thus permitting their occurrence in degraded and disturbed forests, which are the dominant tropical habitats now and in future [27]. We use a unique data set comprising published records on frog species occurrences in tropical forests, forest fragments and more intense altered landscapes such as plantations or settlements. For these species we gathered life-history (e.g. body size, clutch size) and ecological traits (e.g. habitat use) as well as distribution data, which are known to affect the susceptibility of species in general [8, 26, 28] and thus might likewise influence a species response to forest degradation. We ask whether these candidate traits could predict the forest dependency of tropical frog species and whether a particular set of traits makes species less vulnerable to changes in their natural habitat and decreases their risk of extinction.

Methods

Data acquisition

We combined a comprehensive data set on anuran occurrences across tropical forests and human altered forest habitats with detailed information on species traits. To cover all research published on anuran distribution in pristine versus altered environments in the tropics, we did a comprehensive literature research using Google, Google Scholar, Web of Science and data bases included therein (January to August 2013). Queries using different combinations of appropriate keywords (e.g. frog, amphibian, anuran, disturbance, alteration, fragmentation, logging etc.) were applied to all data bases. Appropriate data sets covered a description of the study sites and information on the presence (and absence) of each species in the different habitat types. In addition to already published studies we added our own data on anuran occurrences from the forest zone of Cameroon (M. Hirschfeld et al. unpublished data). The survey amounted to 61 studies (see Additional file 1: anuran distribution references) covering all continents that include a tropical climate: Africa, Asia, Central- and South-America, and Australia with a total of more than 750 different anuran taxa. For our analysis we only included records with species level identifications. Species names were checked and updated if necessary according to Frost [30]. If a taxonomic name

could not be unambiguously assigned to a valid species, i.e. due to cryptic species complexes, the record was not included. This resulted in a data set with 672 species.

For each valid species from the occurrence data set, its life-history and ecological traits (hereafter referred to as traits) were gathered using published literature, suitable data bases reviewed by specialists, and further web resources (see Additional file 2: anuran traits references). Additionally we included our own unpublished data, collected either in the field or from museum specimens (Museum für Naturkunde Berlin, e.g. body size, ripe eggs in female ovaries). Traits collected and used in the analysis comprised information on species distribution, morphology, biology, and ecology. We also noted the geographic (i.e. continent) and phylogenetic (family) origin of each species (see Table 1 for details). As we only considered species for our analysis where at least information on body size (either male or female) was available, the data set was reduced further to 619 species.

Data preparation

Some of the collected trait data required processing for subsequent analysis. We used the elevational range calculated as the difference from the maximum to the minimum elevation where a species is known to occur. Regarding body size, we used the maximum body length known per species and sex or, if not available, mean values plus standard deviation. Only if maximum and/or standard deviation were not available, mean or single values were used. We supplemented the data set with sexual dimorphism, calculated as male divided by female body size. Clutch size was only available for a subset of species (345). The available data on clutch sizes were grouped objectively into ten size classes (A: 4–98, B: 100–265, C: 290–549, D: 563–905, E: 979–1652, F: 1900–3320, G: 3607–6701, H: 8357–12940, I: 17000–25000, J: 36100–40000) and species without information on clutch size were subsequently assigned to a class based on body size (see Additional file 3: clutch size classes for more details).

Studies included in our analyses focused on the comparison of anuran distribution among various landscapes. Hence, broader habitat categories were necessary to combine the results within one analysis. Based on all information available we chose three major habitat categories along a human altered degradation gradient: forest, secondary growth, and non-forest. The habitat category “forest” comprises primary forests, primary forest fragments, and selectively logged or exploited areas; “secondary growth” subsumes secondary forests, edges of primary forests, abandoned plantations (>5 years) and agricultural habitats with remaining forests (e.g. shaded coffee plantations); non-forests comprise simple structured plantations (single strata), pasture or inhabited areas such as villages.

Table 1 Life-history and ecological traits used in the study

Trait	Definition	Scale	Unit/level
Range size ^a	Natural area of occurrence	Ratio	km ²
Elevation	Min. and max. elevation in the entire area of occurrence	Ratio	m asl
SVL male/female	Body length, measured as snout vent length	Ratio	mm
Dimorphism	Calculated as male divided by female body size	Ratio	Proportion
Clutch size	Maximal number of total eggs deposited or maximal number of ripe eggs in the uteri of dissected females	Ratio	#
Clutch size class	Clutch sizes assigned to size classes	Ordinal	Ten size classes, see “Methods” for details
Reproduction	Development	Nominal	Direct, indirect
Adult habitat	Habitat where adults are usually encountered, perch height	Nominal	Aquatic, semi-aquatic, fossorial, litter (<1 m), semi-arboreal (1–3 m), arboreal (>3 m)
Larval habitat	Habitat where the larvae develop	Nominal	None (direct development), terrestrial, semi-aquatic, lentic, lentic and lotic, lotic, phytotelmata (plant associated water bodies, e.g. tree holes, bromeliad tank), skin ^b
Egg deposition	Habitat where the eggs are deposited	Nominal	Terrestrial, semi-terrestrial, aquatic, arboreal, skin ^b
Family	Taxonomic origin, affiliation to family	Nominal	Anuran families according to Frost [30]
Region of origin	Broad geographic region (i.e. continent)	Nominal	

Given is the trait, its definition, the scale of measurement, and the unit (ratio) or levels (nominal, ordinal) of the respective trait


^a Range size according to the IUCN Red List [29] or, if not available, for West African species to the calculated environmental niche model [70]

^b Carried in or on adult male or female

Categorization was realized in accordance with comparative studies [4, 31, 32]. However, in consideration of the modified forest types examined in our data set, slight adaptations and a reduction of categories were necessary. We only took species into account which had information on the presence and absences in these major habitat categories. If a species was detected in several studies, its single occurrence per habitat category (although absent in other studies) was crucial to assign the species to that habitat type. Combining this reduced data set with the available trait data, the final data set amounted to 243 different species with only a few gaps for some traits. As multivariate statistics often require complete data sets, missing values in the trait data set were replaced by dummy variables. This prevents a high loss of information by excluding a trait or a species. For traits with a ratio scale we used the mean, and for traits with a nominal scale the level which occurred most often (compare Table 1). Numbers of required dummy variables in the final data set: range size = 2 (mean = 1,795,153 km²), elevational range: 35 (1217.6 m), snout-vent length (SVL) males: 4 (10.5 mm), SVL females: 21 (18 mm), reproductive mode: 1 (most frequent: indirect development), adult habitat: 2 (litter); larval habitat: 11 (lentic), egg deposition site: 34 (aquatic). All analyses were conducted with the completed data set (see Additional file 4).

Based on the species occurrences in the three major habitat categories, a forest dependency index (FDI) with four levels was established (Fig. 1): dependent species solely detected in forests (D), slightly dependent species occurring in forests and habitats with secondary growth

(SD), forest independent species occurring not in primary forests, i.e. only in habitats with secondary growth and/or non-forested habitats (I), and species with no response occurring in all three habitat categories or forest and non-forest habitats (NR).



Statistical analysis

The distribution and trait data (ratio scale) were non-normal distributed (Shapiro–Wilk test, R package ‘stats’). We thus applied the non-parametric Kruskal–Wallis test and subsequent pairwise Wilcoxon tests with false discovery rate (*fdr*) correction for parameter comparison among species with different forest dependency indices (R package ‘stats’). To filter for species traits explaining the presence or absence of a species in differently degraded habitats and thus their assignment to a particular FDI we performed a Random Forest (RF) classification [33] where 1000 classification trees on bootstrap samples of the data were grown (*randomForest*, R package ‘randomForest’ [34]). The number of candidate variables at each node (*mtry*) was the square root of the total number of variables in the analysis (default setting). To correct for different sample sizes in the training data set, *sampsiz*e was adjusted according to the minimum sample size per analysis. RF was performed for the whole data set and four subsets, three comparing forest dependent species (D) with one of the other FDIs and a comparison of the groups NR and I. We incorporated all available information for a species in RF, including species distribution (range size, elevation range, region of origin) and seven traits (see Table 1). As families were evenly distributed among the different FDIs (see Additional file 5), the affiliation to a family was excluded from the analysis. Binomial Generalized Linear Models (GLM) were performed to filter for potential traits explaining the habitat dependency of a species (*glm*, R package ‘stats’). Therefore, species not responding to habitat changes (NR) were defined as ‘0’ and compared to forest dependent species (D) as well as forest independent species (I), both defined as ‘1’. Numerical variables (body size and sexual dimorphism) were scaled from 0 to 1. To avoid multi-collinearity among explaining variables within one model, generalized variance inflation factors (GVIF) were calculated (*vif*, R package ‘car’). Each model contained the covariates: SVL females, sexual dimorphism, clutch size class, larval habitat, adult habitat, reproductive mode, and egg deposition site. After reducing the co-linearity among the explaining variables and eliminating those with a GVIF higher than five [35], the full model only contained: SVL females, sexual dimorphism, clutch size class, and larval habitat. To test for any influence on the forest dependency of species distribution we fitted Generalized Linear Mixed Effect Models (GLMM) with range size and elevational range (both scaled from 0 to 1) as fixed and the region of origin as random factor (*glme*, R package ‘lme4’). Here, a reduction of covariates due to co-linearity was not necessary. Based on the models, we predicted whether a species is either dependent on forest (non-forest) or occurs in all available habitats (≥ 0.5

for forest, D or non-forest, I; < 0.5 for habitat independent species, NR). All statistical analysis were applied using R 3.2.1 [36].

Results

Taxonomy

The 243 anuran species included in the analysis belonged to 26 different families. The most common families were Rhacophoridae and Hylidae, the latter representing 10–30% of the species in all forest dependency indices (FDIs). The families were equally distributed among the different FDIs (see Additional file 5), ruling out any phylogenetic influence in the data.

Species distribution

Range sizes ranged from 6.17 to 12,217,676 km² and varied highly within each FDI (Table 1 for information on gathered traits; see Table 2; Fig. 2 for results). It differed significantly between forest dependent species (D) and species not responding to habitat alteration (NR) as well as between species slightly depending on forests (SD) and NR. All FDIs covered species with limited and wide altitudinal distribution (see Fig. 2). NR species had the broadest distribution and differed significantly from the others (see Table 2). Species in the final data set originated from Africa, Madagascar, America and Asia. The indices NR, I and SD comprised species from all four regions, only D was lacking Malagasy species (Fig. 3). The region of origin did not differ significantly between the FDIs (Pearson’s χ^2 test: $\chi^2 = 5.89$, *df* = 3, *p* = 0.12).

Habitat

Overall, most species preferred litter as well as shrubs and lower tree strata (1–3 m) as adult habitat (Fig. 3). Almost 75% of the species belonging to D and I live in trees (categories semi-arboreal and arboreal); SD and NR species were mostly found on the ground. Aquatic habitats were not inhabited by SD species, while the other FDIs covered all types. The habitat use differed slightly among the FDIs (Pearson’s χ^2 test: $\chi^2 = 27.28$, *df* = 15, *p* = 0.03). Lentic waters constitute 35–60% of the tadpoles’ habitat per FDI (Fig. 3). Lotic waters were of high importance in SD species, but less in other FDIs. All other categories were only sparsely presented, apart from no larval habitat, representing direct developing species. The larval habitat differed significantly between species assigned to different FDIs ($\chi^2 = 45.23$, *df* = 21, *p* = 0.002).

Body size

Maximum body sizes ranged from 10 to 187 mm for males and from 18 to 287 mm for females, respectively, with a high variation for both sexes within each FDI (see

Table 2 Distribution pattern and life history traits

Trait	D n = 33		SD n = 108		NR n = 83		I n = 19	
	Mean ± SD	Range	Mean ± SD	Range	Mean ± SD	Range	Mean ± SD	Range
a								
Range size (km ²) ^a	1,795,153 ± 2,784,023	6.17–12,217,676	1,511,311 ± 2,996,433	14,67–12,217,676	1,086,942 ± 2,069,589	6.17–10,932,823	2,850,720 ± 3,163,102	21.62–11,045,631
Elevational range (m) ^a	1217.60 ± 591.78	1–3100	994.04 ± 582.89	72–3000	1123.0 ± 557.65	1–2500	1446.33 ± 561.84	400–3002
SVL males (mm) ^a	47.13 ± 30.25	10.5–187	45.83 ± 26.96	18.4–146	45.90 ± 31.85	10.5–180	50.34 ± 31.29	17.0–187
SVL females (mm) ^a	56.37 ± 34.04	18–287	63.06 ± 40.50	24–228.9	50.80 ± 26.95	18–185.0	61.99 ± 40.10	18–287.0
Sexual dimorphism ^a	0.86 ± 0.26	0.09–3.19	0.79 ± 0.29	0.09–2.03	0.90 ± 0.31	0.25–3.19	0.83 ± 0.18	0.49–1.99
Clutch size ^b	1609.62 ± 5186.19	4–40,000	1296.36 ± 3701.47	10–17,000	748.99 ± 1158.77	6–5018	2578.56 ± 7423.93	7–40,000
b								
Trait	Kruskal–Wallis test		Pairwise Wilcoxon test (p)					
	χ ²	df	p	D vs SD	D vs NR	D vs I	SD vs. NR	SD vs. I
Range size (km ²) ^a	19.72	3	<0.001	0.29	<0.001	0.32	<0.01	0.50
Elevational range (m) ^a	21.7	3	<0.0001	0.26	<0.001	0.46	<0.001	0.82
SVL males (mm) ^a	2.67	3	0.45	–	–	–	–	–
SVL females (mm) ^a	5.72	3	0.13	–	–	–	–	–
Sexual dimorphism ^a	8.92	3	<0.05	0.05	0.38	0.38	0.11	0.46
Clutch size ^b	6.92	3	0.07	–	–	–	–	–

Given are the respective mean, standard deviation (sd), and range in general, and for each dependency index separately (a) and comparisons of traits between species of different forest dependency indices using the Kruskal–Wallis test and a pairwise Wilcoxon test with fdr correction as posthoc (b); forest dependency index: D = dependent (n = 33), SD = slightly dependent (n = 108), NR = non-responding (n = 83), I = forest independent (n = 19)

^a Incorporate calculated dummy variables (see “Methods”)

^b Only measured values and therewith differing sample sizes: general = 152, D = 22, SD = 52, NR = 66, I = 12; compare Figs. 2 and 3




Fig. 2 Comparison of distribution patterns and life history traits among anuran species with different forest dependency indices. Given is the range size (a), distribution range along the elevation (b), the maximum body length of males (c) and females (d), size dimorphism between sexes (males/females, e) and the maximum clutch size (f); in addition to the available data, dummy variables were calculated (see “Methods”) and added (a–e); forest dependency index: *D* dependent (n = 33), *SD* slightly dependent (n = 108), *NR* non-responding (n = 83), *I* forest independent (n = 19), for clutch size (f) only measured values are shown, differing sample sizes are given in the graph; see Table 2 for statistical comparisons


Table 2; Fig. 2). It did not differ between the FDIs (see Table 2). Sexual dimorphism also did not show large differences between the indices, but the comparison between *D* and *SD* species showed a trend towards *D* hosting species with greater dimorphism. As female and male body size were highly correlated (Spearman Rank Correlation: $\rho = 0.88$, $p < 0.0001$, $n = 243$), we only used female body size and dimorphism in subsequent analysis.

Reproduction

Clutch size varied between 4 and 40,000 eggs and did not differ between the FDIs (see Table 2; Fig. 2). Independent of the FDI, most clutches were in the first two size classes (4–98 and 100–265 eggs). Species belonging to *I* did not have clutches greater than 6700 eggs. The clutch size measured in categories likewise did not differ significantly between the FDIs (Pearson's χ^2 test: $\chi^2 = 34.96$,

$df = 27$, $p = 0.14$). Most species deposited their eggs in aquatic habitats (see Fig. 3). The second most common habitat was terrestrial, followed by arboreal deposition sites. There were no significant differences in egg deposition site between the FDIs ($\chi^2 = 11.52$, $df = 12$, $p = 0.48$). Almost 80% of the investigated species showed a biphasic development with free swimming tadpoles (see Fig. 3); *D* species had the highest proportion of direct developers (>30%). The reproductive mode did not differ significantly between the FDIs ($\chi^2 = 5.89$, $df = 3$, $p = 0.12$).

Classification by RF on the whole data set resulted in an overall error rate of 50.2%, the misclassification per FDI varied between 42.2 and 94.4% (see Table 3). Classification of subsets performed better, with an overall error rate of 20.7% (*D* vs. *NR*), 22.0% (*D* vs. *I*), 30.4% (*NR* vs. *I*), and 40.4% (*D* vs. *I*). Range size was important in



all, sexual dimorphism and elevational range in four, and clutch size category in three models (Table 4).

Generalized linear models (Table 5) revealed larval habitat and clutch size class as important factors explaining the dependency to forests (D vs. NR species) with the development in lotic waters being significant and clutches of class G (3607–6701 eggs) being almost significant. Based on this model, 77% of all species could be correctly assigned to the original FDIs (matches: D: 12, $n = 33$; NR: 77, $n = 83$; sample size from original data). The model for forest independent species (I vs. NR species) revealed likewise larval habitat as being important with development in lotic waters being significant. The model assigned 82% of all the species correctly to the FDI derived from field observation (I: 4, $n = 19$; NR: 81,

$n = 83$). Generalized Linear Mixed Models (Table 5) fitted with species distribution revealed elevational range as being important factors when comparing both, D and NR as well as I and NR species (the latter barely non-significant). The model contrasting D and NR species assigned 78% of the species to correct FDIs (D = 13; NR = 77), based on the model comparing I and NR species 71% were correctly classified compared to the original FDIs (I = 0; NR = 83). Results of the different approaches confirm each other at least partly: RF vs. GLM: forest dependent species (D): classification overlap of 72%, matches: D: 9; NR: 74; forest independent species (I): 75%, I = 5; NR: 71; RF vs. GLMM: D: 84%, D = 16, NR = 81; I: 71%, I = 0, NR = 72; GLM vs. GLMM: D: 77%, D = 5, NR = 84; I: 94%, I = 0; NR = 96.

Table 3 Confusion matrices of Random Forest analysis

	D	I	NR	SD	CE (%)	OE (%)
Complete data set						50.2
D	15	4	9	5	54.5	
I	5	1	7	6	94.4	
NR	12	6	48	17	42.2	
SD	9	12	29	57	47.2	
Subset D vs. SD						22.0
D	17	–	–	16	48.5	
SD	15	–	–	93	12.0	
Subset D vs. NR						20.7
D	21	–	12	–	36.4	
NR	12	–	71	–	14.5	
Subset D vs. I						40.4
D	23	10	–	–	30.3	
I	11	8	–	–	57.9	
Subset I vs. NR						30.4
I	–	9	10	–	52.6	
NR	–	21	62	–	25.3	

Confusion matrices with per class error (CE) rate and overall error (OE) rate per Random Forest analysis (complete data set and different subsets); analysis were performed with *ntree* = 1000, *mtry* = 3 and *samplesize* adjusted to the smallest sample size (R package 'randomForest'); forest dependency index: *D* dependent (*n* = 33), *SD* slightly dependent (*n* = 108), *NR* non-responding (*n* = 83), *I* forest independent (*n* = 19)

Table 4 Importance of each variable in Random Forest analysis

Variable	Complete	D vs. SD	D vs. NR	D vs. I	I vs. NR
SVL females	6.73	3.41	3.42	2.15	2.08
Adult habitat	4.70	3.23	3.64	0.91	2.37
Sexual dimorphism	9.02	4.00	2.98	2.63	2.59
Clutch size category	6.15	3.49	3.39	4.20	2.63
Egg deposition site	1.13	1.04	0.46	0.50	0.48
Reproductive mode	0.14	0.47	0.07	0.10	0.14
Larval habitat	3.46	2.63	2.10	0.89	1.12
Region	4.54	4.95	2.62	2.11	1.58
Range size	12.52	5.91	7.52	3.27	3.06
Elevational range	8.59	3.37	6.81	2.23	2.87

Importance of each variable per Random Forest analysis (complete data set and different subsets); the four most important variables contributing to the classification are in *italics*; analysis were performed with *ntree* = 1000, *mtry* = 3 and *samplesize* adjusted based on the smallest sample size for each analysis respectively (R package 'randomForest'); forest dependency index: *D* dependent (*n* = 33), *SD* slightly dependent (*n* = 108), *NR* non-responding (*n* = 83), *I* forest independent (*n* = 19)

Discussion

Geographic range size has been identified as a vital factor predicting a species' susceptibility and extinction risk, including birds [11], mammals [10, 16], and amphibians [8, 37]. Species tolerating a wide range of abiotic factors, different habitats [38], or not responding to forest degradation (this study) likewise have the widest distribution. Here, we assign species to one of four levels of

forest dependency, according to their occurrence in habitats with differently strong disturbance. Species belonging to *D* (forest dependent) depend on pristine forests, species assigned to the other categories (*NR*, *I*, or *SD*) can cope with habitat disturbances to different extents. We determined the most important traits explaining the forest dependency of a species using RF classification, GLM, and GLMM techniques. Since range size and extinction risk or habitat breadth might directly depend on each other, making it a single criterion to assess species as critically endangered in the IUCN Red List [29], we excluded it in the GLM filtering for species traits, but analyzed it separately (GLMM). Here, however, only elevational range was important for distinguishing *NR* from *D* and *NR* from *I* species. This is consistent with previous results where a wide altitudinal distribution decreases a species' vulnerability [9, 39, 40], as such species are naturally adapted to varying environmental factors (e.g. vegetation, climate) and hence might also cope better with changes of these factors caused by forest disturbances.

Body size is a central trait, usually correlated with factors such as population size, range size, clutch size or rate of exploitation, all influencing the extinction risk of a species [41–43]. It was thus typically taken into consideration when estimating a species' susceptibility. With increasing body size, studies revealed an increase (amphibians: [39], mammals: [41], birds: [44]), or, as in our data, no change in the extinction risk (amphibians: [28], birds: [40], bats: [45]). These converse results

Table 5 Effects of species traits and distribution on habitat dependency

	Forest dependent species (D vs. NR)				Non-forest species (I vs. NR)			
	Estimate	Std. error	z	p	Estimate	Std. error	z	p
GLM on species traits								
Intercept	0.085	1.47	0.058	0.95	−2.98	1.37	−2.18	0.03
SVL females	−0.08	2.70	−0.03	0.98	0.05	4.59	0.01	0.99
Sexual dimorphism	−3.73	3.11	−1.20	0.23	2.71	2.83	0.96	0.34
Larval habitat								
Lentic/lotic	−17.13	1852.28	−0.009	0.99	−1.13	1.30	−0.87	0.38
Lotic	1.66	0.77	2.16	0.03	3.00	1.34	2.23	0.03
None	0.39	0.64	0.61	0.54	0.26	0.91	0.28	0.78
Phytotelma	18.95	6522.64	0.003	0.99	22.54	10,750	0.002	0.99
Semi-terrestrial	−17.22	4611.48	−0.004	0.99	−16.31	6635	−0.002	0.99
Skin	19.79	6522.64	0.003	0.99	–	–	–	–
Terrestrial	−0.08	2.70	−0.03	0.98	2.50	1.67	1.49	0.14
Clutch size class								
B	0.67	0.67	1.01	0.31	1.14	0.86	1.34	0.18
C	−0.57	1.22	−0.47	0.64	1.73	1.07	1.61	0.11
D	−0.19	1.02	−0.19	0.85	0.20	1.36	0.15	0.88
E	−0.02	0.79	−0.03	0.98	−18.48	2343	−0.01	0.99
F	−0.10	1.10	−0.09	0.93	0.85	1.32	0.64	0.52
G	3.16	1.86	1.70	0.09	−17.69	10,750	−0.002	0.99
H	37.40	6780.54	0.01	0.99				
I	0.59	1.57	0.38	0.71	−17.37	7585	−0.002	0.99
J	−16.74	3995.08	−0.004	0.99	−16.46	7482	−0.002	0.99
GLMM on species distribution								
Fixed effects								
Intercept	1.23	0.68	1.80	0.07	−0.004	0.69	−0.006	0.99
Range size	−1.59	1.13	−1.41	0.16	−1.18	1.38	−0.86	0.39
Elevational range	−4.97	1.45	−3.44	0.0006	−3.03	1.56	−1.85	0.06

Binomial models for forest dependent and non-forest species were conducted and full models (*glm*, R package ‘stats’; *glme*, R packages ‘lme4’) after eliminating multicollinearity (*vif*, R package ‘car’) are presented; Generalized Linear Model (GLM): variables included: SVL females, sexual dimorphism, clutch size class, larval habitat; removed due to co-linearity: adult habitat, reproductive mode, and egg deposition site; Generalized Linear Mixed Model (GLMM): range size and elevation range as fixed and region of continent random factors (no co-linearity among explaining variables); significant effects are in italics; *D* forest dependent species (*n* = 33), *NR* non-responding species (*n* = 83), *I* forest independent species (*n* = 19)

emphasize the complex effects of body size and explain the variation in its influence on the vulnerability of species, differing with study systems [14] but also with the source of extinction risk [46]. According to our results, neither body size nor sexual size dimorphism seem to influence forest dependency.

Although the number of offspring explains the extinction risk in several taxa [12, 13], traits related to reproduction only had minor effects on degradation susceptibility of a frog species in our data set. Species belonging to *I*, however, do not deposit bigger clutches (separating *I* from *NR* species in RF). This could either be related to the larger number of *I* species using flowing, not stagnant, waters as larval habitat and the fact that stream breeders tend to have bigger eggs and thus smaller clutches [47], or to the absence of bigger females,

depositing larger clutches (see Figure in Additional file 3) in *I*. A higher percentage of direct developers among forests dependent species (this study, but see [48, 49]) and an increased extinction risk of ovoviparous anuran species in general ([8], but see [50]) can be explained by the required moist microhabitat for a direct development [51], available in pristine forests, but not necessarily in degraded or fragmented habitats [52, 53].

A species’ microhabitat preferences affect its vulnerability, i.e. the availability of breeding sites, particular soil conditions or vegetation structure can be crucial for the presence of an amphibian species [e.g. 49, 54, 55]. Modified forests are accompanied by an open canopy which facilitates the growth of herbaceous strata and leads to an advantageous humid microclimate for some leaf-litter anurans. This structured understory, including

downed woody debris, has been identified as an important habitat feature for amphibian populations in altered forests [56, 57] and explains the increase of ground dwelling species among degradation tolerant species [25, this study]. Degradation with an accompanying loss of canopy cover generates the most prominent microclimatic shifts in the mid-story, forming the upper strata after disturbances. The resulting decreased humidity, stronger temperature extremes, and increased solar radiation [58–60] have adverse effects on amphibians and explain the high number of semi-arboreal species in our study being forest dependent and the low number being degradation tolerant.

Forest degradation negatively impacts riparian habitats for amphibians by decreasing the amount of woody debris or leaf litter, resulting in less dissolved organic carbon [61] and by a reduction of the canopy cover, leading to higher temperatures and solar radiation [62, 63]. These unfavorable changes explain the higher number of stream breeders among species prone to degradation (this study) and the higher susceptibility of species dependent on lotic breeding sites [54] and riparian species in general [39, 50]. Although forest degradation potentially cause similar changes in lentic habitats, pond breeding amphibians might be less vulnerable or, due to different life-history strategies, even benefit from the consequences: higher temperatures for example increase the developmental rate [64, 65] and higher solar radiation favors the growth of algae [62], the primary food resource for many pond dwelling tadpoles. Compared to species not responding to habitat changes, also a higher number of non-forest species strongly depend on rivers for their tadpole development. These species might be already accustomed to open riparian habitats and thus do not suffer from the prevailing conditions like species occurring in all habitat types.

When contrasting the classification of RF, GLM, and GLMM based on the comparisons D vs. NR, ten species were always wrongly assigned. For example two species, known to occur in strongly degraded habitats [66] and to reproduce in artificial ponds [67] were assigned to D but predicted to belong to NR. Hence the models predicted the species correctly and only the incorporated information from the field was limited and did not cover the occurrences in altered habitats.

Conclusions

Generalist species were identified as the winners in human-dominated landscapes [18, 68], but particular traits facilitating this adaptation were not yet determined. Our pan-tropical approach revealed that the dependency to forested habitats is explained by traits similar to those generally recognized for high species

extinction risk. Indirect developing species exhibiting a big range size, wide elevational range, being independent of streams, and inhabiting the leaf litter are less prone to modifications of their natural habitats. As the effect of a particular trait on the vulnerability of a species might differ among threats [17, 69] and study scales (local vs. global), the generality of our results needs to be treated with caution. However, our findings point to the traits persisting in degraded habitats and thus help to identify future frog communities in our human-dominated world.

Additional files

Additional file 1. Anuran distribution references. References to studies appropriate for the study. Respective data were incorporated in the primary data set on anuran occurrence in different habitat types.

Additional file 2. Anuran traits references. References to journal articles, books and web resources containing information on species traits for the species included in the primary data set on anuran occurrence.

Additional file 3. Clutch size classes. Additional methods describing the objective grouping of clutch sizes.

Additional file 4. Final data set. Given is the species, the traits looked at and the forest dependency index (FDI) derived from the occurrence data. Dummy variables are highlighted in grey. See Table 1 for information on the respective traits.

Additional file 5. Affiliation to different anuran families. Number and relative frequency of species belonging to a particular family per forest dependency index.

Authors' contributions

MH collected data and performed statistical analysis; MH and MOR wrote the text. Both authors read and approved the final manuscript.

Acknowledgements

M. Dahmen provided additional data on species occurrence in Cameroon. M. F. Barej, M. Dahmen, M. Emmrich, H. C. Liedtke, J. Penner, and J. C. Riemann contributed to the trait data set. F. Tillack assisted during literature research and L. Sandberger-Loua gave valuable advices analyzing the data set. A. Channing helped with a thorough language check. All supports are gratefully acknowledged.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

All data generated or analyzed during this study are included in this published article and its additional information files.

Ethics approval and consent to participate

Only Cameroonian frog data were collected for this study. Research permits covered all areas visited and the entire study protocol. They were issued by the Cameroonian Ministry of Scientific Research and Innovation (MINRESI) and the Ministry of Forestry and Wildlife (MINFOP). This study was not approved by any ethics committee as this was not required by Cameroonian or German law. However, all work complied with the guidelines for the use of live amphibians and reptiles in field research compiled by the American Society of Ichthyologists and Herpetologists (ASIH), The Herpetologists' League (HL) and the Society for the Study of Amphibians and Reptiles (SSAR). During amphibian surveys, individuals were identified to species level and immediately released at the same site.

Funding

Fieldwork of MH in Cameroon was supported by scholarships from the Federal State of Berlin (Elsa-Neumann-Stipendium) and the German Academic Exchange Service (DAAD). The publication of this article was funded by the Leibniz Open Access Publishing Fund.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 8 January 2017 Accepted: 20 June 2017

Published online: 28 June 2017

References

- Gibson L, Lee TM, Koh LP, Brook BW, Gardner TA, Barlow J, et al. Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature*. 2011;478(7369):378–81.
- Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, Tyukavina A, et al. High-resolution global maps of 21st-century forest cover change. *Science*. 2013;342(6160):850–3.
- Wright SJ. Tropical forests in a changing environment. *Trends Ecol Evol*. 2005;20(10):553–60.
- Nichols E, Uriarte M, Bunker DE, Favila ME, Slade EM, Vulinec K, et al. Trait-dependent response of dung beetle populations to tropical forest conversion at local and regional scales. *Ecology*. 2013;94(1):180–9.
- Williams NM, Crone EE, T'ai HR, Minckley RL, Packer L, Potts SG. Ecological and life-history traits predict bee species responses to environmental disturbances. *Biol Conserv*. 2010;143(10):2280–91.
- Öckinger E, Schweiger O, Crist TO, Debinski DM, Krauss J, Kuussaari M, et al. Life-history traits predict species responses to habitat area and isolation: a cross-continental synthesis. *Ecol Lett*. 2010;13(8):969–79.
- Bregman TP, Sekercioglu CH, Tobias JA. Global patterns and predictors of bird species responses to forest fragmentation: implications for ecosystem function and conservation. *Biol Conserv*. 2014;169:372–83.
- Sodhi NS, Bickford D, Diesmos AC, Lee TM, Koh LP, Brook BW, et al. Measuring the meltdown: drivers of global amphibian extinction and decline. *PLoS ONE*. 2008;3(2):e1636.
- Rickart EA, Balete DS, Rowe RJ, Heaney LR. Mammals of the northern Philippines: tolerance for habitat disturbance and resistance to invasive species in an endemic insular fauna. *Divers Distrib*. 2011;17(3):530–41.
- Cardillo M, Mace GM, Gittleman JL, Jones KE, Bielby J, Purvis A. The predictability of extinction: biological and external correlates of decline in mammals. *Proc R Soc Lond B Biol Sci*. 2008;275(1641):1441–8.
- Lee TM, Jetz W. Unravelling the structure of species extinction risk for predictive conservation science. *Proc R Soc Lond B Biol Sci*. 2011;278(1710):1329–38.
- Bennett PM, Owens IP. Variation in extinction risk among birds: chance or evolutionary predisposition? *Proc R Soc Lond B Biol Sci*. 1997;264(1380):401–8.
- Siliceo I, Díaz JA. A comparative study of clutch size, range size, and the conservation status of island vs. mainland lacertid lizards. *Biol Conserv*. 2010;143(11):2601–8.
- Henle K, Davies KF, Kleyer M, Margules C, Settele J. Predictors of species sensitivity to fragmentation. *Biodivers Conserv*. 2004;13(1):207–51.
- Davidson AD, Hamilton MJ, Boyer AG, Brown JH, Ceballos G. Multiple ecological pathways to extinction in mammals. *Proc Natl Acad Sci*. 2009;106(26):10702–5.
- Peñaranda DA, Simonetti JA. Predicting and setting conservation priorities for Bolivian mammals based on biological correlates of the risk of decline. *Conserv Biol*. 2015;29(3):834–43.
- Murray KA, Rosauer D, McCallum H, Skerratt LF. Integrating species traits with extrinsic threats: closing the gap between predicting and preventing species declines. *Proc R Soc Lond B Biol Sci*. 2011;278(1711):1515–23.
- McKinney ML, Lockwood JL. Biotic homogenization: a few winners replacing many losers in the next mass extinction. *Trends Ecol Evol*. 1999;14(11):450–3.
- Myers N. Mass extinction and evolution. *Science*. 1997;278(5338):597.
- Stuart SN, Chanson JS, Cox NA, Young BE, Rodrigues ASL, Fischman DL, et al. Status and trends of amphibian declines and extinctions worldwide. *Science*. 2004;306(5702):1783–6.
- Wake DB, Vredenburg VT. Are we in the midst of the sixth mass extinction? A view from the world of amphibians. *Proc Natl Acad Sci*. 2008;105:11466–73.
- Catenazzi A. State of the world's amphibians. *Annu Rev Environ Resour*. 2015;40(1):91–119.
- Ernst R, Linsenmair KE, Rödel M-O. Diversity erosion beyond the species level: dramatic loss of functional diversity after selective logging in two tropical amphibian communities. *Biol Conserv*. 2006;133(2):143–55.
- Riemann JC, Ndriantsoa SH, Raminosoa NR, Rödel M-O, Glos J. The value of forest fragments for maintaining amphibian diversity in Madagascar. *Biol Conserv*. 2015;191:707–15.
- Dixo M, Martins M. Are leaf-litter frogs and lizards affected by edge effects due to forest fragmentation in Brazilian Atlantic forest? *J Trop Ecol*. 2008;24(5):551–4.
- Bielby J, Cooper N, Cunningham AA, Garner TWJ, Purvis A. Predicting susceptibility to future declines in the world's frogs. *Conserv Lett*. 2008;1(2):82–90.
- Laurance WF, Sayer J, Cassman KG. Agricultural expansion and its impacts on tropical nature. *Trends Ecol Evol*. 2014;29(2):107–16.
- Williams SE, Hero J-M. Rainforest frogs of the Australian wet tropics: guild classification and the ecological similarity of declining species. *Proc R Soc Lond B Biol Sci*. 1998;265(1396):597–602.
- Amphibians on the IUCN Red List. <http://www.iucnredlist.org/initiatives/amphibians/analysis/geographic-patterns>. Accessed 3 July 2014.
- Frost DR. Amphibian species of the world: an online reference. Version 6.0. <http://research.amnh.org/herpetology/amphibia/index.html>. Accessed 17 June 2014.
- Nichols E, Larsen T, Spector S, Davis AL, Escobar F, Favila M, et al. Global dung beetle response to tropical forest modification and fragmentation: a quantitative literature review and meta-analysis. *Biol Conserv*. 2007;137(1):1–19.
- Don A, Schumacher J, Freibauer A. Impact of tropical land-use change on soil organic carbon stocks—a meta-analysis. *Glob Change Biol*. 2011;17(4):1658–70.
- Breiman L. Random forests. *Machine learning*. 2001;45(1):5–32.
- Liaw A, Wiener M. Classification and regression by randomForest. *R News*. 2002;2(3):18–22.
- Zuur AE, Ieno EN, Walker NJ, Saveliev AA, Smith GM. Mixed effects models and extensions in ecology with R. New York: Springer; 2009.
- R Core Team. A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2015. <http://www.R-project.org>.
- Cooper N, Bielby J, Thomas GH, Purvis A. Macroecology and extinction risk correlates of frogs. *Glob Ecol Biogeogr*. 2008;17(2):211–21.
- Slatyer RA, Hirst M, Sexton JP. Niche breadth predicts geographical range size: a general ecological pattern. *Ecol Lett*. 2013;16(8):1104–14.
- Lips KR, Reeve JD, Witters LR. Ecological traits predicting amphibian populations declines in Central America. *Conserv Biol*. 2003;17(4):1078–88.
- White RL, Bennett PM. Elevational distribution and extinction risk in birds. *PLoS ONE*. 2015;10(4):e0121849.
- Cardillo M, Mace GM, Jones KE, Bielby J, Bininda-Emonds OR, Sechrest W, et al. Multiple causes of high extinction risk in large mammal species. *Science*. 2005;309(5738):1239–41.
- McKinney ML. Extinction vulnerability and selectivity: combining ecological and paleontological views. *Annu Rev Ecol Syst*. 1997;28:495–516.
- Blüeweiß L, Fox H, Kudzma V, Nakashima D, Peters R, Sams S. Relationships between body size and some life history parameters. *Oecologia*. 1978;37(2):257–72.
- Pavlacky DC Jr, Possingham HP, Goldizen AW. Integrating life history traits and forest structure to evaluate the vulnerability of rainforest birds along gradients of deforestation and fragmentation in eastern Australia. *Biol Conserv*. 2015;188:89–99.
- Jones KE, Purvis A, Gittleman JL. Biological correlates of extinction risk in bats. *Am Nat*. 2003;161(4):601–14.
- Owens IPF, Bennett PM. Ecological basis of extinction risk in birds: habitat loss versus human persecution and introduced predators. *Proc Natl Acad Sci*. 2000;97(22):12144–8.

47. Wells KD. The ecology and behaviour of amphibians. Chicago: University of Chicago Press; 2007.
48. Ernst R, Rödel M-O. Anthropogenically induced changes of predictability in tropical anuran assemblages. *Ecology*. 2005;86(11):3111–8.
49. Hillers A, Veith M, Rödel M-O. Effects of forest fragmentation and habitat degradation on West African leaf-litter frogs. *Conserv Biol*. 2008;22(3):762–72.
50. Lips KR. Decline of a tropical montane amphibian fauna. *Conserv Biol*. 1998;12(1):106–17.
51. Hödl W. Reproductive diversity in Amazonian lowland frogs. In: Hanke W, editor. *Biology and physiology of amphibians*. Stuttgart: Gustav Fischer Verlag; 1990. p. 41–60.
52. Murcia C. Edge effects in fragmented forests: implications for conservation. *Trends Ecol Evol*. 1995;10(2):58–62.
53. Martius C, Höfer H, García MV, Römbke J, Förster B, Hanagarth W. Microclimate in agroforestry systems in central Amazonia: does canopy closure matter to soil organisms? *Agrofor Syst*. 2004;60(3):291–304.
54. Ernst R, Rödel M-O, Arjoon D. On the cutting edge—the anuran fauna of the Mabura Hill Forest Reserve, central Guyana. *Salamandra*. 2005;41(4):179–94.
55. Urbina-Cardona JN, Olivares-Pérez M, Reynoso VH. Herpetofauna diversity and microenvironment correlates across a pasture–edge–interior ecotone in tropical rainforest fragments in the Los Tuxtlas Biosphere Reserve of Veracruz, Mexico. *Biol Conserv*. 2006;132(1):61–75.
56. Spear SF, Crisafulli CM, Storfer A. Genetic structure among coastal tailed frog populations at Mount St. Helens is moderated by post-disturbance management. *Ecol Appl*. 2012;22(3):856–69.
57. Rittenhouse TAG, Harper EB, Rehard LR, Semlitsch RD. The role of microhabitats in the desiccation and survival of anurans in recently harvested oak-hickory forest. *Copeia*. 2008;2008(4):807–14.
58. Chen J, Saunders SC, Crow TR, Naiman RJ, Broszofsky KD, Mroz GD, et al. Microclimate in forest ecosystem and landscape ecology: variations in local climate can be used to monitor and compare the effects of different management regimes. *BioScience*. 1999;49(4):288–97.
59. Szarzynski J, Anhuf D. Micrometeorological conditions and canopy energy exchanges of a neotropical rain forest (Surumoni-Crane Project, Venezuela). *Plant Ecol*. 2001;153(1–2):231–9.
60. Turlon SM, Siegenthaler DT. Immediate impacts of a severe tropical cyclone on the microclimate of a rain-forest canopy in north-east Australia. *J Trop Ecol*. 2004;20(5):583–6.
61. Castelle AJ, Johnson AW. Riparian vegetation effectiveness. Technical Bulletin 799: National Council for Air and Stream Improvement, Inc.; 2000.
62. Kelly DJ, Bothwell ML, Schindler DW. Effects of solar ultraviolet radiation on stream benthic communities: an intersite comparison. *Ecology*. 2003;84(10):2724–40.
63. Johnson SL, Jones JA. Stream temperature responses to forest harvest and debris flows in western Cascades, Oregon. *Can J Fish Aquat Sci*. 2000;57(52):30–9.
64. Álvarez D, Nicieza AG. Effects of temperature and food quality on anuran larval growth and metamorphosis. *Funct Ecol*. 2002;16(5):640–8.
65. Atkinson D. Ectotherm life history responses to developmental temperature. In: Johnston IA, Bennett AF, editors. *Animals and temperature: phenotypic and evolutionary adaptation*. Cambridge: Cambridge University Press; 1996. p. 419.
66. Iskandar D, Mumpuni, Richards S. *Limnonectes grunniens*. The IUCN Red List of Threatened Species. Version 2015.2. <http://www.iucnredlist.org>. Accessed 19 Aug 2015.
67. Shunqing L., Datong Y, Liang F, van Dijk PP, Chanard T, Sengupta S, et al. *Rhacophorus maximus*. The IUCN Red List of Threatened Species. Version 2015.2. <http://www.iucnredlist.org>. Accessed 19 Aug 2015.
68. Devictor V, Julliard R, Jiguet F. Distribution of specialist and generalist species along spatial gradients of habitat disturbance and fragmentation. *Oikos*. 2008;117(4):507–14.
69. González-Suárez M, Gómez A, Revilla E. Which intrinsic traits predict vulnerability to extinction depends on the actual threatening processes. *Ecosphere*. 2013;4(6):76.
70. Penner J. Macroecology of West African amphibians. Ph. D. thesis. Berlin: Humboldt University Berlin; 2014.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit



Additional file 1. Anuran distribution references

References to studies appropriate for the study. Respective data were incorporated in the primary data set on anuran occurrence in different habitat types.

- Adum, G. B., M. P. Eichhorn, W. Oduro, C. Ofori-Boateng & M.-O. Rödel (2013): Two-stage recovery of amphibian assemblages following selective logging of tropical forests. *Conservation Biology* 27: 354-363.
- Akani, G. C., E. Politano & L. Luiselli (2004): Amphibians recorded in forest swamp areas of the River Niger Delta (southeastern Nigeria), and the effects of habitat alteration from oil industry development on species richness and diversity. *Applied Herpetology* 2: 1-22.
- Bell, K. E. & M. A. Donnelly (2006): Influence of forest fragmentation on community structure of frogs and lizards in Northeastern Costa Rica. *Conservation Biology* 20: 1750-1760.
- Bickford, D., T. How Ng, L. Qie, E. P. Kudavidanage & C. J. A. Bradshaw (2010): Forest fragment and breeding habitat characteristics explain frog diversity and abundance in Singapore. *Biotropica* 42:119-125.
- Cabrera-Guzmán, E. & V. H. Reynoso (2012): Amphibian and reptile communities of rainforest fragments: minimum patch size to support high richness and abundance. *Biodiversity and Conservation* 21: 3243-3265.
- Cáceres-Andrade, S. P. & J. N. Urbina-Cardona (2009): Esamblajes de anuros de sistemas productivos y bosques en el piedemonte Llanero, Departamento del Meta, Colombia. *Caldasia* 31: 175-194.
- Campos, V. A., F. H. Oda, L. Juen, A. Barth & A. Dartora (2013): Composition and species richness of anuran amphibians in three different habitat in an agrosystem in Central Brazilian Cerrado. *Biota Neotropica* 13: 124-132.
- Cortés, A. M., M. P. Ramírez-Pinilla, H. A. Suárez & E. Tovar (2008): Edge effects on richness, abundance and diversity of frogs in Andean cloud forest fragments. *South American Journal of Herpetology* 3: 213-222.
- da Silva, F. R. & d. C. Rossa-Feres (2011): Influence of terrestrial habitat isolation on the diversity and temporal distribution of anurans in an agricultural landscape. *Journal of Tropical Ecology* 27: 327-331.
- de Lima, M. G. & C. Gascon (1999): The conservation value of linear forest remnants in central Amazonia. *Biological Conservation* 91: 241-247.
- Dixo, M. & M. Martins (2008): Are leaf-litter frogs and lizards affected by edge effects due to forest fragmentation in Brazilian Atlantic forest? *Journal of Tropical Ecology* 24: 551-554.
- Ernst, R., K. E. Linsenmair & M.-O. Rödel (2006): Diversity erosion beyond the species level: dramatic loss of functional diversity after selective logging in two tropical amphibian communities. *Biological Conservation* 133: 143-155.
- Ernst, R. & M.-O. Rödel (2005): Anthropogenically induced changes of predictability in tropical anuran assemblages. *Ecology* 86: 3111-3118.
- Ernst, R. & M.-O. Rödel (2008): Patterns of community composition in two tropical tree frog assemblages: separating spatial structure and environmental effects in disturbed and undisturbed forests. *Journal of Tropical Ecology* 24: 111-120.
- Faria, D., M. L. Barradas Paciencia, M. Dixo, R. R. Laps & J. Baumgarten (2007): Ferns, frogs, lizards, birds and bats in forest fragments and shade cacao plantations in two contrasting landscapes in the Atlantic forest, Brazil. *Biodiversity and Conservation* 16: 2335-2357.
- Ficetola, G. F., D. Furlani, G. Colombo & F. de Bernardi (2008): Assessing the value of secondary forest for amphibians: *Eleutherodactylus* frogs in a gradient of forest alteration. *Biodiversity and Conservation* 17: 2185-2195.
- Fredericksen, N. J. & T. S. Fredericksen (2002): Terrestrial wildlife responses to logging and fire in a Bolivian tropical humid forest. *Biodiversity and Conservation* 11: 27-38.

- Furlani, D., G. F. Ficetola, G. Colombo, M. Ugurlucan & F. de Bernardi. (2009): Deforestation and the structure of frog communities in the Humedale Terraba-Sierpe, Costa Rica. *Zoological Science* 26: 197-202.
- Gardner, T. A., M. A. Ribeiro-Júnior, J. Barlow, T. Cristina, S. Ávila-Pires, M. S. Hoogmoed & C. A. Peres (2007): The value of primary, secondary, and plantation forests for a neotropical herpetofauna. *Conservation Biology* 21: 775-787.
- Gascon, C. (1993): Breeding-habitat use by five Amazonian frogs at forest edge. *Biodiversity and Conservation* 2: 438-444.
- Germano, J. M., J. M. Sander, R. W. Henderson & R. Powell (2003): Herpetofaunal communities in Grenada: a comparison of altered sites, with an annotated checklist of Grenadian amphibians and reptiles. *Caribbean Journal of Science* 39: 68-76.
- Gillespie, G., S. Howard, D. Lockle, M. Scroggie & Boedl (2005): Herpetofaunal richness and community structure of offshore islands of Sulawesi, Indonesia. *Biotropica* 37: 279-290.
- Gillespie, G. R., E. Ahmad, B. Elahan, A. Evans, M. Ancrenaz, B. Goossens & M. P. Scroggie (2012): Conservation of amphibians in Borneo: relative value of secondary tropical forest and non-forest habitats. *Biological Conservation* 152: 136-144.
- Heinen, J. T. (1992): Comparison of the leaf litter herpetofauna in abandoned cacao plantations and primary rain forest in Costa Rica: some implications for faunal restoration. *Biotropica* 24: 431-439.
- Hilje, B. & T. Mitchell Aide (2012): Recovery of amphibian species richness and composition in a consequence of secondary forests, northeastern Costa Rica. *Biological Conservation* 146: 170-176.
- Hillers, A., M. Veith & M.-O. Rödel (2008): Effects of forest fragmentation and habitat degradation on West African leaf-litter frogs. *Conservation Biology* 22: 762-772.
- Hinde, R., G. Corti, E. Fanning & R. K. B. Jenkins (2001): Anurans in the Kilombero Valley, Tanzania: comparison between miombo woodland, evergreen forest and teak plantations. *African Journal of Herpetology* 50: 35-39.
- Inger, R. F. & R. K. Colwell (1977): Organization of contiguous communities of amphibians and reptiles in Thailand. *Ecological Monographs* 47: 229-253.
- Isaacs Cubides, P. J. & J. N. Urbina Cardona (2011): Anthropogenic disturbance and edge effects on anuran assemblages inhabiting cloud forest fragments in Colombia. *Brazilian Journal of Nature Conservation* 9: 39-46.
- Katwate, U., D. Apte & R. Raut (2013): Diversity and distribution of anurans in Phansad Wildlife Sanctuary (PWS), northern Western Ghats of India. *Journal of Threatened Taxa* 5: 3589-3602.
- Krishna, S. N., S. B. Krishna & K. K. Vijayalaxmi (2005): Variation in anuran abundance along the streams of the Western Ghats, India. *Herpetological Journal* 15: 167-172.
- Krishnamurthy, S. V. (2003): Amphibian assemblages in undisturbed and disturbed areas of Kudremukh National Park, central Western Ghats, India. *Environmental Conservation* 30: 274-282.
- Kudavidangage, E. P., T. C. Wanger, C. de Alwis, S. Sanjeewa & S. W. Kotagama (2011): Amphibian and butterfly diversity across a tropical land-use gradient in Sri Lanka; implications for conservation decision making. *Animal Conservation* 15: 253-265.
- Lieberman, S. S. (1986): Ecology of the leaf litter herpetofauna of a neotropical rain forest: La Selva, Costa Rica. *Acta Zoologica Mexicana* 15: 1-74.
- Malzona, P. K. & M. Veith (2012): Amphibian community along elevational and habitat disturbance gradients in the Taita Hills, Kenya. *Herpetotropicalos* 7: 7-16.
- Maritz, B. (2007): The distribution and abundance of herpetofauna on a quaternary aeolian dune deposit: implications for strip mining. University of the Witwatersrand, Johannesburg.
- Murali, R. & T. R. S. Raman (2012): Streamside amphibian communities in plantations and a rainforest fragment in the Anamalai hills, India. *Journal of Threatened Taxa* 4: 2849-2856.
- Murrieta-Galindo, R., A. González-Romero, F. López-Barrera & G. Parra-Olea (2013): Coffee agrosystems: an important refuge for amphibians in central Veracruz, Mexico. *Agroforestry Systems* 87: 1-13.
- Nath, N., S. Sutradhar, A. K. Mani, V. Vijayan, K. Kumar, B. L. Narayana, B. Naresh, G. Baburao, S. Dharwadkar, G. SKrishnan, B. Vinoth, R. Maniraj, D. M. Reddy, D. Adi mallaiah & K. Swamy (2012):

- Herpetofaunal assemblage with special emphasis on community structure and spatiality in amphibians of Cauvery delta region, Tamil Nadu. *Asian Journal of Conservation Biology* 1: 78-85.
- Ofori-Boateng, C., W. Oduro, A. Hillers, K. Norris, S. K. Oppong, G. B. Adum & M.-O. Rödel (2012): Differences in the effects of selective logging on amphibian assemblages in three West African forest types. *Biotropica* 45: 94-101.
- Parris, K. M. & D. B. Lindenmayer (2004): Evidence that creation of a *Pinus radiata* plantation in south-eastern Australia has reduced habitat for frogs. *Acta Oecologica* 25: 93-101.
- Pawar, S. (1999): Effect of habitat alteration on herpetofaunal assemblages of evergreen forest in Mizoram, North-East India. Saurashtra University, Rajkot.
- Pawar, S., G. S. Rawat & B. C. Choudhury (2004): Recovery of frog and lizard communities following primary habitat alteration in Mizoram, Northeast India. *BMC Ecology* 4: 10.
- Pineda, E. & G. Halffter (2004): Species diversity and habitat fragmentation: frogs in a tropical montane landscape in Mexico. *Biological Conservation* 117: 499-508.
- Pineda, E., C. Moreno, F. Escobar & G. Halffter (2005): Frog, bat, and dung beetle diversity in the cloud forest and coffee agroecosystems of Veracruz, Mexico. *Conservation Biology* 19: 400-410.
- Reichert, M. (2004): A comparison of anuran species richness between primary and secondary forest in São Francisco do Pará. Independent Study Project (ISP) Collection 524.
- Ribeiro-Júnior, M. A., T. A. Gardner & T. C. S. Ávila-Pires (2008): Evaluating the effectiveness of herpetofaunal sampling techniques across a gradient of habitat change in a tropical forest landscape. *Journal of Herpetology* 42: 733-749.
- Russel, C. & C. T. Downs (2012): Effect of land use on anuran species composition in north-eastern KwaZulu-Natal, South Africa. *Applied Geography* 35: 247-256.
- Santos-Barrera, G. & J. N. Urbina-Cardona (2011): The role of the matrix-edge dynamics of amphibian conservation in tropical montane fragmented landscapes. *Revista mexicana de biodiversidad* 82: 679-687.
- Schlaepfer, M. A. & T. A. Gavin (2001): Edge effects on lizards and frogs in tropical forest fragments. *Conservation Biology* 15: 1079-1090.
- Tocher, M. D., C. Gascon & J. Meyer (2001): Community composition and breeding success of Amazonian frogs in continuous forest and matrix habitat aquatic sites. pp. 235-247 in R. O. J. Bierregaard, C. Gascon, T. E. Lovejoy & R. C. G. Mesquita (eds.). *Lessons from Amazonia — the ecology and conservation of a fragmented forest*. Yale University Press, New Haven & London.
- Tocher, M. D., C. Gascon & B. L. Zimmerman (1997): Fragmentation effects on a Central Amazonian frog community: a ten year study. pp 124-137 in W. F. Laurance & R. O. J. Bierregaard (eds.). *Tropical forest remnants - ecology, management, and conservation of fragmented landscapes*. The University of Chicago Press, Chicago & London.
- Toral C., E., P. Feinsinger & M. L. Crump (2002): Frogs and a cloud-forest edge in Ecuador. *Conservation Biology* 16: 735-744.
- Urbina-Cardona, J. N. & M. C. Londoño-M. (2003): Distribución de la comunidad de herpetofauna asociada a cuatro áreas con diferente grado de perturbación en la Isla Gorgona, Pacífico colombiano. *Academia Colombiana de Ciencias* 27: 105-113.
- Urbina-Cardona, J. N., M. Olivares-Pérez & V. H. Reynoso (2006): Herpetofauna diversity and microenvironment correlates across a pasture–edge–interior ecotone in tropical rainforest fragments in the Los Tuxtlas Biosphere Reserve of Veracruz, Mexico. *Biological Conservation* 132: 61-75.
- Vallan, D. (2000): Influence of forest fragmentation on amphibian diversity in the nature reserve of Ambohitantely, highland Madagascar. *Biological Conservation* 96: 31-43.
- Vallan, D. (2002): Effects of anthropogenic environmental changes on amphibian diversity in the rain forests of eastern Madagascar. *Journal of Tropical Ecology* 18: 725-742.
- Vallan, D., F. Andreone, V. H. Raherisoa & R. Dolch (2004): Does selective wood exploitation affect amphibian diversity? The case of An'Ala, a tropical rainforest in eastern Madagascar. *Oryx* 38: 410-417.
- Vonesh, J. R. (2001): Patterns of richness and abundance in a tropical African leaf-litter herpetofauna. *Biotropica* 33: 502-510.

- Wanger, T. C., D. T. Iskandar, I. Motzke, B. W. Brook, N. S. Sodhi, Y. Clough & T. Tschardtke (2011): Effects of landuse change on community composition of tropical amphibians and reptiles in Sulawesi, Indonesia. *Conservation Biology* 24: 795-802.
- Watling, J. I. & M. A. Donnelly (2007): Multivariate correlates of extinction proneness in a naturally fragmented landscape. *Diversity and Distribution* 13: 372-378.

Additional file 2. Anuran traits references.

References to journal articles, books and web resources containing information on species traits for the species included in the primary data set on anuran occurrence.

Journal articles and Books:

- Abravaya, J. P. & J. F. Jackson (1978): Reproduction in *Macrogenioglottus alipioi* Carvalho (Anura, Leptodactylidae). Serial Publications of the Natural History Museum of Los Angeles County 298: 1-9.
- Adler, K. (1965): Three new frogs of the genus *Hyla* from the Sierra Madre del Sur of Mexico. Occasional Papers of the Museum of Zoology University of Michigan 642: 1-18.
- Adum, B. A., C. Ofori-Boateng, W. Oduro & M.-O. Rödel (2011): Re-discovery of the giant West African squeaker, *Arthroleptis krokosua* Ernst, Agyei & Rödel, 2008 (Amphibia: Anura: Arthroleptidae) in two forests of southwestern Ghana with observations on the species' variability and habitat preferences. Zootaxa 2744: 34-38.
- Alcala, A. C. & W. C. Brown (1998): Philippine amphibians: an illustrated field guide. Bookmark, Makati City, Philippines.
- Amiet, J.-L. (1977): Les *Astylosternus* du Cameroun (Amphibia anura, Astylosterninae). Annales de la Faculté des Science de Yaoundé 23-24: 99-237.
- Amiet, J.-L. (1991): Images d'amphibiens camerounais. III. Le comportement de garde des oeufs Alytes 9: 15-22.
- Amiet, J.-L. (2012). Les rainettes du Cameroun. J.-L. Amiet and La Naf des Livre, Sain-Nazaire.
- Anahi Güizado-Rodriguez, M., U. Omar Garcia-Vazquez & J. L. Aguilar-Lopez (2010): Thermoregulation of *Craugastor berkenbuschii* (Peters, 1870). Herpetological bulletin 112: 4-6.
- Andreone, F. & R. A. Nussbaum (2006): A revision of *Mantidactylus microtis* and *M. microtympanum*, and a comparison with other large Madagascan stream frogs (Anura: Mantellidae: Mantellinae). Zootaxa 1105: 49-68.
- Andreone, F., G. M. Rosa & A. P. Raselimanana (2014). Les amphibiens des zones arides de l'ouest et du sud de Madagascar. Association Vahatra, Antananarivo.
- Angel, F. (1940): Description de trois amphibiens nouveaux du Cameroun, matériaux de la mission *P. lepesme*, *R. paulian* et *A. villiers* (2e note). Bulletin du Muséum national d'histoire naturelle Paris 12: 238-243.
- Anstis, M. (2013): Tadpoles and frogs of Australia. New Holland, Sindney.
- Ao, M., S. Bordoloi, A. Ohler & S. Grosjean (2006): *Rana khare* (Kiyasetuo & Khare, 1986): present distribution, redescription of holotype and morphology of adults and tadpoles. Alytes 24: 22-39.
- Ariyasiri, K., G. Bowatte, U. Menike, S. Meegaskumbura & M. Meegaskumbura (2011): Predator-induced plasticity in tadpoles of *Polypedates cruciger* (Anura: Rhacophoridae). Amphibian and Reptile Conservation 5: 14-21.
- Artenga, A., L. Bustamante & J. M. Guayasamin (2014): The amphibians and reptiles of Mindo — life in the cloudforest. Universidad Tecnológica Indoamérica, Quito.
- Ávila, R. W., A. Pansonato & C. Strüssmann (2010): A new species of the *Rhinella margaritifera* group (Anura: Bufonidae) from Brazilian Pantanal. Zootaxa 2339: 57-68.
- Bahir, M. M., M. Meegaskumbura, K. Manamendra-Arachchi, C. J. Schneider & R. Pethiyagoda (2005): Reproduction and terrestrial direct development in Sri Lankan shrub frogs (Ranidae: Rhacophorinae: *Philautus*). The Raffles Bulletin of Zoology 12: 339-350.
- Barej, M. F., M.-O. Rödel, N. L. Gonwouo, O. S. G. Pauwels, W. Böhme, & A. Schmitz (2010): Review of the genus *Petropedetetes* Reichenow, 1874 in Central Africa with the description of three new species (Amphibia: Anura: Petropedetidae). Zootaxa 2340: 1-49.

- Barej, M. F., A. Schmitz, M. Menegon, A. Hillers, H. Hinkel, W. Böhme & M.-O. Rödel (2011): Dusted off — the African *Amietophrynus superciliaris*-species complex of giant toads. *Zootaxa* 2772: 1-32.
- Barreto, L. & G. V. Andrade (1995): Aspects of the reproductive biology of *Physalaemus cuvieri* (Anura: Leptodactylidae) in northeastern Brazil. *Amphibia-Reptilia* 16: 67-76.
- Barrio-Amorós, C. L., J. M. Guayasamin & S. B. Hedges (2012): A new minute Andean *Pristimantis* (Anura: Strabomantidae) from Venezuela. *Phyllomedusa* 11: 83-93.
- Barrio-Amorós, C. L. & H. Kaiser (2008): Distribution of *Strabomantis biporcatus* (Terrarana: Strabomantidae) in northern Venezuela, with comments on its phenotypic variation. *Salamandra* 44: 248-254.
- Barros, F. B., L. Vicente & H. M. Pereira (2010): Amphibians of the Riozinho do Anfrísio Extractive Reserve, Pará. Rapid Color Guide.
- Bennett, D., K. Hampson, K. Sanders & M. Anderson (1998): Frogs of Coorg, Karnataka, India. Aberdeen University Western Ghats Project 1998.
- Berry, P. Y. & J. R. Hendrickson (1963): *Leptobrachium nigrops*, a new pelobatid frog from the Malay Peninsula, with remarks on the genus *Leptobrachium* in southeastern Asia. *Copeia* 1963: 643-648.
- Biju, S. D. (2003): Reproductive mode in the shrub frog *Philautus glandulosus* (Jerdon, 1853) (Anura: Rhacophoridae). *Current Science* 84: 283-284.
- Biju, S. D. & F. Bossuyt (2009): Systematics and phylogeny of *Philautus* Gistel, 1848 (Anura, Rhacophoridae) in the Western Ghats of India, with descriptions of 12 new species. *Zoological Journal of the Linnean Society* 155: 374-444.
- Biju, S. D., I. van Bocxlaer, V. B. Giri, S. P. Loader & F. Bossuyt (2009): Two new endemic genera and a new species of toad (Anura: Bufonidae) from the Western Ghats of India. *BMC research notes* 2: 241.
- Biju, S. D., I. van Bocxlaer, S. Mahony, K. P. Dinesh, C. Radhahrishnan, A. Zachariah, V. Giri & F. Bossuyt (2011): A taxonomic review of the night frog genus *Nyctibatrachus* Boulenger, 1882 in the Western Ghats, India (Anura: Nyctibatrachidae) with description of twelve new species. *Zootaxa* 3029: 1-96.
- Bogert, C. M. (1969) The eggs and hatchlings of the Mexican leptodactylid frog *Eleutherodactylus decoratus* Taylor. *American Museum novitates* 2376: 1-9.
- Bolívar-G, W., A. Giraldo & J. Mendez (2011): Amphibia, Anura, Strabomantidae, *Pristimantis palmeri* Boulenger, 1912: distribution extension for the Central Cordillera, Colombia. *CheckList* 7: 9-10.
- Bopage, M. M., K. Wewalwala, M. Krvavac, O. Jovanovic, G. Safarek & V. Pushpamal (2011): Species diversity and threat status of amphibians in the Kanneliya Forest, lowland Sri Lanka. *Salamandra* 47: 173-177.
- Boulenger, G. A. (1896): II. — Descriptions of new Reptiles and Batrachians from Colombia. *Journal of Natural History* 17: 16-21.
- Boulenger, G. A. (1898): Fourth report on additions to the batrachian collection in the Natural-History Museum Proceedings of the Zoological Society of London 66: 473-482.
- Boulenger, G. A. (1920): A monograph of the South Asian, Papuan, Melanesian and Australian frogs of the genus *Rana*. *Zoological Survey of India, Calcutta*.
- Bourne, G. R. (1997): Reproductive behaviour of terrestrial breeding frogs *Eleutherodactylus johnstonei* in Guyana. *Journal of Herpetology* 31: 221-229.
- Brauer, K. (1991): Kröten. *Urania*, Leipzig. Buchacher, C. (1993): Field studies on the Small Surinam toad *Pipa arrabali* near Manaus, Brazil. *Institut für Zoologie der Universität Wien, Wien*.
- Caldwell, J. P. (1991). A new species of toad in the genus *Bufo* from Pará, Brazil, with an unusual breeding site. *Papéis Avulsos de Zoologia* 37: 389-400.
- Caldwell, J. P. & P. T. Lopez (1989): Foam-generating behavior in tadpoles of *Leptodactylus mystaceus*. *Copeia* 1989: 498-502.
- Campbell, J. A., J. C. Blancas-Hernández & E. N. Smith (2009): A new species of stream-breeding treefrog of the genus *Charadrahyla* (Hylidae) from the Sierra Madre del Sur of Guerrero, Mexico. *Copeia* 2009: 287-295.
- Canedo, C., M. Dixo & J. P. Pombal Jr. (2004): A new species of *Chiasmocleis* Méhely, 1904 (Anura, Microhylidae) from the Atlantic rainforest of Bahia, Brazil. *Herpetologica* 60: 495-501.

- Cannatella, D. C. (1986): A new species of *Osornophryne* (Anura: Bufonidae) from the Andes of Ecuador. *Copeia* 1986. Caramaschi, U. & C. A. G. da Cruz (1997): Redescription of *Chiasmocleis albopunctata* (Boettger) and description of a new species of *Chiasmocleis* (Anura: Microhylidae). *Herpetologica* 53 : 259-268.
- Caramaschi, U. & J. P. Pombal Jr. (2006): A new species of *Rhinella Fitzinger*, 1826 from the Atlantic Rain Forest, Eastern Brazil (Amphibia, Anura, Bufonidae). *Papéis Avulsos de Zoologia* 46: 251-259.
- Cardozo, D. & F. Lobo (2009): *Pseudopaludicola mirandae* Mercadal de Barrio and Barrio, 1994 (Anura, Leiuperidae) is a junior synonym of *Pseudopaludicola boliviana* Parker, 1927. *Journal of Herpetology* 43: 685-687.
- Carruthers, V. (2001): *Frogs and frogging in southern Africa*. Struik Publishers, Cape Town.
- Chanda, S. K. (2002): *Handbook. Indian Amphibians*. Zoological Survey of India, Calcutta.
- Chandramouli, S. R. & S. R. Ganesh (2010): Herpetofauna of Southern Western Ghats, India — reinvestigated after decades. *Taprobanica* 2: 72-85.
- Chandramouli, S. R., S. R. Ganesh & N. Baskaran (2011a). On recent sightings of a little-known south Indian toad, *Duttaphrynus hololius* (Günther, 1876) with notes on its morphological characterization and ecology. *Herpetology Notes* 4: 271-274.
- Chandramouli, S. R., S. Harikrishnan S. & K. Vasudevan (2011b): Little known endemic frogs of the Andaman Islands. *FrogLog* 98: 16-17.
- Chandramouli, S. R. & A. Kalaimani (2014): Description of the larvae of Günther's toad *Duttaphrynus hololius* (Günther, 1876) (Anura: Bufonidae) with notes on development and oral ultra-structure. *Alytes* 31: 3-12.
- Channing, A. (2001): *Amphibians of Central and Southern Africa*. Cornell University Press, Ithaca.
- Channing, A. & K. M. Howell (2006): *Amphibians of East Africa*. Cornell University Press, Ithaca.
- Channing, A., M.-O. Rödel & J. Channing (2012): Tadpoles of Africa. The biology and identification of all known tadpoles in sub-Saharan Africa. Edition Chimaira, Frankfurt/M.
- Chávez, G., C. H. Cosmópolis & L. Luján (2013): Annotated checklist and ecological notes of anurans from the Southern region of Yanachaga Chemillen National Park, Central Andes of Peru. *Herpetotropics* 8: 23-18.
- Chochran, D. M. (1927): New reptiles and batrachians collected by Dr. Hugh M. Smithn Siam. *Proceedings of the Biological Society of Washington* 40: 179-192.
- Cochran, D. M. & C. J. Coin (1970): *Frogs of Colombia*. Smithsonian Institution Press, Washington.
- Coe, M. J. (1967): Co-operation of three males in nest construction by *Chiromantis rufescens* Günther (Amphibia: Rhacophoridae). *Nature* 214: 112-113.
- Crump, M. L. (1974): Reproductive strategies in a tropical anuran community. University of Kansas Museum of Natural History Miscellaneous Publications 61: 1-68.
- da Cruz, C. A. G., U. Caramaschi & E. Izecksohn (1997): The genus *Chiasmocleis* Méhely, 1904 (Anura, Microhylidae) in the Atlantic Rain Forest of Brazil, with description of three new species. *Alytes* 15: 49-71.
- da Cruz, C. A. G. & M. F. Napoli (2010): A new species of smooth horned frog, genus *Proceratophrys* Miranda-Ribeiro (Amphibia: Anura: Cycloramphidae), from the Atlantic Rainforest of eastern Bahia, Brazil. *Zootaxa* 2660: 57-67.
- Daniels, R. J. R. (2005): *Amphibians of peninsular India*. Indian Academy of Science & University Press, Hyderabad.
- Das, I. (2007): *The amphibians and reptiles of Brunei. A pocket guide*. Natural History Publications (Borneo), Koto Kinabalu.
- Das, I. & S. K. Dutta (2006): New species of *Polypedates* (Anura: Rhacophoridae) from the Western Ghats, Southwest India. *Journal of Herpetology* 40: 214-220.
- Das, I. & A. Haas (2010): New species of *Microhyla* from Sarawak: Old World's smallest frogs crawl out of miniature pitcher plants on Borneo (Amphibia: Anura: Microhylidae). *Zootaxa* 2571: 37-52.
- Das, I., A. Jankowski, M. I. B. Makmor & A. Haas (2007): Species diversity, elevational distribution and reproductive modes in an amphibian community at the Matang Range, Sarawak (Borneo). *Mitteilungen des Hamburger Zoologischen Museums und Instituts* 104: 141-171.

- de Almeida Prado, C., L. F. Toledo, J. Zina & C. F. B. Haddad (2005): Trophic eggs in the foam nests of *Leptodactylus labyrinthicus* (Anura, Leptodactylidae): an experimental approach. The Herpetological Journal 15: 279-284.
- de Almeida Prado, C. P., M. Uetanabaro, C. F. B. Haddad & C. Guyer (2002): Description of a new reproductive mode in *Leptodactylus* (Anura, Leptodactylidae), with a review of the reproductive specialization toward terrestriality in the genus. Copeia 2002: 1128-1133.
- de Almeida Prado, C. P., M. Uetanabaro & F. S. Lopes (2000): Reproductive strategies of *Leptodactylus chaquensis* and *L. podicipinus* in the Pantanal, Brazil. Journal of Herpetology 34: 135-139.
- de Carvalho, A. L. (1946): Um novo gênero de Ceratophridideo do sudeste Baiano. Boletim do Museu Nacional. Zoologia 73: 1-19.
- de la Riva, I. (1999): A new *Phyllomedusa* from southwestern Amazonia (Amphibia: Anura: Hylidae). Revista española de herpetología 13: 123-131.
- de Mattos Brito, L. B., L. Bezerra, F. Aguiar, C. Moura-Neto, C. A. Zucco & P. Cascon (2013): Diet, activity patterns, microhabitat use and defensive strategies of *Rhinella hoogmoedi* Caramaschi & Pombal, 2006 from a humid forest in northeast Brazil. The Herpetological Journal 23: 29-37.
- de Oliveira Filho, J. C., H. C. de Moura Costa & Ú. M. L. Braga (2005): Egg-laying and foam-beating in *Leptodactylus fuscus* (Anura, Leptodactylidae). Biota Neotropica 5: 1-2.
- de Oliveira Filho, J. C. & A. A. Giaretta (2008): Reproductive behavior of *Leptodactylus mystacinus* (Anura, Leptodactylidae) with notes on courtship call of other *Leptodactylus* species. Iheringia. Série Zoologia 98: 508-515.
- Dehling, J. M. & D. M. Dehling (2013) Range extension of *Rhacophorus dulitensis* Boulenger, 1892 (Amphibia: Anura: Rhacophoridae) in western Borneo. Check List 9: 425-426.
- Deichmann, J. L., W. E. Duellman & B. G. Williamson (2008): Predicting biomass from snout-vent length in New World frogs. Journal of Herpetology 42: 238-245.
- Dixo, M. (2004): Range extension of *Rhacophorus dulitensis* Boulenger, 1892 (Amphibia: Anura: Rhacophoridae) in western Borneo. Phyllomedusa 3: 77-79.
- Downie, J. R. (1996): A new example of female parental behaviour in *Leptodactylus validus*, a frog of the *Leptodactylid melanotus*' species group. Herpetological Journal 6: 32-34.
- du Preez, L. & V. Carruthers (2007): Complete Guide to the Frogs of Southern Africa. Struik Publishers, Cape Town.
- Duellman, W. E. (1960): Synonymy, variation, and distribution of *Ptychohyla leonhard-schultzei* Ahl. Studies of American Hylid frogs. IV. Herpetologica 16: 191-197.
- Duellman, W. E. (2001): Hylid Frogs of Middle America (Contribution to Herpetology). 2 edition. Society for the study of amphibians and reptiles, Ithaca, New York.
- Duellman, W. E. (2005): Cusco Amazónico — the lives of amphibians and reptiles in an Amazonian rainforest. Comstock Publishing Associates, Ithaca.
- Duellman, W. E. & M. L. Crump (1974): Speciation in frogs of the *Hyla parviceps* group in the upper Amazon Basin. Occasional Papers of the Museum of Natural History, The University of Kansas 23: 1-40.
- Duellman, W. E. & E. Lehr. (2009): Terrestrial-breeding frogs (Strabomantidae) in Peru. Natur & Tier Verlag, Münster.
- Duellman, W. E. & J. Lescure (1973): Life history and ecology of the Hylid frog *Osteocephalus taurinus*, with observations on larval behavior. Occasional Papers of the Museum of Natural History, The University of Kansas 13: 1-12.
- Duellman, W. E. & M. Lizana (1994): Biology of a sit-and-wait predator, the Leptodactylid frog *Ceratophrys cornuta*. Herpetologica 50: 51-64.
- Duellmann, W. E. (1978): The biology of an equatorial herpetofauna in Amazonian Ecuador. Miscellaneous Publication, Museum of Natural History, University of Kansas 65: 1-352.
- Ernst, R., A. C. Agyei & M.-O. Rödel (2008): A new giant species of *Arthroleptis* (Amphibia: Anura: Arthroleptidae) from the Krokosua Hills Forest Reserve, south-western Ghana. Zootaxa 1697: 58-68.

- Fabrezi, M., S. I. Quinzio & J. Goldberg (2009): Giant tadpole and delayed metamorphosis of *Pseudis platensis* Gallardo, 1961 (Anura, Hylidae). *Journal of Herpetology* 43: 228-243.
- Fernando, P. & N. Dayawansa (1995): Description of the larval stages and notes on the reproduction of *Polypedates longinasus* (Ahl, 1927) (Amphibia, Rhacophoridae). *Journal South Asian Natural History* 1: 235-240.
- Fernando, P., N. Dayawansa & M. Siriwardena (1994): *Bufo kotagamai*, a new toad (Bufonidae) from Sri Lanka. *Journal of South Asian natural History* 1: 119-124.
- Firschein, I. L. (1950): A new toad from Mexico with a redefinition of the cristatus group. *Copeia* 1950: 81-87.
- Frétey, T., M. Dewynter & C. P. Blanc (2011): Amphibiens d'Afrique centrale et d'Angola. Clé de détermination Illustrée des Amphibiens du Gabon et du Mbini. Biotope, Mèze (Collection Parthénopé) & Muséum National d'Histoire Naturelle, Paris.
- Garda, A. A., G. M. Biavati & G. C. Costa (2006): Sexual dimorphism, female fertility, and diet of *Pipa arrabali* (Anura, Pipidae) in Serra do Cachimbo, Pará, Brazil. *South American Journal of Herpetology* 1: 20-24.
- Garda, A. A., D. J. Santana & V. A. de Avelar São-Pedro (2010): Taxonomic characterization of paradoxical frogs (Anura, Hylidae, Pseudae): geographic distribution, external morphology, and morphometry. *Zootaxa* 2666: 1-28.
- Gaulke, M. (2011): The herpetofauna of Panay Island, Philippines. Edition Chimaira, Frankfurt/M.
- Gawor, A., K. van der Straeten, D. Karbe, U. Manthey & T. Ziegler (2011): Reproduction and development of the dark-sided frog *Hylarana nigrovittata* sensu lato at the Cologne Zoo. *Salamandra* 47: 1-8.
- Giaretta, A. A. & M. N. de C. Kokubum (2004): Reproductive ecology of *Leptodactylus furnarius* Sazima & Bokermann, 1978, a frog that lays eggs in underground chambers. *Herpetozoa* 16: 115-126.
- Giaretta, A. A. & K. G. Facure (2006): Terrestrial and communal nesting in *Eupemphix nattereri* (Anura, Leiuperidae): interactions with predators and pond structure. *Journal of Natural History* 40: 2577-2587.
- Giaretta, A. A. & K. G. Facure (2009): Habitat, egg-laying behaviour, eggs and tadpoles of four sympatric species of *Pseudopaludicola* (Anura, Leiuperidae). *Journal of Natural History* 43: 995-1009.
- Gillespie, G. R., M. Anstis, S. D. Howard & D. Lockie (2007): Description of the tadpole of the Rhacophorid frog *Rhacophorus georgii* Roux (Rhacophoridae) from Sulawesi, Indonesia. *Journal of Herpetology* 41: 150-153.
- Gillespie, G. R., D. Lockie, M. P. Scroggi & D. T. Iskandar (2004) Habitat use by stream-breeding frogs in southeast Sulawesi, with some preliminary observations on community organization. *Journal of Tropical Ecology* 20: 439-448.
- Girish, S. & S. K. Saidapur (2003): Density-dependent growth and metamorphosis in the larval bronze frog *Rana temporalis* is influenced by genetic relatedness of the cohort. *Journal of biosciences* 28: 489-496.
- Glaw, F. & M. Vences (1992): Zur Kenntnis der Gattungen *Boophis*, *Aglyptodactylus* und *Mantidactylus* (Amphibia: Anura) aus Madagaskar, mit Beschreibung einer neuen Art. *Bonner Zoologische Beiträge* 43: 45-77.
- Glaw, F. & M. Vences (1997): Neue Ergebnisse zur *Boophis goudoti*-Gruppe aus Madagaskar: Bioakustik, Fortpflanzungsstrategien und Beschreibung von *Boophis rufioculis* sp. nov. *Salamandra* 32: 225-242.
- Glaw, F. & M. Vences (1999): Resurrection and redescription of *Mantidactylus tricinctus* from eastern Madagascar. *Journal of Herpetology* 33: 639-647.
- Glaw, F. & M. Vences (2002): A new cryptic frog species of the *Mantidactylus boulengeri* group with a divergent vocal sac structure. *Amphibia-Reptilia* 23: 293-304.
- Glaw, F. & M. Vences (2004): A preliminary review of cryptic diversity in frogs of the subgenus *Ochthomantis* based on mtDNA sequences and morphology (Anura, Mantellidae, *Mantidactylus*). *Spixiana* 27: 83-91.

- Glaw, F. & M. Vences. (2007): A field guide to the amphibians and reptiles of Madagascar. 3 edition. Vences & Glaw Verlags GbR.
- Glaw, F., M. Vences & V. Gossmann (2000): A new species of *Mantidactylus* (subgenus *Guibemantis*) from Madagascar, with a comparative survey of internal femoral gland structure in the genus (Amphibia: Ranidae: Mantellinae). *Journal of Natural History* 34: 1135-1154.
- Gomez-Mestre, I., R. A. Pyron & J. J. Wiens (2012): Phylogenetic analyses reveal unexpected patterns in the evolution of reproductive modes in frogs. *Evolution* 66: 3687-3700.
- Gopalan, S. V., A. Nair, K. S. Kumar, J. Merilä & S. George (2012): Morphology of *Indirana semipalmata* (Boulenger, 1882) (Amphibia; Anura) adults and tadpoles from the Western Ghats, India. *Herpetology Notes* 5: 263-273.
- Goris, R. C. & N. Maeda (2004): Guide to the amphibians and reptiles of Japan. Krieger Publishing Company, Malabar.
- Grismer, L. L. (2011): Amphibians and reptiles of the Seribuat Archipelago. Edition Chimaira, Frankfurt/M.
- Grismer, L. L., T. Neang, T. Chav, P. L. Wood Jr, J. R. Oaks, J. Holden, J. L. Grismer, T. R. Szutz & T. M. Youmans (2008): Additional amphibians and reptiles from the Phnom Samkos Wildlife Sanctuary in northwestern Cardamom Mountains, Cambodia, with comments on their taxonomy and the discovery of three new species. *The Raffles Bulletin of Zoology* 56: 161-175.
- Grosjean, S., A. Strauß, J. Glos, R.-D. Randrianiana, A. Ohler & M. Vences (2011): Morphological and ecological uniformity in the funnel-mouthed tadpoles of Malagasy litter frogs, subgenus *Chonomantis*. *Zoological Journal of the Linnean Society* 162: 149-183.
- Grosjean, S., M. Thomas, F. Glaw & M. Vences (2006): The tadpole of the Malagasy treefrog *Boophis rufiocularis*: molecular identification and description. *Spixiana* 29: 73-76.
- Guimarães, T. C. S., G. B. de Figueiredo, D. O. Mesquita & M. M. Vasconcellos (2011): Ecology of *Hypsiboas albopunctatus* (Anura: Hylidae) in a neotropical savanna. *Journal of Herpetology* 45: 244-250.
- Gururaja, K. V. (2010a): Novel reproductive mode in a torrent frog *Micrixalus saxicola* (Jerdon) from the Western Ghats, India. *Zootaxa* 2642: 45-52.
- Gururaja, K. V. (2010b): Pictorial guide to frogs and toads of the Western Ghats. Sharadh Enterprises, Karnataka.
- Haas, A. & I. Das (2008): Larval identities of *Ansonia hanitschi* Inger, 1960 (Amphibia: Bufonidae) and *Polypedates colletti* (Boulenger, 1890) (Amphibia: Rhacophoridae) from East Malaysia (Borneo). *Salamandra* 44: 85-100.
- Haddad, C. F. B., L. F. Toledo, C. P. A. Prado, D. Loebmann, J. L. Gasparini & I. Sazima (2013): Guia dos Anfíbios da Mata Atlântica: Diversidade e Biologia. Anolisbooks, Sao Paulo.
- Hailey, A. (ed.) (2011): OGATT: The online guide to the animals of Trinidad and Tobago. Department of Life Sciences, The University of the West Indies, St. Augustine.
- Hamidy, A. & M. Matsui (2010): A new species of blue-eyed *Leptobrachium* (Anura: Megophryidae) from Sumatra, Indonesia. *Zootaxa* 2395: 34-44.
- Harper, E. B., G. J. Measey, D. A. Patrick, M. Menegon & J. R. Vonesh (2010): Field Guide to the Amphibians of the Eastern Arc Mountains and Coastal Forests of Tanzania and Kenya. Camerapix Publishers International, Nairobi.
- Hartmann, T., F. Ihlow, S. Edwards, S. Sothanin, M. Handschuh & W. Böhme (2013): A preliminary annotated checklist of the amphibians and reptiles of the Kulen Promtep Wildlife Sanctuary in Northern Cambodia. *Asian Herpetological Research* 4: 36-55.
- Herrmann, H.-W., W. Böhme, P. A. Herrmann, M. Plath, A. Schmitz & M. Solbach (2005): African biodiversity hotspots: the amphibians of Mt. Nlonako, Cameroon. *Salamandra* 41: 61-81.
- Heyer, W. R. (1973): Ecological interactions of frog larvae at a seasonal tropical location in Thailand. *Journal of Herpetology* 7: 337-361.
- Heyer, W. R. (1978): Systematics of the fuscus group of the frog genus *Leptodactylus* (Amphibia, Leptodactylidae). *Science Bulletin. Natural History Museum of Los Angeles County* 29: 1-85.

- Heyer, W. R. (1983): Clarification of the names *Rana mysacea* Spix, 1824, *Leptodactylus amazonicus* Heyer, 1978 and a description of a new species, *Leptodactylus spixi* (Amphibia: Leptodactylidae). *Proceedings of the Biological Society of Washington* 96: 270-272.
- Heyer, W. R. (1988): A notable collection of *Cycloramphus* (Amphibia: Leptodactylidae) from Bahia, Brazil, with a description of a new species (*Cycloramphus migueli*). *Proceedings of the Biological Society of Washington* 101: 151-154.
- Heyer, W. R. (1994): Variation within the *Leptodactylus podicipinus-wagneri* complex of frogs (Amphibia: Leptodactylidae). *Smithsonian Contributions to Zoology* 546: 1-124.
- Heyer, W. R. & M. S. Bellin (1973): Ecological notes on five sympatric *Leptodactylus* (Amphibia, Leptodactylidae) from Ecuador. *Herpetologica* 29: 66-72.
- Heyer, W. R. & A. A. Giaretta (2009): Advertisement calls, notes on natural history, and distribution of *Leptodactylus chaquensis* (Amphibia: Anura: Leptodactylidae) in Brasil. *Proceedings of the Biological Society of Washington* 122: 292-305.
- Heyer, W. R. & M. M. Heyer (2002): *Leptodactylus elenae* Heyer. *Catalogue of American Amphibians and Reptiles* 742: 1-5.
- Heyer, W. R. & M. M. Heyer (2006): *Leptodactylus knudseni* Heyer. Knudsen's thin-toed frog. *Catalogue of American Amphibians and Reptiles* 807: 1-12.
- Heyer, W. R. & M. M. Heyer (2012): Systematics, distribution, and bibliography of the frog *Leptodactylus validus* (Amphibia: Leptodactylidae). *Proceedings of the Biological Society of Washington* 125: 276-294.
- Heyer, W. R., M. M. Heyer & R. O. de Sá (2010): *Leptodactylus savagei* Heyer. Savage's thin-toed frog. *Catalogue of American Amphibians and Reptiles* 867: 1-19.
- Heyer, M. M., W. R. Heyer, S. Spear & R. O. de Sá (2003): *Leptodactylus mystacinus* (Burmeister), Mustached Frog. *Catalogue of American Amphibians and Reptiles* 767: 1-11.
- Heyer, W. R. & A. J. Wolf (1989): *Physalaemus crombiei* (Amphibia, Leptodactylidae), a new frog species from Espírito-Santo, Brazil with comments on the *Physalaemus signifer* group. *Proceedings of the Biological Society of Washington* 102: 500-506.
- Hirschfeld, M., M. F. Barej, S. P. Loader & M.-O. Rödel (2012): Description of two *Werneria* tadpoles from Cameroon (Amphibia: Anura: Bufonidae). *Zootaxa* 3172: 65-68.
- Hödl, W. (1990): Reproductive diversity in Amazonian lowland frogs. pp 41-60 in W. Hanke (ed.). *Biology and physiology of amphibians*. Gustav Fischer Verlag, Stuttgart, New York.
- Hoffmann, P. (1995): Zur Kenntniss des Engmaulfrosches *Chaperina fusca* Moquaard, 1892 vom Mt. Kinabalu auf Borneo. *Herpetofauna* 17: 27-29.
- Hoogmoed, M., D. Borges & P. Cascon (1994): Three new species of the genus *Adelophryne* (Amphibia: Anura: Leptodactylidae) from northeastern Brazil, with remarks on the other species of the genus. *Zoologische Mededelingen* 68: 271-300.
- Inger, R. F., H. B. Shaffer, M. Koshy & R. Bakde (1984): A report on a collection of amphibians and reptiles from the Ponmudi, Kerala, South India. *The Journal of the Bombay Natural History Society* 81: 551-570.
- Inger, R. F., B. L. Stuart & D. T. Iskandar (2009): Systematics of a widespread Southeast Asian frog, *Rana chalconota* (Amphibia: Anura: Ranidae). *Zoological Journal of the Linnean Society* 155: 123-147.
- Inger, R. F. & R. B. Stuebing (1997): A field guide to the frogs of Borneo. 2 edition. Natural History Publications (Borneo), Koto Kinabalu.
- Iskandar, D. T., B. J. Evans & J. A. McGuire (2014): A novel reproductive mode in frogs: a new species of fanged frog with internal fertilization and birth of tadpoles. *PLoS ONE* 9: e115884.
- Jansen, M., J. Q. Vidoz & O. Helmig (2008): *Physalaemus cuvieri*. *Herpetological Review* 39: 106-107.
- Jones, C. (1971): Notes on hairy frogs (*Trichobatrachus robustus* Boulenger) collected in Rio Muni, West Africa. *Herpetologica* 27: 51-54.
- Jungfer, K.-H. & W. Hödl (2002): A new species of *Osteocephalus* from Ecuador and a redescription of *O. lepieurii* (Duméril & Bibron, 1841) (Anura: Hylidae). *Amphibia-Reptilia* 23: 21-46.
- Jungfer, K.-H. & P. Weygoldt (1999): Biparental care in the tadpole-feeding Amazonian treefrog *Osteocephalus oophagus*. *Amphibia-Reptilia* 20: 235-249.

- Kabir, M., K. Hasa & M. Ahmed (2010): *Chiromantis vittatus* recorded for the first time in Bangladesh. frog leg 14: 2-3.
- Kadadevaru, G. G. & R. D. Knamandi (2000): Courtship and nesting behaviour of the Malabar gliding frog *Rhacophorus malabaricus* (Jerdon, 1870). Current Science 79: 377-380.
- Kaplan, M. & P. Heimes (2011): Tadpole of *Plectrohyla arborescandens*, with comments on the identity of the tadpole of *Plectrohyla cyclada*. Journal of Herpetology 45: 463-464.
- Kaplan, M. & P. M. Ruíz (1997): Two new species of *Hyla* from the Andes of Central Colombia and their relationships to other small Andean Hyla. Journal of Herpetology 31: 230-244.
- Karsen, S. J., M. W.-N. Lai & A. Bogadek (1986): Hong Kong amphibians and reptiles. Urban council, Hong Kong.
- Kehr, A. I. & V. I. Gómez (2009): Intestinal, body and tail plasticity in *Rhinella schneideri* (Bufonidae) tadpoles induced by a predator insect (*Belostoma elegans*). Advanced Studies in Biology 1: 85-94.
- Knoll, A., J. Köhler, F. Glaw, M. Teschke & M. Vences (2007): Larval morphology in four species of Madagascan frogs of the subgenus *Brygroomantis* (Mantellidae: Mantidactylus). Zootaxa 1616: 49-59.
- Kobel, H. R., L. du Pasquier & H. Gloor (1980): *Xenopus amieti* sp. nov (Anura: Pipidae) from the Cameroon, another case of tetraploidy. Revue suisse de Zoologie 87: 919-926.
- Koch, A., E. Arida, A. Riyanto & W. Böhme (2009): Islands between the realms: a revised checklist of the herpetofauna of the Talaud Archipelago, Indonesia, with discussion about its biogeographic affinities. Bonner Zoologische Beiträge 56: 107-129.
- Köhler, G. (2011): Amphibians of Central America. Herpeton, Offenbach.
- Köhler, J. (2000): Amphibian diversity in Bolivia: a study with special reference to montane forest regions. Bonner Zoologische Monographien 48: 1-243.
- Köhler, J., M. Vences, N. D'Cruze & F. Glaw (2010): Giant dwarfs: discovery of a radiation of large-bodied 'stump-toed frogs' from karstic cave environments of northern Madagascar. Journal of Zoology 282: 21-38.
- Kok, P. J. R. & M. Kalamandeen (2008): Introduction to the taxonomy of the amphibians of Kaieteur National Park, Guyana. ABC Taxa 5: 1-278.
- Kouamé, N. G. N., B. Tohé, N. G. E. Assemian, G. Gourène & M.-O. Rödel (2008): Prey composition of two syntopic *Phrynobatrachus* species in the swamp forest of Banco National Park, Ivory Coast. Salamandra 44: 177-186.
- Krishna, S. N. & S. B. Krishna (2005): Female courtship calls of the litter frog (*Rana curtipes*) in the tropical forests of Western Ghats, South India. Amphibia-Reptilia 26: 431-435.
- Krishna, S. N., S. B. Krishna & K. K. Vijayalaxmi (2004): Breeding ecology of a rare microhylid, *Ramanella montana*, in the forests of Western Ghats, India. Current Science 87: 80-82.
- Krishnamurthy, S. V., A. H. Manjunatha Reddy & K. V. Gururaja (2001): A new species of frog in the genus *Nyctibatrachus* (Anura: Ranidae) from Western Ghats, India. Current Science 80: 887-891.
- Kubicki, B. (2007): Ranas de vidrio — Costa Rica — glass frogs. Instituto Nacional de Biodiversidad, Santo Domingo de Heredia.
- Lambiris, A. J. L. (1989): The frogs of Zimbabwe. Museo Regionale di Scienze Naturali, Torino.
- Laufer, G. & J. M. Barreneche (2008): Re-description of the tadpole of *Pseudopaludicola falcipes* (Anura: Leiuperidae), with comments on larval diversity of the genus. Zootaxa 1760: 50-58.
- Lee, J. C. (1996): The amphibians and reptiles of the Yucatan Peninsula. Comstock Publishing Associates., Ithaca, New York.
- Legler, J. M. (1964): A narrow-mouthed toad (*Gastrophryne usta*) in El Salvador. Herpetologica 19: 286-287.
- Lehtinen, R. M. (2003): Parental care and reproduction in two species of *Mantidactylus* (Anura: Mantellidae). Journal of Herpetology 37: 766-768.
- Lemos-Espinal, J. A. & J. R. Dixon (2013): Amphibians and reptiles of San Luis Potosi. Eagle Mountain Publishing, Eagle Mountain.
- Lescure, J. & C. Marty (2000): Atlas des amphibiens de Guyane. Collections Patrimoines Naturels. Paris 45: 1-388.

- Lima, A. P., W. E. Magnusson, M. Menin, L. K. Erdtmann, D. J. Rodrigues, C. Keller & W. Hödl (2005): Guia de Sapos da Reserva Adolpho Ducke, Amazonia Central. Instituto Nacional de Pesquisas da Amazônia, Manaus.
- Lisboa, B. S., F. A. C. do Nascimento & G. O. Skuk (2011): Redescription of the tadpole of *Macrogenioglottus alipioi* (Anura: Cycloramphidae), a rare and endemic species of the Brazilian Atlantic Forest. *Zootaxa* 3046: 67-68.
- Loader, S. P., G. J. Measey, R. O. de Sa & P. K. Malonza (2009): A new brevipitid species (Brevipitidae: *Callulina*) from the fragmented forests of the Taita Hills, Kenya. *Zootaxa* 2123: 55-68.
- Lobo, F. (1994): Descripción de una nueva especie de *Pseudopaludicola* (Anura: Leptodactylidae), redescrpción de *P. falcipes* (Hensel, 1867) y *P. saltica* (Cope, 1887) y osteología de las tres especies. *Cuadernos de herpetología* 8: 177-199.
- Lötters, S., W. Haas, S. Schick & W. Böhme (2002): On the systematics of the harlequin frogs (Amphibia: Bufonidae: Atelopus) from Amazonia. II: Redescription of *Atelopus pulcher* (Boulenger, 1882) from the eastern Andean versant in Peru. *Salamandra* 38: 165-184.
- Lötters, S., K.-H. Jungfer, F. W. Henkel & W. Schmidt (2007): Pfeilgiftfrösche - Biologie, Haltung, Arten. Edition Chimaira, Frankfurt/M.
- Lucas, E. M., C. A. Brasileiro, H. M. Oyamaguchi & M. Martins (2008): The reproductive ecology of *Leptodactylus fuscus* (Anura, Leptodactylidae): new data from natural temporary ponds in the Brazilian Cerrado and a review throughout its distribution. *Journal of Natural History* 42: 2305-2320.
- Lynch, J. D. (1968): Systematic status of some Andean leptodactylid frogs with a description of a new species of *Eleutherodactylus*. *Herpetologica* 24: 289-300.
- Lynch, J. D. (1969): Identity of two Andean *Eleutherodactylus* with the description of a new species (Amphibia: Leptodactylidae). *Journal of Herpetology* 33: 135-143.
- Lynch, J. D. (1970): A taxonomic revision of the leptodactylid frog genus *Syrrophus* Cope. University of Kansas Publications, Museum of Natural History 20: 1-45.
- Lynch, J. D. (1975): A review of the broad-headed eleutherodactyline frogs of South America (Leptodactylidae). *Occasional Papers of the Museum of Natural History, The University of Kansas* 38: 1-46.
- Lynch, J. D. (1984) New frogs (Leptodactylidae: *Eleutherodactylus*) from cloud forests in the Northern Cordillera Oriental, Colombia. Milwaukee Public Museum. *Contributions in Biology and Geology* 60: 1-19.
- Lynch, J. D. (1994): Two new species of the *Eleutherodactylus conspicillatus* group (Amphibia: Leptodactylidae) from the Cordillera Oriental of Colombia. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 19: 187-193.
- Lynch, J. D. (1998): New species of *Eleutherodactylus* from the Cordillera Occidental of western Colombia with a synopsis of the distribution of species in western Colombia. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 22: 117-148.
- Lynch, J. D. (2000): The relationships of an ensemble of Guatemalan and Mexican frogs (*Eleutherodactylus*: Leptodactylidae: Amphibia). *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 24: 129-156.
- Lynch, J. D. (2001): Three new rainfrogs of the *Eleutherodactylus diastema* group from Colombia and Panama. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 25: 287-297.
- Lynch, J. D. & M. C. Ardila-Robayo (1999): The *Eleutherodactylus* of the taeniatus complex in western Colombia: taxonomy and distribution. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 23: 615-624.
- Lynch, J. D. & W. E. Duellman (1980): The *Eleutherodactylus* of the Amazonian slopes of the Ecuadorian Andes (Anura: Leptodactylidae). The University of Kansas. Museum of Natural History. *Miscellaneous Publications*. 69: 1-86.

- Lynch, J. D. & W. E. Duellman (1997): Frogs of the Genus *Eleutherodactylus* in Western Ecuador. Systematics, Ecology, and Biogeography. The University of Kansas. Museum of Natural History. Special Publications. 23: 1-236.
- Lynch, J. D. & M. S. Hoogmoed (1977): Two new species of *Eleutherodactylus* (Amphibia: Leptodactylidae) from northeastern South America. *Proceedings of the Biological Society of Washington* 90: 424-439.
- Lynch, J. D. & L. Trueb (1980): A new species of *Eleutherodactylus* (Leptodactylidae) from the cloud forests of western Ecuador. *Copeia* 1980: 392-396.
- Maffei, F. & F. K. Ubaid (2014): Amphibians of Rio Claro Farm. Lençóis Paulista, São Paulo, Brazil. Canal 6, Bauru, SP. Maiditsch, I., H. C. Liedtke, J. M. Ng'wava & W. Hoedl (2011): Advertisement and close-range encounter call of *Arthroleptis schubotzi* Nieden, 1911, with notes on phonotaxis and sexual dimorphism in the third manual digit. *Herpetozoa* 24: 23-31.
- Malkmus, R. & J. M. Dehling (2008): Anuran amphibians of Borneo as phytotelm-breeders—a synopsis. *Herpetozoa* 20: 165-172.
- Malkmus, R., U. Manthey, G. Vogel, P. Hoffmann & J. Kosuch (2002): Amphibians and reptiles of Mount Kinabalu (North Borneo). A.R.G. Gantner Verlag K.G, Ruggell.
- Malone, J. H. (2004): Reproduction in three species of *Smilisca* from Costa Rica. *Journal of Herpetology* 38: 27-35.
- Manamendra-Arachchi, K. & D. Gabadage (1996): *Limnonectes kirtisinghei*, a new species of ranid frog from Sri Lanka. *Journal of South Asian natural History* 2: 31-42.
- Manamendra-Arachchi, K. & R. Pethiyagoda (2005): The Sri Lankan shrub-frogs of the genus *Philautus* Gistel, 1848 (Ranidae: Rhacophorinae), with description of 27 new species. *Raffles Bulletin of Zoology* 12: 163-303.
- Manthey, U. & W. Grossmann (1997): Amphibien & Reptilien Südostasiens. Natur und Tier Verlag, Münster.
- Marin da Fonte, L. F. (2010): Variacao morfológica e na estrutura do canto em *Scinax granulatus* (Peters, 1871) (Anura, Hylidae). Universidade Federal do Rio Grande do Sul, Porto Alegre.
- Martins, I. A. (2001): Parental care behaviour in *Leptodactylus podicipinus* (Cope, 1862) (Anura, Leptodactylidae). *Herpetological Journal* 11: 29-32.
- Mathe, R. & N. Sen (2010): Pictorial guide to amphibians of North East India. Zoological Survey of India, Kolkata.
- Measey, G. J., P. K. Malonza & V. Muchai (2009): Amphibians of the Taita Hills. SANBI Biodiversity Series 12. South African National Biodiversity Institute, Pretoria.
- Mendelson III, J. R. (1997): A new species of toad (Anura: Bufonidae) from Oaxaca, Mexico with comments on the status of *Bufo cavifrons* and *Bufo cristatus*. *Herpetologica* 53: 268-286.
- Menin, M., D. J. Rodrigues & A. P. Lima (2007): Clutches, tadpoles and advertisement calls of *Synapturanus mirandaribeiroi* and *S. cf. salseri* in Central Amazonia, Brazil. *The Herpetological Journal* 17: 86-91.
- Mercurio, V. (2011): Amphibians of Malawi. An analysis of the richness and community diversity in a changing landscape. Edition Chimaira, Frankfurt/M.
- Mercurio, V., W. Böhme & B. Streit (2009): Reproductive diversity of Malawian anurans. *Herpetology Notes* 22: 175-183.
- Mertens, R. (1965): Die Amphibian von Fernando Poo. *Bonner Zoologische Beiträge* 16: 14-29.
- Minter, L. R., M. Burger, J. A. Harrison, H. H. Braack, B. P. J. & D. Knoepfer (2004): Atlas and red data book of the frogs of South Africa, Lesotho & Swaziland. SI/MAB Biodiversity Program Series #9, Smithsonian Institution, Washington D.C.
- Moura, G. J. B., E. V. E. Andrade & E. M. X. Freire (2010): Amphibia, Anura, Microhylidae, *Stereocyclops incrassatus* Cope, 1870: Distribution extension. *Check List* 6: 71-72.
- Muniz, K. P. R., A. A. Giaretta, W. R. Silva & K. G. Facure (2008): Auto-ecologia de *Hypsiboas albopunctatus* (Anura, Hylidae) em área de Cerrado no sudeste do Brasil. *Iheringia, Série Zoologia* 98: 254-259.

- Murrieta-Galindo, R., A. González-Romero, F. López-Barrera & G. Parra-Olea (2013): Coffee agrosystems: an important refuge for amphibians in central Veracruz, Mexico. *Agroforestry Systems* 87: 1-13.
- Myers, C. W. (1969): The ecological geography of cloud forest in Panama. *American Museum novitates* 2396: 1-52.
- Myers, C. W. & M. A. Donnelly (2001): Herpetofauna of the Yutaje-Corocoro massif, Venezuela: second report from the Robert G. Goellet American Museum-Terramar expedition to the northwestern Tepuis. *Bulletin of the American Museum of Natural History* 261: 1-85.
- Nair, A., S. V. Gopalan, S. George, K. S. Kumar, A. G. F. Teacher & J. Merilä (2012): Endemic Indirana frogs of the Western Ghats biodiversity hotspot. *Annales Zoologici Fennici* 49: 257-286.
- Narzary, J. & S. Bordoloi (2013): Study of normal development and external morphology of tadpoles of *Microhyla ornata* and *Uperodon globulosus* of the family Microhylidae (Amphibia: Anura) from North East India. *International Journal of Advanced Biological Research* 3: 61-73.
- Nascimento, L. B., B. V. S. Pimenta, C. A. G. Cruz & U. Caramaschi (2006): Taxonomic status of *Gomphobates marmoratus* Reinhardt and Lütken, 1862 "1861" and *Eupemphix fuscomaculatus* Steindachner, 1864 (Amphibia, Anura, Leptodactylidae). *South American Journal of Herpetology* 1: 166-174.
- Norman, D. R. (1994): *Anfibios y Reptiles del Chaco Paraguayo*, Tomo 1. (Amphibians and reptiles of the Paraguayan Chaco, Volume 1.). Private printing, San José, Costa Rica.
- Ohler, A., S. R. Swan & J. C. Daltry (2002): A recent survey of the amphibian fauna of the Cardomom Mountains, Southwest Cambodia with descriptions of three new species. *Raffles Bulletin of Zoology* 50: 465-482.
- Orlow, N. L., S. A. Ryabov, N. B. Ananjeva & A. A. Evsyunin (2010): Asian treefrogs of the genus *Theloderma* Tschudi, 1838 (Amphibia: Anura: Rhacophoridae: Rhacophorinae). *Russian Academy of Sciences, Zoological Institute, St. Petersburg*.
- Ouboter, P. E. & R. Jairam (2012): *Amphibians of Suriname*. Brill, Leiden.
- Peracca, D. M. G. (1904): Rettili ed Anfibi. Facsimiles. *Bollettino dei Musei di Zoologia ed Anatomia comparata della R. Università di Torino* 19: 429-469.
- Perret, J.-L. (1966): Les amphibiens du Cameroun. *Zoologische Jahrbücher (Systematik)* 8: 289-464.
- Perret, J.-L. (1977): Les Hylarana (Amphibiens, Ranidés) du Cameroun. *Revue suisse de Zoologie* 84: 841-868.
- Perret, J.-L. (1983): Nouvelles données sur *Hylarana occidentalis* Perret (Amphibia, Ranidae). *Bulletin de la Société Neuchâteloise des Sciences Naturelles* 106: 109-113.
- Perret, J.-L. (1988): Les espèces de *Phrynobatrachus* (Anura, Ranidae) à éperon palpébral. *Archives des Sciences Genève* 41: 275-294.
- Pickersgill, M. (2007): A redefinition of *Afraxalus fulvovittatus* (Cope, 1860) and *Afraxalus vittiger* (Peters, 1876) (Amphibia, Anura Hyperoliidae). *African Journal of Herpetology* 56: 23-37.
- Pillai, R. S. & R. Pattabiraman (1990): Amphibians from Sabarigiri forest, Western Ghats, Kerala, including a new species of *Micrixalus*. *Records of Zoological Survey India* 86: 383-390.
- Pimenta, B. V. S., C. A. G. da Cruz & M. Dixo (2002): *Chiasmocleis carvalhoi*. *Herpetological Review* 33: 219.
- Pineda, E. & C. A. Rodríguez-Mendoza (2010): Distribución y abundancia de *Craugastor vulcani*: una especie de rana en riesgo de Los Tuxtlas, Veracruz, México. *Revista mexicana de biodiversidad* 81: 133-141.
- Poynton, J. C. (1964): Amphibia of southern Africa; a faunal study. *Annals of the Natal Museum* 17: 1-334.
- Preininger, D., M. Boeckle, A. Freudmann, I. Starnberger, M. Sztatecsny & W. Hödl (2013): Multimodal signaling in the small torrent frog (*Micrixalus saxicola*) in a complex acoustic environment. *Behavioral ecology and sociobiology* 67: 1449-1456.
- Proy, C. (1992): Zur Biologie von *Agalychnis saltator* Taylor, 1955. *Herpetozoa* 5: 99-107.
- Pupin, N. C., J. L. Gasparini, R. P. Bastos, C. F. B. Haddad & C. de Almeida Prado (2010): Reproductive biology of an endemic *Physalaemus* of the Brazilian Atlantic forest, and the trade-off between

- clutch and egg size in terrestrial breeders of the *P. signifer* group. The Herpetological Journal 20: 147-156.
- Rabb, G. B. & M. S. Rabb (1963): On the behavior and breeding biology of the African pipid frog: *Hymenochirus boettgeri*. Zeitschrift für Tierpsychologie 20: 215-241.
- Randrianiaina, R.-D., L. Raharivololoniaina, C. Preuss, A. Strauss, F. Glaw, M. Teschke, J. Glos, N. Raminosoa & M. Vences (2009): Descriptions of the tadpoles of seven species of Malagasy treefrogs, genus *Boophis*. Zootaxa 2021:23-41.
- Randrianiaina, R. D., K. C. Wollenberg, T. Rasolonjatovo Hiobiarilanto, A. Strauß, J. Glos & M. Vences (2011): Nidicolous tadpoles rather than direct development in Malagasy frogs of the genus *Gephyromantis*. Journal of Natural History 45: 2871-2900.
- Rivero, J. A. (1971): Tres nuevos records y una nueva especie de anfibios de Venezuela. Caribbean Journal of Science 11: 1-9.
- Roberto, I. J., L. Brito & P. Cascon (2011): Temporal and spatial patterns of reproductive activity in *Rhinella hoogmoedi* (Anura: Bufonidae) from a tropical rainforest in Northeastern Brazil, with the description of its advertisement call. South American Journal of Herpetology 6: 87-97.
- Rödel, M.-O. (2000): Herpetofauna of West Africa. Vol I Amphibians of the West African savanna. Edition Chimaira, Frankfurt.
- Rödel, M.-O. & W. R. Branch (2002): Herpetological survey of the Haute Dodo and Cavally forests, western Ivory Coast, Part I: amphibians. Salamandra 38: 245-268.
- Rödel, M.-O. & R. Ernst (2001): Description of the tadpole of *Kassina lamottei* Schiøtz, 1967. Journal of Herpetology 35: 678-681.
- Rödel, M.-O. & R. Ernst (2002a): A new *Phrynobatrachus* from the Upper Guinean rain forest, West Africa, including a description of a new reproductive mode for the genus. Journal of Herpetology 36: 561-571.
- Rödel, M.-O. & R. Ernst (2002b): A new reproductive mode for the genus *Phrynobatrachus*: *Phrynobatrachus alticola* has nonfeeding, nonhatching tadpoles. Journal of Herpetology 36: 121-125.
- Rödel, M.-O., M. Gil, A. Cudjoe Agyei, A. D. Leaché, R. E. Diaz, M. K. Fujita & R. Ernst (2005): The amphibians of the forested parts of south-western Ghana. Salamandra 41: 107-127.
- Rödel, M.-O., J. Kosuch, M. Veith & R. Ernst (2003): First record of the genus *Acanthixalus* Laurent, 1944 from the upper Guinean rain forest, West Africa, with the description of a new species. Journal of Herpetology 37: 43-52.
- Rödel, M.-O., D. Krätz & R. Ernst (2002a): The tadpole of *Ptychadena aequiplicata* (Werner, 1898) with the description of a new reproductive mode for the genus (Amphibia, Anura, Ranidae). Alytes 20: 1-12.
- Rödel, M.-O., F. Range, J.-T. Seppänen & R. Noë. (2002b): Caviar in the rain forest: monkeys as frog-spawn predators in Taï National Park, Ivory Coast. Journal of Tropical Ecology 18: 289-294.
- Rödel, M.-O., V. H. W. Rudolf, S. Frohshammer, K. E. Linsenmair & R. M. Lehtinen (2004a): Life history of a West-African tree-hole breeding frog, *Phrynobatrachus guineensis* Guibé and Lamotte, 1961 (Amphibia: Anura: Petropedetidae). Miscellaneous Publications Museum of Zoology University of Michigan 193: 31-44.
- Rödel, M.-O., A. Schmitz, O. S. G. Pauwels & W. Böhme (2004b): Revision of the genus *Werneria* Poche, 1903, including the description of two new species from Cameroon and Gabon (Amphibia: Anura: Bufonidae). Zootaxa 720: 1-28.
- Rödel, M.-O. & M. Spieler (2000): Trilingual keys to the savannah-anurans of the Comoé National Park, Ivory Coast. Stuttgarter Beiträge zur Naturkunde Serie A 620: 1-31.
- Rodrigues, D. J., M. Menin & A. P. Lima. 2007. Redescription of the tadpole of *Leptodactylus rhodomystax* (Anura: Leptodactylidae) with natural history notes. Zootaxa 1509: 61-67.
- Rodrigues, D. J., M. Uetanabaro & F. S. Lopes (2005): Reproductive patterns of *Trachycephalus venulosus* (Laurenti, 1768) and *Scinax fuscovarius* (Lutz, 1925) from the Cerrado, Central Brazil. Journal of Natural History 39: 3217-3226.
- Rodriguez, L. O. & W. E. Duellman (1994): Guide to the frogs of the Iquitos region, Amazonian Peru. Natural History Museum, University of Kansas, Lawrence.

- Roelke, C. E., E. Greenbaum, C. Kusamba, M. M. Aristote & E. N. Smith (2011): Systematics and conservation status of two distinct Albertine Rift treefrogs, *Leptopelis karissimbensis* and *Leptopelis kivuensis* (Anura: Arthroleptidae). *Journal of Herpetology* 45: 343-351.
- Rojas-Rivera, A., S. Cortés-Bedoya & P. D. A. Gutiérrez-Cárdenas (2011): *Pristimantis achantinus* (cachabi robber frog). Parental care and clutch size. *Herpetological Review* 42: 588-589.
- Roux, J. (1904): Reptilien und Amphibien aus Celebes. *Verhandlungen der Naturforschenden Gesellschaft in Basel* 15: 425-433.
- Ruas, D. S., C. V. M. Mendes, B. B. Szpeiter & M. Sole (2012): The tadpole of *Rhinella crucifer* (Wied-Neuwied, 1821) (Amphibia: Anura: Bufonidae) from southern Bahia, Brazil. *Zootaxa* 3299: 66-68.
- Rueda Almonacid, J. V., J. V. Rodríguez Mahecha, E. La Marca, S. Lötters, T. Kahn & A. Angulo (eds.) (2005): *Ranas Arlequines*. Conservación Internacional.
- Sabater-Pi, J. (1984): Contribution to the biology of the giant frog (*Conraua goliath*, Boulenger). *Amphibia-Reptilia* 6: 143-153.
- Sánchez, D. A. (2010): Larval development and synapomorphies for species groups of *Hyloscirtus* Peters, 1882 (Anura: Hylidae: Cophomantini). *Copeia* 2010: 351-363.
- Santana, D. J., A. P. Motta, R. M. Pirani, E. T. da Silva & R. N. Feio (2012): Advertisement call and tadpole of *Chiasmocleis mantiqueira* Cruz, Feio and Cassini, 2007 (Anura, Microhylidae). *Journal of Herpetology* 46: 14-18.
- Sarkar, A. K. & D. P. Sanyal (1985): Amphibia. *Records of Zoological Survey India* 82: 285-295.
- Savage, J. M. (1984): A new species of montanae rain frog, genus *Eleutherodactylus* (Leptodactylus) from Guerrero, Mexico. *Amphibia-Reptilia* 5: 253-260.
- Savage, J. M. (2002): *The amphibians and reptiles of Costa Rica*. University of Chicago Press, Chicago.
- Schiøtz, A. (1999): *Treefrogs of Africa*. Edition Chimaira, Frankfurt/M.
- Schlüter, A. & A. W. Salas (1991): Reproduction, tadpoles, and ecological aspects of three syntopic microhylid species from Peru (Amphibia: Microhylidae). *Stuttgarter Beiträge zur Naturkunde* 458: 1-17.
- Schmidt, H., A. Strauss, F. Glaw, M. Teschke & M. Vence. (2009): Description of tadpoles of five frog species in the subgenus *Brygoomantis* from Madagascar (Mantellidae: Mantidactylus). *Zootaxa* 1988: 48-60.
- Shimizu, S. & H. Ota (2003): Normal development of *Microhyla ornata*: the first description of the complete embryonic and larval stages for the microhylid frogs (Amphibia: Anura). *Current Herpetology* 22: 73-90.
- Smith, H. M. (1939): Mexican herpetological novelties. *Proceedings of the Biological Society of Washington* 52: 187-196.
- Sokol, O. M. (1962): The tadpole of *Hymenochirus boettgeri*. *Copeia* 1962: 272-284.
- Spawls, S., K. Howell & R. C. Drewes (2006): *Pocket guide to the reptiles and amphibians of East Africa*. A. & C. Black., London.
- Staniszewski, M. (2001): *Mantellas*. Edition Chimaira, Frankfurt/M.
- Stewart, M. M. (1967): *The amphibians of Malawi*. State University Press, New York.
- Tapley, B., C. Babburjung Purushotham & S. Girgin (2011): *Indirana semipalmata* (brown leaping frog). Reproduction. *Herpetological Review* 42: 87-88.
- Targino, M. & J. P. Pombal (2011): Redescription and variation of *Hyophryne histrio* Carvalho, 1954, an enigmatic microhylid frog from the Atlantic Rainforest of Brazil. *Amphibia-Reptilia* 32: 465-475.
- Taylor, E. H. (1938): New species of Mexican tailless amphibia. *The University of Kansas. Science Bulletin* 25: 285-405.
- Taylor, E. H. (1950): A new bromeliad frog from the Mexican state of Veracruz. *Copeia* 1950: 274-276.
- Taylor, E. H. (1962): The amphibian fauna of Thailand. *University of Kansas Science Bulletin* 43: 265-599.
- Teixeira, R. L., D. Vrcibradic & G. I. Almeida (2006): Food habits of *Stereocyclops incrassatus* (Anura, Microhylidae) from Povoação, Espírito Santo State, southeastern Brazil. *Boletim do Museu Mello Leitão* 19: 53-58.
- Tessa, G., F. Mattioli, V. Mercurio & F. Andreone (2009): Egg numbers and fecundity traits in nine species of *Mantella* poison frogs from arid grasslands and rainforests of Madagascar (Anura: Mantellidae). *Madagascar Conservation & Development* 4: 113-119.

- Toledo, L. F., L. M. Castanho & C. F. B. Haddad (2005): Recognition and distribution of *Leptodactylus mystaceus* (Anura; leptodactylidae) in the state of São Paulo, southeastern Brazil. *Biota Neotropica* 5: 57-62.
- Trueb, L. & D. Massemin (2001): The osteology and relationships of *Pipa aspera* (Amphibia: Anura: Pipidae), with notes on its natural history in French Guiana. *Amphibia-Reptilia* 22: 33-54.
- Truong, N. Q. & N. V. Khoi (2008): First record of *Theloderma stellatum* Taylor, 1962 from Phu Quoc Island, Kien Giang Province, southern Vietnam. *Herpetology Notes* 1: 61-62.
- Tyler, M. J. & F. Knight (2009): Field guide to the frogs of Australia. CSIRO Publishing, Collingwood, Victoria.
- Ukuwela, K. D. B. (2009): Description of the tadpole of *Hylarana aurantiaca* (Anura: Ranidae) from Sri Lanka. *Zootaxa* 2016: 67-68.
- Ukuwela, K. D. B. & I. N. Bandara (2009): The first description of the tadpole of *Lankanectes corrugatus* (Peters, 1864) (Anura: Nyctibatrachidae) from Sri Lanka. *Russian Journal of Herpetology* 16: 213-216.
- Urbina-Cardona, J. N. & V. H. Reynoso (2009): Uso del microhábitat por hembras grávidas de la rana de hojarasca *Craugastor loki* en la selva alta perennifolia de Los Tuxtlas, Veracruz, México. *Revista mexicana de biodiversidad* 80: 571-573.
- Veeranagoudar, D. K., R. S. Radder, B. A. Shanbhag & S. K. Saidapur (2009): Jumping behavior of semiterrestrial tadpoles of *Indirana beddomii* (Günth.): relative importance of tail and body size. *Journal of Herpetology* 43: 680-684.
- Veeranagoudar, D. K., B. A. Shanbhag & S. K. Saidapur (2004): Foraging behaviour in tadpoles of the bronze frog *Rana temporalis*: experimental evidence for the ideal free distribution. *Journal of Biosciences* 29: 201-207.
- Vejarano, S., M. Thomas & M. Vences (2006): Comparative tadpole morphology in three species of frogs of the genus *Spinomantis* (Amphibia: Mantellidae). *Contributions to Zoology* 75: 99-108.
- Venâncio, N. M. & P. R. Melo-Sampaio (2010): Reproductive behavior of the giant leaf frog *Phyllomedusa bicolor* (Anura: Hylidae) in the western Amazon. *Phyllomedusa* 9: 63-67.
- Vences, M., F. Glaw & C. Zapp (1999): Bemerkungen zu Eizahlen und Eigrößen bei madagassischen Fröschen der Gattungen *Tomopterna*, *Aglyptodactylus*, *Boophis* und *Mantidactylus* (Amphibia: Ranidae). *Salamandra* 35: 77-82.
- Vences, M., C. J. Raxworthy, R. A. Nussbaum & F. Glaw (2003): New microhylid frog (Plethodontohyla) from Madagascar, with semi-arboreal habits and possible parental care. *Journal of Herpetology* 37: 629-636.
- Verdade, V. K. & M. T. Rodrigues (2007): Taxonomic review of *Allobates* (Anura, Aromobatidae) from the Atlantic Forest, Brazil. *Journal of Herpetology* 41: 566-580.
- Vitt, L. J. & J. P. Caldwell (2013): Herpetology: an introductory biology of amphibians and reptiles. 4 edition. Academic Press, London.
- Vonesh, J. R. & S. Ross. (2000): Dipteran predation on the arboreal eggs of four Hyperolius frog species in western Uganda. *Copeia* 2000: 560-566.
- Wang, Y.-Y., T.-D. Zhang, J. Zhao, Y.-H. Sung, J.-H. Yang, H. Pang & Z. Zhang (2012): Description of a new species of the genus *Xenophrys* Günther, 1864 (Amphibia: Anura: Megophryidae) from Mount Jinggang, China, based on molecular and morphological data. *Zootaxa* 3546: 53-67.
- Werner, F. (1899): Über Reptilien und Batrachier aus Columbien und Trinidad. *Verhandlungen der kaiserlich-königlichen zoologisch-botanischen Gesellschaft in Wien* 49: 470-484.
- Wickramasinghe, D. D., K. L. Oseen, S. W. Kotagama & R. J. Wassersug (2004): The terrestrial breeding biology of the ranid rock frog *Nannophrys ceylonensis*. *Behaviour* 141: 899-913.
- Wickramasinghe, L. J. M., D. R. Vidanapathirana, M. D. G. Rajeev, S. C. Ariyaratne, A. W. A. Chanaka, L. L. D. Priyantha, I. N. Bandara & N. Wickramasinghe (2013): Eight new species of *Pseudophilautus* (Amphibia: Anura: Rhacophoridae) from Sripada World Heritage Site (Peak Wilderness), a local amphibian hotspot in Sri Lanka. *Journal of Threatened Taxa* 5: 3789-3920.
- Wildenhues, M. J., A. Gawor, T. Q. Nguyen, T. T. Nguyen, A. Schmitz & T. Ziegler (2010): First description of larval and juvenile stages of *Rhacophorus maximus* Gunther, 1859 "1858" (Anura: Rhacophoridae) from Vietnam. *Revue suisse de Zoologie* 117: 679-696.

Wilkinson, J. A., T. Thin, K. S. Lwin & A. K. Shein (2005): A new species of *Rhacophorus* (Anura: Rhacophoridae) from Myanmar (Burma). *Proceedings-California Academy of Sciences* 56: 42-52.

Wogan, G. O. U. (2012): A new species of *Leptobrachium* from Myanmar (Anura: Megophryidae). *Zootaxa* 3415: 23-36.

Wolf, S. (2013): A case of male egg guarding behaviour in a stream-dwelling frog, *Mantidactylus* (Ochthomantis) sp (Anura: Mantellidae), from northeastern Madagascar. *Salamandra* 49: 45-47.


Zina, J., G. R. Silva, D. Loebmann & V. G. D. Orrico (2014): The recognition of *Dendropsophus minusculus* (Rivero, 1971) (Hylidae, Dendropsophini) as a highly polymorphic, multi-domain distributed species. *Brazilian Journal of Biology* 74: 146-153.

Webpages:

African amphibians Lifedesk	http://africanamphibians.lifedesks.org/
Amphibia Web	http://amphibiaweb.org/
Amphibians of Panama	http://biogeodb.stri.si.edu/amphibians/en/
Amphibiaweb	http://amphibiaweb.org/
Animal Diversity Web	http://animaldiversity.ummz.umich.edu
Dendroworld	http://www.dendroworld.co.uk
Frog Atlas	http://www.frogatlas.com.au/
Frog Forum	http://www.frogforum.net/content/
Frog.org	http://talkto.thefrog.org/
Frogs of Borneo	http://frogsofborneo.org/
Hepetofauna Costa Ricas	http://www.herpetologia.de
IUCN	http://www.iucnredlist.org/
Online field guide	http://online-field-guide.com/
Reptiles	http://www.reptilesmagazine.com/
Tropical Herping	http://www.tropicalherping.com/default.html
Wikipedia	https://en.wikipedia.org

Additional file 3. Clutch size classes

Clutch size was only available for a subset (345 out of 619) of species. An objective grouping by applying K-means clustering (*kmeans*, R package 'stats'; 1000 iterations) on the available data revealed the following clutch size classes (mean, range, sample size = species with this clutch size): A: 38.6, 4-98, n = 121; B: 160.5, 100-265, n = 68; C: 392.0, 290-549, n = 37; D: 712.0, 563-905, n = 24; E: 1201.6, 979-1652, n = 37; F: 2408.7, 1900-3320, n = 32; G: 4724.5, 3607-6701, n = 13; H: 10420.0, 8357-12940, n = 5; I: 21400.0, 17000-25000, n = 5; J: 38050.0, 36100-40000, n = 2. Ten classes were chosen to achieve a fine scale fractionation but at the same time at least a number of five records per class; except for the biggest size class. Species without information on clutch size were subsequently assigned to a single class based female (or male if female size was not available) body size, as the maximal number of eggs per clutch was significantly correlated with body size (females: $\rho = 0.58$, $p < 0.0001$, $n = 334$; males: $\rho = 0.58$, $p < 0.0001$, $n = 341$). Crucial was the minimal distance of the body size of a species lacking information to the mean body size of a particular clutch size class. 266 species were assigned based on female, 34 based on male body size. Calculation of clutch size classes and the assignment to a category was conducted independently of the reproductive mode, i.e. direct and indirect developing species together. This was permitted as data distribution of clutch size as well as the data distribution of clutch size related to female body size did not differ between the modes, i.e. data of direct developing species were within the range of clutch sizes of indirect developing species with similar body size (Kolmogorov-Smirnov test: clutch size: $D = 0.08$, $p = 0.23$; clutch size / female body size: $D = 0.07$, $p = 0.46$; N: direct developing species = 47, indirect developing species = 297; see Figure below).



Given are the maximum clutch size (log transformed) and the maximum female body size (snout vent length, svl) of 344 species. Direct developing species ($n = 47$) are marked with black, indirect developing species ($n = 297$) with white dots.

Additional file 4. Final Data set

Given is the species, the traits looked at and the forest dependency index (FDI) derived from the occurrence data. Dummy variables are highlighted in grey. See Table 1 for information on the respective traits.

Family	Genus	Species	Region of origin	Range size	Elevational range	SVL m	f	Adult habitat	Larval habitat	Clutch size	Clutch size class	Egg depos. site	Develop- ment	FDI
Aromobatidae	<i>Allobates</i>	<i>marchesianus</i>	America	1913449,48	1217,6	16,3	26,4	lit	len	14	A	ter	i	SD
Aromobatidae	<i>Anomaloglossus</i>	<i>stepheni</i>	America	24168,75	1217,6	17,0	18,0	lit	ter	6	A	ter	i	SD
Arthroleptidae	<i>Arthroleptis</i>	<i>adelphus</i>	Africa	471769,61	2003	29,0	33,1	lit	none	51	A	ter	d	SD
Arthroleptidae	<i>Arthroleptis</i>	<i>poecilonotus</i>	Africa	738639	1199	27,0	31,0	lit	none	29	A	ter	d	SD
Arthroleptidae	<i>Arthroleptis</i>	<i>stenodactylus</i>	Africa	2386661,02	1960	33,0	44,0	lit	none	80	A	ter	d	NR
Arthroleptidae	<i>Arthroleptis</i>	<i>sylvaticus</i>	Africa	1013495,74	1900	21,2	21,9	lit	none	20	A	ter	d	SD
Arthroleptidae	<i>Arthroleptis</i>	<i>variabilis</i>	Africa	1795153,2	1359	10,5	42,0	lit	none	68	A	ter	d	SD
Arthroleptidae	<i>Arthroleptis</i>	<i>wahlbergii</i>	Africa	50859,91	1000	18,0	22,0	lit	none	30	A	ter	d	NR
Arthroleptidae	<i>Arthroleptis</i>	<i>xenodactyloides</i>	Africa	1481,64	2000	17,0	27,0	lit	none	20	A	ter	d	NR
Arthroleptidae	<i>Cardioglossa</i>	<i>gracilis</i>	Africa	660912,32	1446	37,8	40,0	lit	lot	48	A	ter	i	SD
Arthroleptidae	<i>Leptopelis</i>	<i>brevirostris</i>	Africa	543127,2	1100	45,0	64,0	s-arb	none	NA	E	ter	i	SD
Arthroleptidae	<i>Leptopelis</i>	<i>calcaratus</i>	Africa	1884906,04	1500	42,0	57,0	s-arb	len	NA	D	ter	i	SD
Arthroleptidae	<i>Leptopelis</i>	<i>flavomaculatus</i>	Africa	805972,28	1217,6	50,0	70,0	arb	len	NA	E	ter	i	D
Arthroleptidae	<i>Leptopelis</i>	<i>mossambicus</i>	Africa	379280,3	1217,6	52,0	63,0	arb	len	NA	E	ter	i	D
Arthroleptidae	<i>Leptopelis</i>	<i>rufus</i>	Africa	236932,47	1757	55,0	87,0	arb	lot	NA	F	ter	i	SD
Arthroleptidae	<i>Trichobatrachus</i>	<i>robustus</i>	Africa	217523,72	1458	130,0	90,0	lit	lot	721	D	aqu	i	SD
Brevicipitidae	<i>Breviceps</i>	<i>adpersus</i>	Africa	2329163,99	1400	47,0	60,0	fos	none	95	A	ter	d	D
Brevicipitidae	<i>Breviceps</i>	<i>mossambicus</i>	Africa	1606225,44	1800	33,0	52,0	fos	ter	25	A	ter	i	I
Bufo	<i>Adenomus</i>	<i>kelaartii</i>	Asia	4928,03	1670	33,0	50,0	s-arb	lot	1000	E	aqu	i	NR
Bufo	<i>Amietophrynus</i>	<i>garmani</i>	Africa	1754477,81	2000	74,0	115,0	lit	len / lot	2000	F	aqu	i	NR
Bufo	<i>Amietophrynus</i>	<i>gutturalis</i>	Africa	5933446,92	1800	90,0	120,0	lit	len	25000	I	aqu	i	NR
Bufo	<i>Atelopus</i>	<i>pulcher</i>	America	9259,31	300	29,3	35,1	lit	lot	600	D	aqu	i	SD
Bufo	<i>Bufo</i>	<i>meghalayanus</i>	Asia	305,09	20	41,0	56,4	lit	len	NA	C	aqu	i	I
Bufo	<i>Duttaphrynus</i>	<i>kotagamai</i>	Asia	30,31	920	40,3	62,7	lit	lot	NA	E	aqu	i	D
Bufo	<i>Duttaphrynus</i>	<i>melanostictus</i>	Asia	600,82	1800	83,0	150,0	lit	len	40000	J	aqu	i	NR
Bufo	<i>Incilius</i>	<i>cristatus</i>	America	100,08	100	54,8	87,3	lit	lot	NA	F	aqu	i	D
Bufo	<i>Incilius</i>	<i>valliceps</i>	America	603359,39	1850	76,0	82,0	lit	len	NA	F	aqu	i	NR
Bufo	<i>Ingerophrynus</i>	<i>celebensis</i>	Asia	180051,98	1217,6	59,0	80,0	lit	len / lot	NA	F	aqu	i	NR
Bufo	<i>Mertensophryne</i>	<i>loveridgei</i>	Africa	56624,03	1217,6	35,0	38,0	lit	len	131	B	aqu	i	NR
Bufo	<i>Pedostibes</i>	<i>kempi</i>	Asia	6,17	1	30,0	56,4	arb	lot	NA	A	aqu	i	SD
Bufo	<i>Rhinella</i>	<i>granulosa</i>	America	10992958,86	1217,6	70,0	90,0	lit	len	900	D	aqu	i	NR
Bufo	<i>Rhinella</i>	<i>marina</i>	America	9689511,58	3000	187,0	287,0	lit	len / lot	36100	J	aqu	i	NR
Bufo	<i>Schismaderma</i>	<i>carens</i>	Africa	2955848,16	1217,6	88,0	92,0	lit	len	20000	I	aqu	i	NR
Centrolenidae	<i>Hyalinobatrachium</i>	<i>fleischmanni</i>	America	405988,89	1800	25,5	32,0	arb	lot	30	A	arb	i	SD
Centrolenidae	<i>Vitreorana</i>	<i>oyampiensis</i>	America	822656,57	810	21,0	24,0	arb	lot	20	A	arb	i	D
Ceratophryidae	<i>Ceratophrys</i>	<i>cornuta</i>	America	5285910,29	400	77,0	120,0	lit	len	2270	F	aqu	i	SD
Craugastoridae	<i>Craugastor</i>	<i>alfredi</i>	America	33650,37	600	36,0	45,0	lit	none	NA	B	ter	d	SD
Craugastoridae	<i>Craugastor</i>	<i>berkenbuschii</i>	America	56056,05	1500	69,5	93,0	s-aqu	none	NA	F	ter	d	SD
Craugastoridae	<i>Craugastor</i>	<i>decoratus</i>	America	5298,29	1378	45,0	60,0	s-arb	none	46	A	ter	d	NR
Craugastoridae	<i>Craugastor</i>	<i>fitzingeri</i>	America	273537,28	1520	35,0	52,5	s-arb	none	85	A	ter	d	NR
Craugastoridae	<i>Craugastor</i>	<i>mexicanus</i>	America	43635,86	1920	28,1	39,5	lit	none	NA	A	ter	d	SD
Craugastoridae	<i>Craugastor</i>	<i>pygmaeus</i>	America	16205,21	2000	18,0	18,0	lit	none	NA	A	ter	d	NR
Craugastoridae	<i>Craugastor</i>	<i>spatulatus</i>	America	85,95	1217,6	23,6	32,0	s-arb	none	NA	A	ter	d	SD
Craugastoridae	<i>Craugastor</i>	<i>uno</i>	America	321,31	1217,6	47,1	62,5	lit	none	NA	E	ter	d	SD

Craugastoridae	<i>Craugastor</i>	<i>vulcani</i>	America	1593,35	800	47,1	73,0	lit	none	NA	E	ter	d	D
Craugastoridae	<i>Pristimantis</i>	<i>achatinus</i>	America	184759	2330	36,0	46,0	s-arb	none	57	A	ter	d	NR
Craugastoridae	<i>Pristimantis</i>	<i>chloronatus</i>	America	7932,73	1500	30,3	36,9	s-arb	none	40	A	ter	d	NR
Craugastoridae	<i>Pristimantis</i>	<i>fenestratus</i>	America	3557857,79	1850	39,3	51,0	s-arb	none	48	A	ter	d	NR
Craugastoridae	<i>Pristimantis</i>	<i>festae</i>	America	1795153,2	1340	21,5	23,0	lit	none	NA	A	aqu	d	SD
Craugastoridae	<i>Pristimantis</i>	<i>zimmermanae</i>	America	39885,11	50	21,0	26,0	s-arb	none	NA	A	ter	d	SD
Dendrobatidae	<i>Allobates</i>	<i>femorialis</i>	America	5607495,67	1000	33,0	35,0	lit	len	43	A	ter	i	NR
Dicroglossidae	<i>Euphlyctis</i>	<i>cyanophlyctis</i>	Asia	4439313,99	2500	45,0	72,0	aqu	len / lot	844	D	aqu	i	NR
Dicroglossidae	<i>Euphlyctis</i>	<i>hexadactylus</i>	Asia	1477319,91	800	86,8	120,0	aqu	len	2000	F	aqu	i	NR
Dicroglossidae	<i>Fejervarya</i>	<i>limnocharis</i>	Asia	8896740,71	2000	52,0	67,0	s-aqu	len / lot	1400	E	aqu	i	NR
Dicroglossidae	<i>Fejervarya</i>	<i>cancrivora</i>	Asia	1342960,13	1300	70,0	82,0	aqu	len	2000	F	aqu	i	I
Dicroglossidae	<i>Hoplobatrachus</i>	<i>tigerinus</i>	Asia	3495177,78	1975	80,0	121,0	aqu	len / lot	2400	F	aqu	i	NR
Dicroglossidae	<i>Limnonectes</i>	<i>grunniens</i>	Asia	330324,38	1217,6	146,0	146,0	s-aqu	len / lot	NA	H	aqu	i	D
Dicroglossidae	<i>Limnonectes</i>	<i>laticeps</i>	Asia	341740,01	1450	47,0	46,0	s-aqu	lot	NA	B	aqu	i	SD
Dicroglossidae	<i>Limnonectes</i>	<i>malesianus</i>	Asia	140359,73	150	150,0	95,0	lit	lot	NA	I	aqu	i	SD
Dicroglossidae	<i>Limnonectes</i>	<i>modestus</i>	Asia	276147,2	1217,6	72,0	70,0	lit	lot	NA	E	ter	i	SD
Dicroglossidae	<i>Occidozyga</i>	<i>laevis</i>	Asia	507731,45	500	31,0	48,0	aqu	len / lot	140	B	aqu	i	NR
Dicroglossidae	<i>Sphaerotheca</i>	<i>breviceps</i>	Asia	3279685,84	1500	55,0	57,0	fos	len	75	A	aqu	i	NR
Dicroglossidae	<i>Zakerana</i>	<i>kirtisinghei</i>	Asia	8078,04	1430	39,0	44,0	lit	len	NA	B	aqu	i	I
Eleutherodactylidae	<i>Diasporus</i>	<i>diastema</i>	America	109484,13	1619	21,0	25,0	arb	none	10	A	arb	d	NR
Eleutherodactylidae	<i>Eleutherodactylus</i>	<i>cystignathoides</i>	America	24906,65	1200	14,0	26,0	lit	none	13	A	ter	d	SD
Eleutherodactylidae	<i>Eleutherodactylus</i>	<i>leprus</i>	America	2031,47	795	27,0	30,0	lit	none	NA	A	ter	d	NR
Hemisotidae	<i>Hemisus</i>	<i>guttatus</i>	Africa	51667,84	1000	51,0	80,0	fos	len	200	B	ter	i	NR
Hemisotidae	<i>Hemisus</i>	<i>marmoratum</i>	Africa	9850888,49	1850	45,0	55,0	fos	len	200	B	ter	i	SD
Hylidae	<i>Charadrahyla</i>	<i>taeniopus</i>	America	7538,77	1217,6	65,9	70,0	s-arb	len	NA	E	aqu	i	D
Hylidae	<i>Dendropsophus</i>	<i>brevifrons</i>	America	513088,37	1217,6	21,4	23,2	arb	len	115	B	arb	i	SD
Hylidae	<i>Dendropsophus</i>	<i>ebraccatus</i>	America	232873,74	1600	28,0	37,0	arb	len	155,5	B	arb	i	SD
Hylidae	<i>Dendropsophus</i>	<i>leucophyllata</i>	America	5525350,6	1217,6	36,0	44,0	s-arb	len	771	D	arb	i	NR
Hylidae	<i>Dendropsophus</i>	<i>marmorata</i>	America	4845441,2	1217,6	44,0	56,0	arb	len	979	E	aqu	i	NR
Hylidae	<i>Dendropsophus</i>	<i>microcephalus</i>	America	5570825,86	1300	25,0	32,0	s-arb	len	NA	A	aqu	i	NR
Hylidae	<i>Dendropsophus</i>	<i>minutus</i>	America	11045630,91	1217,6	23,0	25,0	arb	len	400	C	arb	i	NR
Hylidae	<i>Dendropsophus</i>	<i>nanus</i>	America	8158117,78	1500	20,0	23,0	s-arb	len	NA	A	aqu	i	I
Hylidae	<i>Dendropsophus</i>	<i>parviceps</i>	America	4478924,19	1217,6	21,9	26,1	s-arb	len	385	C	aqu	i	SD
Hylidae	<i>Ecnomiohyla</i>	<i>miotympanum</i>	America	22082,36	1382	38,4	51,0	arb	len / lot	NA	C	aqu	i	NR
Hylidae	<i>Ecnomiohyla</i>	<i>valancifer</i>	America	1241,71	880	77,7	82,0	arb	lot	NA	F	aqu	i	I
Hylidae	<i>Hypsiboas</i>	<i>boans</i>	America	6899721,81	1000	128,0	123,0	arb	len	4800	G	aqu	i	NR
Hylidae	<i>Hypsiboas</i>	<i>calcaratus</i>	America	4611334,93	600	44,0	54,0	arb	len / lot	1144	E	aqu	i	SD
Hylidae	<i>Hypsiboas</i>	<i>cinerascens</i>	America	5614933,83	500	35,0	37,0	arb	len	400	C	aqu	i	NR
Hylidae	<i>Hypsiboas</i>	<i>geographicus</i>	America	5190041,24	700	55,0	83,0	arb	len / lot	2797	F	aqu	i	SD
Hylidae	<i>Hypsiboas</i>	<i>lanciformis</i>	America	10419166,83	1217,6	80,0	94,0	arb	len	2500	F	aqu	i	I
Hylidae	<i>Hypsiboas</i>	<i>punctatus</i>	America	4621749,79	1400	30,0	39,0	arb	len	325	C	aqu	i	NR
Hylidae	<i>Megastomatohyla</i>	<i>mixomaculata</i>	America	973,43	600	30,0	35,0	s-arb	lot	NA	A	aqu	i	D
Hylidae	<i>Osteocephalus</i>	<i>buckleyi</i>	America	4477069,23	1217,6	50,0	69,0	arb	lot	1000	E	aqu	i	D
Hylidae	<i>Osteocephalus</i>	<i>lepieurii</i>	America	3812252,54	860	48,1	64,0	arb	len	2882	F	aqu	i	SD
Hylidae	<i>Osteocephalus</i>	<i>taurinus</i>	America	6116928,89	1217,6	92,0	101,0	arb	len	2000	F	aqu	i	NR
Hylidae	<i>Phyllomedusa</i>	<i>bicolor</i>	America	5230330,94	800	103,0	119,0	arb	len	1202	E	arb	i	SD
Hylidae	<i>Phyllomedusa</i>	<i>tarsius</i>	America	2694953,8	450	90,0	110,0	s-arb	len	549	C	arb	i	NR
Hylidae	<i>Phyllomedusa</i>	<i>tomopterna</i>	America	5475272	500	54,0	62,0	arb	len	89	A	arb	i	SD
Hylidae	<i>Phyllomedusa</i>	<i>vaillantii</i>	America	4991230,63	450	65,0	83,0	arb	len	645	D	arb	i	NR
Hylidae	<i>Plectrohyla</i>	<i>arborescandens</i>	America	3286,81	1500	38,0	52,0	arb	lot	NA	D	aqu	i	D

Hylidae	<i>Plectrohyla</i>	<i>pentheter</i>	America	2297,22	720	46,0	57,8	s-arb	lot	NA	D	aqu	i	SD
Hylidae	<i>Ptychohyla</i>	<i>leonhardschultzei</i>	America	1123,88	1300	34,9	30,5	lit	lot	60	A	aqu	i	SD
Hylidae	<i>Scinax</i>	<i>cruentommus</i>	America	2460973,09	1217,6	27,0	32,0	arb	len	1200	E	aqu	i	NR
Hylidae	<i>Scinax</i>	<i>garbei</i>	America	3703694,25	1217,6	46,0	48,0	arb	len	905	D	aqu	i	I
Hylidae	<i>Scinax</i>	<i>ruber</i>	America	8136346	2600	37,0	45,0	arb	len	1105	E	aqu	i	NR
Hylidae	<i>Scinax</i>	<i>staufferi</i>	America	569459,34	1530	29,0	32,0	arb	len	NA	A	aqu	i	SD
Hylidae	<i>Smilisca</i>	<i>baudinii</i>	America	1050352,67	1925	76,0	90,0	arb	len	3320	F	aqu	i	NR
Hylidae	<i>Tlalocohyla</i>	<i>loquax</i>	America	348680,87	1585	35,0	47,0	s-arb	len	250	B	aqu	i	NR
Hylidae	<i>Tlalocohyla</i>	<i>picta</i>	America	313559,87	1300	21,4	22,1	s-arb	len	NA	A	aqu	i	SD
Hylidae	<i>Trachycephalus</i>	<i>resinifictrix</i>	America	4649150,62	450	77,0	88,0	arb	phy	2500	F	arb	i	SD
Hyperoliidae	<i>Afrixalus</i>	<i>fulvovittatus</i>	Africa	94244,89	1685	27,0	28,0	s-arb	len	100	B	arb	i	SD
Hyperoliidae	<i>Afrixalus</i>	<i>laevis</i>	Africa	1207641,32	1217,6	23,0	25,0	s-arb	lot	39	A	arb	i	SD
Hyperoliidae	<i>Hyperolius</i>	<i>argus</i>	Africa	708667,09	300	34,0	37,0	s-arb	len	NA	A	arb	i	D
Hyperoliidae	<i>Hyperolius</i>	<i>armoratus</i>	Africa	1212234,73	1600	34,0	34,0	s-arb	len	100	B	aqu	i	NR
Hyperoliidae	<i>Hyperolius</i>	<i>mitchelli</i>	Africa	808554,63	1217,6	27,0	32,0	s-arb	len	290	C	arb	i	I
Hyperoliidae	<i>Hyperolius</i>	<i>nasutus</i>	Africa	1828824,27	1217,6	24,0	25,0	s-arb	len / lot	NA	A	aqu	i	SD
Hyperoliidae	<i>Hyperolius</i>	<i>pusillus</i>	Africa	798478,66	800	20,0	25,0	s-arb	len	200	B	aqu	i	D
Hyperoliidae	<i>Hyperolius</i>	<i>semidiscus</i>	Africa	141664,64	1217,6	25,0	35,0	s-arb	len	17	A	aqu	i	D
Hyperoliidae	<i>Hyperolius</i>	<i>tuberilinguis</i>	Africa	3091192,54	1000	33,0	35,0	s-arb	len / lot	NA	A	arb	i	SD
Hyperoliidae	<i>Kassina</i>	<i>senegalensis</i>	Africa	12217676,34	2000	40,0	40,0	lit	len	10	A	aqu	i	D
Leiuperidae	<i>Physalaemus</i>	<i>cuvieri</i>	America	7282718,66	1217,6	28,0	31,0	lit	len	2465	F	aqu	i	NR
Leptodactylidae	<i>Adenomera</i>	<i>andreae</i>	America	5976957,43	400	20,0	27,0	lit	ter	15	A	ter	i	NR
Leptodactylidae	<i>Leptodactylus</i>	<i>fragilis</i>	America	1433339,34	1700	36,0	40,0	lit	len	250	B	ter	i	NR
Leptodactylidae	<i>Leptodactylus</i>	<i>knudseni</i>	America	5236498,65	1800	170,0	154,0	lit	len	1000	E	ter	i	NR
Leptodactylidae	<i>Leptodactylus</i>	<i>leptodactylodes</i>	America	10932822,52	685	47,9	56,2	lit	len	1053	E	aqu	i	SD
Leptodactylidae	<i>Leptodactylus</i>	<i>mystaceus</i>	America	7133840,42	1217,6	50,0	60,0	lit	len	425	C	ter	i	NR
Leptodactylidae	<i>Leptodactylus</i>	<i>mystacinus</i>	America	5187450,08	1800	65,0	67,0	lit	len	627	D	ter	i	NR
Leptodactylidae	<i>Leptodactylus</i>	<i>pentadactylus</i>	America	4505396,51	1195	177,0	185,0	lit	len	1000	E	ter	i	SD
Leptodactylidae	<i>Leptodactylus</i>	<i>petersii</i>	America	6056228,71	1217,6	40,0	45,0	lit	len	563	D	ter	i	NR
Leptodactylidae	<i>Leptodactylus</i>	<i>rhodomystax</i>	America	5152273,6	500	72,0	85,0	lit	len	290	C	ter	i	NR
Leptodactylidae	<i>Leptodactylus</i>	<i>riveroi</i>	America	836714,79	360	58,0	74,0	lit	len	1000	E	aqu	i	SD
Leptodactylidae	<i>Leptodactylus</i>	<i>stenodema</i>	America	4406321,41	1217,6	94,0	96,0	lit	len	NA	I	s-ter	i	SD
Leptodactylidae	<i>Lithodytes</i>	<i>lineatus</i>	America	442,1	1800	43,0	56,0	lit	len	260	B	ter	i	NR
Mantellidae	<i>Aglyptodactylus</i>	<i>madagascariensis</i>	Madagascar	152848,58	2000	41,0	53,0	lit	len	2865	F	aqu	i	SD
Mantellidae	<i>Blommersia</i>	<i>blommersae</i>	Madagascar	17718,31	400	21,0	20,0	s-arb	len	NA	A	arb	i	NR
Mantellidae	<i>Blommersia</i>	<i>grandisonae</i>	Madagascar	64754,59	800	23,0	20,0	s-arb	lot	NA	A	arb	i	SD
Mantellidae	<i>Boophis</i>	<i>boehmei</i>	Madagascar	60227,92	1100	30,0	41,0	s-arb	lot	NA	A	aqu	i	I
Mantellidae	<i>Boophis</i>	<i>goudotii</i>	Madagascar	80548,07	1300	70,0	87,0	s-arb	lot	1000	E	aqu	i	NR
Mantellidae	<i>Boophis</i>	<i>lichenoides</i>	Madagascar	57627,21	850	43,0	56,4	s-arb	len	NA	C	aqu	i	SD
Mantellidae	<i>Boophis</i>	<i>luteus</i>	Madagascar	37505,21	1500	60,0	40,0	s-arb	lot	200	B	aqu	i	SD
Mantellidae	<i>Boophis</i>	<i>madagascariensis</i>	Madagascar	91319,12	1700	62,0	75,0	s-arb	len / lot	405	C	aqu	i	SD
Mantellidae	<i>Boophis</i>	<i>marojezensis</i>	Madagascar	41872,73	700	27,0	33,0	s-arb	lot	NA	A	aqu	i	SD
Mantellidae	<i>Boophis</i>	<i>picturatus</i>	Madagascar	28014,9	150	33,0	56,4	s-arb	lot	NA	A	aqu	i	SD
Mantellidae	<i>Boophis</i>	<i>pyrrhus</i>	Madagascar	30001,38	465	32,0	37,0	s-arb	len	NA	A	aqu	i	SD
Mantellidae	<i>Boophis</i>	<i>rappiodes</i>	Madagascar	31797,19	600	25,0	34,0	s-arb	lot	300	C	aqu	i	SD
Mantellidae	<i>Boophis</i>	<i>reticulatus</i>	Madagascar	62346,55	850	35,0	44,0	s-arb	len	NA	B	aqu	i	SD
Mantellidae	<i>Boophis</i>	<i>rufioculis</i>	Madagascar	15852,15	300	35,0	47,0	s-arb	lot	NA	B	aqu	i	I
Mantellidae	<i>Gephyromantis</i>	<i>boulengeri</i>	Madagascar	27478,03	1200	30,0	56,4	lit	none	NA	A	ter	d	SD
Mantellidae	<i>Gephyromantis</i>	<i>cornutus</i>	Madagascar	9606,04	350	40,0	39,0	s-arb	len	NA	A	aqu	i	SD
Mantellidae	<i>Gephyromantis</i>	<i>eiselti</i>	Madagascar	1993,95	400	23,0	35,0	lit	none	4	A	ter	d	I

Mantellidae	<i>Gephyromantis</i>	<i>luteus</i>	Madagascar	64398,49	700	43,0	47,0	s-arb	none	NA	B	ter	d	SD
Mantellidae	<i>Gephyromantis</i>	<i>malagasius</i>	Madagascar	62053,09	1500	23,0	26,0	lit	none	NA	A	ter	d	SD
Mantellidae	<i>Gephyromantis</i>	<i>redimitus</i>	Madagascar	52053,49	850	53,0	56,4	s-arb	len	NA	E	aqu	i	SD
Mantellidae	<i>Guibemantis</i>	<i>bicalcaratus</i>	Madagascar	115691,97	1200	26,0	29,0	arb	phy	54,8	A	arb	i	SD
Mantellidae	<i>Guibemantis</i>	<i>depressiceps</i>	Madagascar	134337,42	1200	45,0	35,0	s-arb	len	70	A	arb	i	I
Mantellidae	<i>Guibemantis</i>	<i>flavobrunneus</i>	Madagascar	62478,52	1000	33,0	38,0	arb	len	NA	A	aqu	i	SD
Mantellidae	<i>Guibemantis</i>	<i>kathrinae</i>	Madagascar	450,46	340	59,0	56,4	arb	len	NA	E	arb	i	SD
Mantellidae	<i>Guibemantis</i>	<i>liber</i>	Madagascar	143658,64	2050	29,0	28,0	arb	len	90	A	arb	i	NR
Mantellidae	<i>Guibemantis</i>	<i>pulcher</i>	Madagascar	62885,3	1400	25,0	28,0	arb	phy	NA	A	arb	i	SD
Mantellidae	<i>Mantella</i>	<i>baroni</i>	Madagascar	32886,41	600	29,0	30,0	lit	len	64	A	ter	i	SD
Mantellidae	<i>Mantella</i>	<i>pulchra</i>	Madagascar	16161,75	650	26,0	28,0	lit	lot	61	A	ter	i	SD
Mantellidae	<i>Mantidactylus</i>	<i>aerumnalis</i>	Madagascar	36593,35	500	27,0	31,0	lit	lot	NA	A	ter	i	SD
Mantellidae	<i>Mantidactylus</i>	<i>albofrenatus</i>	Madagascar	1848,89	50	23,0	27,0	lit	lot	NA	A	ter	i	SD
Mantellidae	<i>Mantidactylus</i>	<i>argenteus</i>	Madagascar	28417,71	600	27,0	31,0	lit	lot	12	A	arb	i	SD
Mantellidae	<i>Mantidactylus</i>	<i>betsileanus</i>	Madagascar	87389,03	1500	28,0	37,0	lit	lot	7	A	s-ter	i	NR
Mantellidae	<i>Mantidactylus</i>	<i>biporus</i>	Madagascar	88413,28	1600	27,0	34,0	s-aqu	len	NA	A	ter	i	SD
Mantellidae	<i>Mantidactylus</i>	<i>femoralis</i>	Madagascar	191431,49	2500	37,0	55,0	s-aqu	lot	76	A	ter	i	SD
Mantellidae	<i>Mantidactylus</i>	<i>grandidieri</i>	Madagascar	76192,23	1500	180,0	56,4	s-aqu	lot	NA	J	aqu	i	SD
Mantellidae	<i>Mantidactylus</i>	<i>mocquardi</i>	Madagascar	126158,93	2500	55,0	67,0	lit	lot	NA	E	ter	i	SD
Mantellidae	<i>Mantidactylus</i>	<i>opiparis</i>	Madagascar	74905,08	900	26,0	33,0	lit	lot	89	A	aqu	i	SD
Mantellidae	<i>Mantidactylus</i>	<i>tricinctus</i>	Madagascar	2229,22	400	19,0	20,0	s-aqu	len / lot	NA	A	aqu	i	SD
Mantellidae	<i>Spinomantis</i>	<i>peraccae</i>	Madagascar	100196,03	1500	41,0	42,0	arb	lot	NA	D	arb	i	SD
Megophryidae	<i>Leptobranchium</i>	<i>smithi</i>	Asia	206534,7	1290	68,2	78,0	lit	lot	NA	F	aqu	i	SD
Microhylidae	<i>Anodonthyla</i>	<i>boulengerii</i>	Madagascar	80806,46	1300	22,0	21,0	s-arb	phy	NA	A	arb	i	SD
Microhylidae	<i>Chiasmocleis</i>	<i>albopunctata</i>	America	2548381,93	1400	32,0	38,0	lit	len	NA	A	aqu	i	SD
Microhylidae	<i>Chiasmocleis</i>	<i>shudikarensis</i>	America	2910745,4	200	25,0	28,0	lit	len	566	D	aqu	i	SD
Microhylidae	<i>Ctenophryne</i>	<i>geayi</i>	America	5940721,75	600	41,7	49,3	fos	len	600	D	ter	i	D
Microhylidae	<i>Gastrophryne</i>	<i>elegans</i>	America	10959404,32	952	25,8	28,9	lit	len	NA	A	aqu	i	NR
Microhylidae	<i>Hypopachus</i>	<i>ustus</i>	America	211031,15	1000	30,6	31,4	lit	len	NA	A	aqu	i	SD
Microhylidae	<i>Kalophrynus</i>	<i>pleurostigma</i>	Asia	754159,24	2200	55,0	60,0	lit	len	6	A	aqu	i	SD
Microhylidae	<i>Kaloula</i>	<i>baleata</i>	Asia	860868,31	1525	61,0	67,0	fos	len	160	B	aqu	i	NR
Microhylidae	<i>Kaloula</i>	<i>pulchra</i>	Asia	2438006,6	750	70,0	75,0	fos	len	NA	E	aqu	i	NR
Microhylidae	<i>Microhyla</i>	<i>berdmorei</i>	Asia	1622527,08	1217,6	28,0	45,0	lit	len	NA	B	aqu	i	NR
Microhylidae	<i>Microhyla</i>	<i>ornata</i>	Asia	3446277,95	2000	27,0	32,0	lit	len	1200	E	aqu	i	NR
Microhylidae	<i>Microhyla</i>	<i>rubra</i>	Asia	1178745,31	700	27,5	29,5	lit	len	NA	A	aqu	i	NR
Microhylidae	<i>Paradoxophyla</i>	<i>palmata</i>	Madagascar	31237,38	950	22,1	25,6	fos	len	408	C	aqu	i	SD
Microhylidae	<i>Platypelis</i>	<i>barbouri</i>	Madagascar	37250,33	1100	23,0	56,4	s-arb	phy	NA	A	aqu	i	I
Microhylidae	<i>Platypelis</i>	<i>grandis</i>	Madagascar	129138,45	1217,6	88,0	61,0	s-arb	len	100	B	aqu	i	SD
Microhylidae	<i>Platypelis</i>	<i>pollicaris</i>	Madagascar	11501,66	1200	28,0	56,4	s-arb	phy	NA	A	aqu	i	SD
Microhylidae	<i>Platypelis</i>	<i>tuberifera</i>	Madagascar	62693,76	800	40,0	56,4	s-arb	phy	NA	C	aqu	i	SD
Microhylidae	<i>Plethodontohyla</i>	<i>inguinalis</i>	Madagascar	71194,35	700	100,0	63,0	s-arb	phy	NA	E	arb	i	SD
Microhylidae	<i>Plethodontohyla</i>	<i>notosticta</i>	Madagascar	102688,81	1200	35,0	38,5	s-arb	phy	120	B	aqu	i	SD
Microhylidae	<i>Rhombophryne</i>	<i>alluaudi</i>	Madagascar	99859,19	1400	60,0	56,4	fos	len	NA	E	ter	i	SD
Microhylidae	<i>Stumpffia</i>	<i>tetradactyla</i>	Madagascar	595,36	100	15,0	56,4	lit	ter	NA	A	ter	i	SD
Microhylidae	<i>Synapturanus</i>	<i>salseri</i>	America	1318002,07	200	30,0	35,0	fos	ter	8	A	ter	d	SD
Petropedetidae	<i>Conraua</i>	<i>crassipes</i>	Africa	555260,86	1758	81,0	76,0	s-aqu	lot	NA	F	aqu	i	SD
Petropedetidae	<i>Conraua</i>	<i>robusta</i>	Africa	12360,4	650	140,0	122,0	s-aqu	lot	52	A	ter	i	SD
Phrynobatrachidae	<i>Phrynobatrachus</i>	<i>acridoides</i>	Africa	1756215,67	700	27,0	25,0	s-aqu	len	105	B	aqu	i	I
Phrynobatrachidae	<i>Phrynobatrachus</i>	<i>africanus</i>	Africa	713611,07	1758	28,0	30,0	lit	len	NA	A	aqu	i	SD
Phrynobatrachidae	<i>Phrynobatrachus</i>	<i>mababiensis</i>	Africa	4355399,29	2410	19,0	22,0	lit	len	775	D	aqu	i	NR

Phrynobatrachidae	<i>Phrynobatrachus</i>	<i>natalensis</i>	Africa	10456437,35	2200	30,0	31,0	lit	len	1652	E	aqu	i	NR
Pipidae	<i>Pipa</i>	<i>arrabali</i>	America	1038533,62	860	42,0	55,0	aqu	skin	16	A	skin	d	D
Pipidae	<i>Xenopus</i>	<i>laevis</i>	Africa	4599862,56	3000	97,5	147,0	aqu	len	17000	I	aqu	i	D
Ptychadenidae	<i>Ptychadena</i>	<i>anchietae</i>	Africa	6641710,13	1780	51,0	62,0	lit	len	300	C	aqu	i	NR
Ptychadenidae	<i>Ptychadena</i>	<i>mascareniensis</i>	Africa	590831,48	2000	57,0	68,0	lit	len	1079	E	aqu	i	NR
Ptychadenidae	<i>Ptychadena</i>	<i>mossambica</i>	Africa	2867084,3	1000	35,0	44,0	lit	len	315	C	aqu	i	D
Ptychadenidae	<i>Ptychadena</i>	<i>oxyrhynchus</i>	Africa	8160886,06	2000	53,0	64,0	lit	len	5018	G	aqu	i	SD
Pyxicephalidae	<i>Amietia</i>	<i>angolensis</i>	Africa	4274988,92	3100	81,0	90,0	s-aqu	len / lot	500	C	aqu	i	I
Pyxicephalidae	<i>Pyxicephalus</i>	<i>edulis</i>	Africa	2933976,25	1500	120,0	120,0	fos	len	4000	G	aqu	i	SD
Pyxicephalidae	<i>Tomopterna</i>	<i>cryptotis</i>	Africa	10857651,05	811	45,0	58,0	fos	len	3000	F	aqu	i	D
Pyxicephalidae	<i>Tomopterna</i>	<i>krugerensis</i>	Africa	1209670,06	1500	45,0	46,0	fos	len	5000	G	aqu	i	D
Pyxicephalidae	<i>Tomopterna</i>	<i>natalensis</i>	Africa	494825,84	2000	39,0	44,0	fos	len / lot	NA	B	aqu	i	SD
Ranidae	<i>Clinotarsus</i>	<i>alticola</i>	Asia	152446,47	1217,6	53,0	56,4	s-arb	lot	NA	E	aqu	i	SD
Ranidae	<i>Hylarana</i>	<i>albolabris</i>	Africa	3669770,03	1939	56,0	70,0	s-arb	len	636	D	aqu	i	SD
Ranidae	<i>Hylarana</i>	<i>aurantiaca</i>	Asia	12416,56	1395	56,0	63,0	s-arb	lot	NA	E	aqu	i	NR
Ranidae	<i>Hylarana</i>	<i>celebensis</i>	Asia	30938,04	1217,6	46,0	56,4	lit	len / lot	NA	D	aqu	i	NR
Ranidae	<i>Hylarana</i>	<i>chalconota</i>	Asia	246684,24	1217,6	45,0	61,0	s-arb	len / lot	3200	F	aqu	i	SD
Ranidae	<i>Hylarana</i>	<i>erythraea</i>	Asia	2091033,97	1200	48,0	75,0	lit	len	1900	F	aqu	i	NR
Ranidae	<i>Hylarana</i>	<i>temporalis</i>	Asia	43280,93	1770	56,1	78,0	lit	lot	1200	E	aqu	i	SD
Ranidae	<i>Lankanectes</i>	<i>corrugatus</i>	Asia	9692,15	1495	35,0	71,0	aqu	len	NA	E	aqu	i	NR
Ranidae	<i>Lithobates</i>	<i>berlandieri</i>	America	1463513,93	3002	112,0	56,4	aqu	len / lot	3000	F	aqu	i	NR
Ranidae	<i>Lithobates</i>	<i>vaillanti</i>	America	507080,67	1097	94,0	125,0	s-aqu	len / lot	1000	E	aqu	i	NR
Ranidae	<i>Nannophrys</i>	<i>ceylonensis</i>	Asia	5598,28	1140	42,5	52,5	aqu	semi-aquatic	43,2	A	ter	i	NR
Ranidae	<i>Pterorana</i>	<i>khare</i>	Asia	5211,62	1400	62,0	56,4	lit	lot	900	D	aqu	i	SD
Ranixalidae	<i>Indirana</i>	<i>beddomii</i>	Asia	50217,45	1800	50,0	60,0	lit	semi-aquatic	NA	E	aqu	i	NR
Ranixalidae	<i>Indirana</i>	<i>leithii</i>	Asia	16246,03	800	38,0	56,4	lit	semi-aquatic	NA	B	s-ter	i	SD
Rhacophoridae	<i>Chiromantis</i>	<i>vittatus</i>	Asia	1279274,39	1500	35,0	32,0	arb	len	NA	A	arb	i	NR
Rhacophoridae	<i>Chiromantis</i>	<i>xerampelina</i>	Africa	2780951,55	1200	75,0	90,0	arb	len	1226	E	arb	i	NR
Rhacophoridae	<i>Nyctixalus</i>	<i>pictus</i>	Asia	205915,69	1700	33,0	38,0	s-arb	phy	15	A	arb	i	SD
Rhacophoridae	<i>Philautus</i>	<i>namdaphaensis</i>	Asia	928,27	200	27,0	56,4	s-arb	len	12	A	ter	d	D
Rhacophoridae	<i>Polypedates</i>	<i>cruciger</i>	Asia	16005,14	1510	59,8	90,0	arb	len	300	C	arb	i	I
Rhacophoridae	<i>Polypedates</i>	<i>leucomystax</i>	Asia	3926220,27	1500	50,0	75,0	arb	len	900	D	arb	i	NR
Rhacophoridae	<i>Polypedates</i>	<i>longinasus</i>	Asia	97,9	1150	47,0	60,0	s-arb	len	42	A	arb	i	D
Rhacophoridae	<i>Pseudophilautus</i>	<i>abundus</i>	Asia	402,43	72	31,0	37,0	s-arb	none	155	B	ter	d	D
Rhacophoridae	<i>Pseudophilautus</i>	<i>auratus</i>	Asia	111,54	757	23,3	26,4	s-arb	none	155	B	ter	d	D
Rhacophoridae	<i>Pseudophilautus</i>	<i>cavirostris</i>	Asia	197,21	900	47,1	48,9	s-arb	none	155	B	ter	d	D
Rhacophoridae	<i>Pseudophilautus</i>	<i>cuspis</i>	Asia	130,28	510	20,3	228,9	s-arb	none	155	B	ter	d	D
Rhacophoridae	<i>Pseudophilautus</i>	<i>folicola</i>	Asia	21,62	1240	23,1	29,4	s-arb	none	155	B	ter	d	NR
Rhacophoridae	<i>Pseudophilautus</i>	<i>hoipolloi</i>	Asia	501,92	685	27,6	28,6	s-arb	none	155	B	ter	d	I
Rhacophoridae	<i>Pseudophilautus</i>	<i>mittermeieri</i>	Asia	14,67	90	18,4	56,4	s-arb	none	155	B	ter	d	D
Rhacophoridae	<i>Pseudophilautus</i>	<i>popularis</i>	Asia	3743,31	1067	21,3	24,7	s-arb	none	155	B	ter	d	I
Rhacophoridae	<i>Pseudophilautus</i>	<i>reticulatus</i>	Asia	96,12	1340	44,1	61,1	arb	none	155	B	ter	d	D
Rhacophoridae	<i>Pseudophilautus</i>	<i>silvaticus</i>	Asia	106,27	760	25,3	31,3	s-arb	none	155	B	ter	d	D
Rhacophoridae	<i>Pseudophilautus</i>	<i>sordidus</i>	Asia	6404,69	1519	36,8	39,4	s-arb	none	155	B	ter	d	NR
Rhacophoridae	<i>Pseudophilautus</i>	<i>stictomerus</i>	Asia	230,04	455	23,2	36,0	s-arb	none	155	B	ter	d	NR
Rhacophoridae	<i>Raorchestes</i>	<i>parvulus</i>	Asia	426431,47	1350	19,6	21,0	s-arb	none	NA	A	aqu	d	NR
Rhacophoridae	<i>Rhacophorus</i>	<i>georgii</i>	Asia	4493,61	1217,6	47,1	70,0	arb	phy	108	B	arb	i	D
Rhacophoridae	<i>Rhacophorus</i>	<i>maximus</i>	Asia	328678,53	1500	114,3	56,4	arb	len	NA	G	arb	i	D
Rhacophoridae	<i>Rhacophorus</i>	<i>monticola</i>	Asia	23126,65	1217,6	36,7	39,9	lit	len	NA	B	arb	i	SD

Additional file 5. Affiliation to different anuran families

Given are number and relative frequency (in color code) of species belonging to a particular family per forest dependency index (higher frequency with darker color, see legend in table); forest dependency index: D = dependent (n = 33), SD = slightly dependent (n = 108), NR = non-responding (n = 83), I = forest independent (n = 19).

Family	Forest dependency Index			
	D	SD	NR	I
Aromobatidae	0	2	0	0
Arthroleptidae	2	9	3	0
Brevicipitidae	1	0	0	1
Bufo	2	2	10	1
Centrolenidae	1	1	0	0
Ceratophryidae	0	1	0	0
Craugastoridae	1	7	6	0
Dendrobatidae	0	0	1	0
Dicoglossidae	1	3	6	2
Eleutherodactylidae	0	1	2	0
Hemisotidae	0	1	1	0
Hylidae	4	13	15	4
Hyperoliidae	4	4	1	1
Leiuperidae	0	0	1	0
Leptodactylidae	0	4	8	0
Mantellidae	0	31	4	4
Megophryidae	0	1	0	0
Microhylidae	1	14	6	1
Petropedetidae	0	2	0	0
Phrynobatrachidae	0	1	2	1
Pipidae	2	0	0	0
Ptychadenidae	1	1	2	0
Pyxicephalidae	2	2	0	1
Ranidae	0	5	7	0
Ranixalidae	0	1	1	0
Rhacophoridae	11	2	7	3

0
< 10%
< 30%
> 30% (33%)