

NORDIC JOURNAL OF BOTANY

Research

- 0 **Determinants of alpha and beta vascular plant diversity in** 61
Mediterranean island systems: the Ionian islands, Greece

5 Anna-Thalassini Valli, Konstantinos Kougioumoutzis, Eleni Iliadou, Maria Panitsa and Panayiotis Trigas 65

10 A.-T. Valli (<http://orcid.org/0000-0002-2085-1174>) (thalassinivalli@yahoo.gr), K. Kougioumoutzis and P. Trigas (<https://orcid.org/0000-0001-9557-7723>), Laboratory of Systematic Botany, Faculty of Crop Science, Agricultural Univ. of Athens, Athens, Greece. – E. Iliadou and M. Panitsa, Division of 75
Plant Biology, Faculty of Biology, Univ. of Patras, Patras, Greece.

15 **Nordic Journal of Botany**
2018: e02156

20 doi: 10.1111/njb.02156

Subject Editor: Rob Lewis
Editor-in-Chief: Torbjörn Tyler
25 Accepted 19 November 2018

The Ionian archipelago is the second largest Greek archipelago after the Aegean, but the factors driving plant species diversity in the Ionian islands are still barely known. We used stepwise multiple regressions to investigate the factors affecting plant species diversity in 17 Ionian islands. Generalized dissimilarity modelling was applied to examine variation in the magnitude and rate of species turnover along environmental gradients, as well as to assess the relative importance of geographical and climatic factors in explaining species turnover. The values of the residuals from the ISAR log₁₀-transformed models of native and endemic taxa were used as a measure of island floristic diversity. Area was confirmed to be the most powerful single explanatory predictor of all diversity metrics. Mean annual precipitation and temperature, as well as shortest distance to the nearest island are also significant predictors of vascular plant diversity. The island of Kalamos constitutes an important plant diversity hotspot in the Ionian archipelago. The recent formation of the islands, the close proximity to the mainland source and the relatively low dispersal filtering of the Ionian archipelago has resulted in islands with a flora principally comprising common species and a low proportion of endemics. Small islands keep a key role in conservation of plant priority sites.

Keywords: species richness, endemism, island biogeography

Introduction

Islands contribute disproportionately to global biodiversity, since they comprise just 3.5% of earth's land area but host 15–20% of all terrestrial species (Whittaker et al. 2017). Thus, islands are of key interest for studies and experimental research in ecology, biogeography and evolution (Denslow 2001). They have additional research advantages compared to mainland areas, as they are comparatively small, have distinct boundaries, their biotas are usually less complex than those of the adjacent mainland, and ecological and evolutionary processes are relatively easier to detect on islands due to their geographical isolation (Whittaker and Fernández-Palacios 2007). Islands also play a major role in the contemporary extinction crisis, as more than



0 60% of the documented terrestrial species extinctions since
1500 CE have been island endemics (Johnson et al. 2017,
Whittaker et al. 2017). Consequently, islands are particu-
larly important for the conservation of global biodiversity
(Caujapé-Castells et al. 2010).

5 The Mediterranean basin is one of the 35 major global
terrestrial biodiversity hotspots (Médail and Myers 2004).
The Mediterranean islands, which are mainly of continen-
tal origin, contain numerous endemic and range-restricted
10 plants (Thompson et al. 2005, Vogiatzakis et al. 2008,
Médail 2013). Their high plant diversity reflects their
palaeogeographical history (Médail and Quézel 1999). Most
Mediterranean islands belong to the two Greek archipelagos,
with ca 7600 islands and islets in the Aegean archipelago and
15 ca 300 islands and islets in the Ionian (Médail 2017).

The Ionian archipelago is located at the eastern half
of the Mediterranean basin, in western Greece (Fig. 1).
The size and elevation of individual islands range from
0.015 to 734 km² and from 12 to 1628 m a.s.l., respec-
tively. They form a distinct phytogeographical region, (IoI
– sensu Dimopoulos et al. 2013) that hosts 2027 plant
taxa, 1827 of which are native and 89 are Greek endemics
(Dimopoulos et al. 2013, 2016, Flora Ionica Working
25 Group 2016–). In the Ionian archipelago, 21 sites have been
included in the Natura 2000 network of protected areas,
underlying the need to protect and preserve the biodiver-
sity in this region. The ecological importance of the Ionian
islands is also highlighted by the existence of two National

61 Parks (the Ainos National Park and the Zakynthos National
Marine Park), while Echinades islets (islands no. 1–4 in
Fig. 1) are part of a National Park that also includes main-
land areas. The Ionian islands belong to the Mediterranean
65 climatic belt, while along this area some of the highest cen-
tral Mediterranean precipitation heights have been recorded
(Kolios and Kalimeris 2017).

The origin and distribution patterns of the Greek flora have
previously been related to palaeogeographical patterns (Turrill
70 1929, Rechinger 1965). The Ionian islands play a major role
in the geodynamic framework of the central Mediterranean,
constituting a multiple junction region (Accordi et al. 2014),
where three tectonic plates meet (African plate, Eurasian plate
and Adriatic micro-plate). The palaeogeographical history of
75 the Ionian islands is rather recent, as most islands became
separated from the adjacent mainland during the Pleistocene
or even later (Perissoratis and Conispoliatis 2003, Triantis
and Mylonas 2009).

During Pleistocene's glacial and interglacial periods, sea-
level fluctuations have repeatedly increased island area, bring-
ing islands closer to each other and to the adjacent mainland
(Shackleton 1987), thus reducing island isolation and favor-
ing species' colonization (Blondel et al. 2010). More specifi-
cally, 21.5 thousand years ago (Kya) the sea-level was ca 120
m lower than today (Perissoratis and Conispoliatis 2003),
resulting in the fusion of the Ionian islands with western
Greek mainland. A narrow strait separated the landmass of
Kefalonia, Ithaki and Zakynthos from Lefkada; the latter
90




Figure 1. The Ionian islands included in the present study. Numbers correspond to the islets that belong to Echinades islets group as follows:
1 = Oxeia, 2 = Petalas, 3 = Provati, 4 = Dragonera.

0 formed a mainland peninsula, and was connected with most
of its small, satellite islets (i.e. Meganisi, Skorpios, Kastos
and Kalamos). Arkoudi and Atokos were the only islets that
remained isolated during this period (Ferentinos et al. 2012,
5 Zavitsanou et al. 2015). At the same period, Kerkyra was con-
nected with the adjacent mainland. Subsequently, 18 Kya the
sea-level started rising rapidly (Fairbanks 1989, Perissoratis
and Conispoliatis 2003). The present day configuration of
the Ionian islands arose during the Holocene (ca 9–8 Kya),
10 when they became permanently separated from the mainland
and were detached from each other.

The Aegean archipelago has attracted the attention
of many biogeographers and factors affecting plant spe-
cies richness in the Aegean Sea are well studied (Strid
15 1996, Panitsa et al. 2006, Kallimanis et al. 2010,
Kagiampaki et al. 2011, Kougioumoutzis and Tiniakou
2014, Kougioumoutzis et al. 2014, 2017). In contrast,
there are only few similar studies in the Ionian archi-
pelago. Only two macroecological studies have been car-
ried out (Panitsa and Iliadou 2013, Iliadou et al. 2014a).
Moreover, the majority of the biogeographical studies on
the Greek islands have focused on island diversity patterns
(Triantis et al. 2008, Panitsa et al. 2010, Kallimanis et al.
2010, Trigas et al. 2012, Sfenthourakis and Triantis 2017),
25 while there are just a few studies investigating factors
affecting beta-diversity (Fattorini 2006, Panitsa et al. 2008,
Sfenthourakis and Panitsa 2012).

Beta-diversity compared to species richness (alpha-
diversity), allows testing of different hypotheses about the
processes driving species distributions and biodiversity. Beta-
diversity can be partitioned into species turnover and nested-
ness (Baselga and Orme 2012) and this is essential for the
30 investigation of the relative importance of environmental and
spatial drivers. Unraveling the drivers of beta-diversity pat-
terns in the Ionian archipelago would increase our knowledge
on the factors shaping species assemblages on continental
island systems.

The aim of the present study is to investigate the effect
of biotic and abiotic factors shaping plant species diversity
in a rather overlooked Greek archipelago, since our knowl-
edge of the ecological and evolutionary factors driving plant
species assemblages in the Ionian Islands remain vague and
45 narrative. To our knowledge, regression modeling tech-
niques have not before been applied to disentangle the role
of several biogeographical factors in shaping native plant
assemblages in the Ionian archipelago. Additionally, for the
first time, we aim to analyze the beta-diversity in the Ionian
islands, as well as to assess the role of individual islands as
50 reservoirs of plant diversity. Particularly, we have addressed
the following questions: 1) how does richness vary as a func-
tion of area and which are the key factors that affect native
and endemic plant species richness in the Ionian islands?
2) which are the most important drivers of beta-diversity
in the Ionian archipelago? and 3) are there any islands that
55 can be regarded as plant diversity hotspots in the Ionian
archipelago?

Material and methods

Data

We compiled a presence/absence island-plant matrix for the
65 Ionian archipelago, containing 17 islands and 1827 native
plant taxa (species and subspecies, N). We excluded from the
analyses all islands with a surface area < 1 km² or with incom-
plete floristic data (i.e. Kastos). All alien and naturalized taxa
70 were also excluded from the analyses. Species nomenclature
followed Dimopoulos et al. (2013).

We used plant distributional data available from the Flora
Ionica Working Group (2016– and references therein), as well
as some additional unpublished plant records reported by the
75 first author, in order to compile our matrix. Endemicity of
plants was estimated at different levels (Panitsa et al. 2010),
as follows: the total number of endemics (TE) of a single
island was the sum of the taxa that have a distribution range
restricted to Greece (i.e. Greek endemics) and Ionian endem-
ics (IoE) were taxa with their distribution range limited to the
Ionian archipelago (Strid and Tan 1997). Ionian endemics
(IoE) were used only for the identification of Ionian plant
diversity hotspots (hotspot analysis), as their number repre-
85 sents only 1.3% of the overall species richness. Furthermore,
we did not include single island endemics in our analyses,
since the vast majority of the Ionian islands do not host
such taxa (with the exception of Zakynthos, Kefalonia and
Lefkada). The numbers of N, TE, (which constitute the
90 response variables) and IoE, as well as the geographical, cli-
matic, geological and topological data of the Ionian Islands
are shown in Table 1.

Best subset regression analysis of alpha- and beta- diversity

In order to explore the effect of key biological, geographical
and climatic variables on species richness of the Ionian islands,
the determinants of the two species richness metrics (N and
TE) were modeled by two complementary approaches: 1) the
island species-area relationships (ISARs) and 2) best subset
regression analysis (Burnham and Anderson 2002). We also
105 applied best subset regression analysis on beta-diversity and
its components for N.

ISARs were investigated for the Ionian islands by fitting
the logarithmic transformation of the Arrhenius (1921)
power model (power function model hereafter), $\text{Log}(S) = c + z$
 $\times \text{Log}(A)$, where S is the value of each of the richness metrics
used (N and TE), A is the area of the respective island in km²
and c, z the fitted parameters. A value of 1 was added to all
the dependent variables except N before log₁₀-transformation,
because zero values were reported for some islands (Brunet
and Medellín 2001). Most of the variables had frequency
distributions that were strongly positively skewed. Hence,
the variables were log₁₀-transformed to normalize their dis-
tribution so that they could be compared with bivariate and
multivariate regression methods without heteroscedastic
115

61

65

70

75

80

85

90

95

100

105

110

115

121

0 Table 1. Plant species richness and values of the explanatory variables used for the Ionian islands included in the present study. N, TE and 61
IoE are the number of total native, Greek endemic and Ionian endemic taxa respectively. A=area (km²); E=maximum elevation (m);
D_m=shortest distance from the nearest mainland (km); D_i=shortest distance to the nearest island (km); G=number of geological substrates;
HD=human population density (people km⁻²); T=mean annual temperature (°C × 10); P=mean annual precipitation (mm).

5	Island	A	E	D _m	D _i	G	HD	T	P	N	TE	IoE	65
	Antipaxoi	4.1	103	21.7	15.5	1	0.0	170.8	1044.8	227	6	3	
	Atokos	4.4	181	17.6	8.9	3	0.0	174.4	956.0	234	8	1	
	Dragonera	2.4	128	1.5	10.7	1	0.0	175.3	922.7	144	0	0	
	Ereikoussa	3.7	137	27	11	1	134.3	166.0	1109.6	348	2	1	70
10	Ithaki	96.0	806	29.3	2.9	7	0.0	166.4	989.4	874	28	8	
	Kalamos	25.0	745	2.5	13.1	5	19.9	163.4	980.8	610	23	5	
	Kefalonia	734.0	1628	32.6	2.9	14	45.3	160.4	1005.2	1311	65	14	
	Kerkyra	592.0	914	2.2	2.2	12	165.5	161.5	1167.5	1484	32	6	
	Lefkada	325.0	1158	0.1	8.4	7	68.1	155.8	1030.8	1180	58	11	75
15	Mathraki	3.1	152	37.7	8	1	60.0	168.5	1067.5	239	1	1	
	Meganisi	22.4	301	8.9	15	3	46.5	172.5	975.0	428	11	3	
	Othonoi	10.8	393	41.3	21	2	66.3	162.3	1042.7	315	4	2	
	Oxeia	4.3	421	1.2	28.6	1	0.0	175.5	885.0	253	3	1	
	Paxoi	25.3	231	15.6	13.4	2	73.7	167.6	1067.7	549	7	3	
	Petalas	5.5	250	1.3	19.2	1	0.0	175.3	901.7	161	1	0	80
20	Provati	1.2	75	2.5	26.9	1	0.0	178.0	907.0	141	0	0	
	Zakynthos	406.0	756	18.4	14.9	10	100.2	172.1	964.0	1122	36	11	

25 biases and improve the linearity of the relationships in the regression models. A goodness-of-fit test (Shapiro-Wilk and 61
Kolmogorov-Smirnov with 95% level of confidence) to a normal distribution was used to confirm that each transformed
variable was successfully transformed to an approximately normal distribution. We compared the models using R² values
as a measure of their goodness-of-fit. As the models have the same number of fitted parameters, the R² values are directly
comparable, without any modification (Triantis et al. 2003, 2005). The z parameter was used for preliminary comparison
between the floristic diversity of the Ionian islands.

35 We used Akaike's information criterion (AIC) to identify the minimum adequate models. This process also allowed
calculating the relative importance of each explanatory variable, which captured the percentage of variation explained by
each factor when the other factors were held constant. The variance inflator factors (VIF) were below 2.5, thus indicat-
ing that multicollinearity was not a problem in any of our obtained models.

45 The explanatory variables were island area (A, km²), maximum elevation (E, m), shortest distance to nearest main-
land and island (D_m and D_i, respectively, km), number of geological substrates (G), human population density (HD,
people km⁻²), as well as mean annual temperature and precipitation [T and P (°C × 10, mm), respectively]. The
variables A, E, D_i, and D_m were determined from 1:50 000 scale digital topographic maps obtained from the Hellenic
Military Geographical Service (web.gys.gr/GeoSearch/). The number of geological substrates (G) was compiled from the
1:500 000 scale geological map of Greece (Bornovas and Rondogianni-Tsiambaou 1983). From these, 18 different
types of geological substrates were considered, while the most dominant substrates are limestones and Triassic breccias
(remnants of weathered gypsum). We used WorldClim

55 30-arc second climate products (Hijmans et al. 2005) to calculate the climatic variables; T and P were obtained from the
CCSM4 model and were statistically downscaled to a 2.5-arc minute resolution (Hijmans et al. 2005). Data regarding
HD was extracted from the Hellenic Statistical Authority's report (2011). By fitting the full model, the total adjusted
coefficient of multiple determination (R²_{adj}) was assessed. Spearman's correlation coefficient was used for all pairwise
correlations among species richness metrics. Two explanatory variables (E and G) were excluded from the analyses
because they were highly collinear with area (Supplementary material Appendix 1 Table A1). In order to examine the rela-
tive importance of each predictor variable in driving species turnover, we also used the geographical distance between the
Ionian islands (G_D).

Finally, in order to compare the floristic diversity between islands of different size and locate possible Ionian plant diver-
sity hotspots, we used the values of the residuals from the ISAR log₁₀-transformed models of species richness metrics
(N and TE) as well as IoE, as these values are interpreted as a measure of island species diversity and do not reflect the
influence of island area (Hobohm's α-index, Hobohm 2000, 2003). Positive and negative values of this Hobohm's α-index
refer to areas with species diversity above and below average, respectively (Hobohm 2000, 2003).

All analyses and graphs were carried out in the R 3.3.2 (R Development Core Team) using core functions and func-
tions from the 'leaps' (Lumley 2009) and 'ggplot2' (Wickham 2009) packages.

Generalized dissimilarity modelling

We used Generalized dissimilarity modelling (GDM – Ferrier et al. 2007) to model pairwise plant community

0 compositional dissimilarity (quantified with the Sørensen
measure) between all the Ionian islands included in our
analysis as a response to environmental and spatial variables.
In the GDM framework, a group of curvilinear monotonic
5 functions is used to regress species turnover along environ-
mental gradients, consequently identifying the non-linear
relationship between ecological and environmental dis-
similarity (Fitzpatrick et al. 2013). The significance of all
variables was assessed through a Monte Carlo permutation
10 test (999 repetitions; Manion et al. 2018). Hence, we were
able to identify the most significant predictor variables.
We extracted for each of these variables the fitted I-spline
(a curvilinear line expressing the relationship between spe-
cies turnover and each predictor – each I-spline has three
15 coefficients). We then plotted the I-splines to investigate
the magnitude and rate of beta diversity variation along the
most significant predictor variables. We used the sum of the
I-spline's coefficients (it defines the proportion of composi-
tional turnover explained by that variable and is determined
20 by the maximum height of its I-spline; Ferrier et al. 2007,
Fitzpatrick et al. 2013) in order to quantify the magnitude of
turnover along each gradient. Model fit was assessed via per-
cent deviance explained by the model (Manion et al. 2018).
25 The relative importance of each gradient in driving species
turnover was explained as the percent change in deviance
explained by the full model and the deviance explained by a
model fit with that variable permuted (999 permutations).
All GDM analyses were performed with the 'gdm' 1.3.7
30 (Manion et al. 2018) package.

Results

Species richness

The total number of taxa (N) occurring in the island-plant
matrix for the Ionian archipelago was 1827. Eighty-nine
40 taxa (5%) are Greek endemics, while 23 of them (1.3%) are
endemics to the Ionian islands. The number of native taxa per

island varied from 141 to 1473 (Table 1). The proportion of
Greek and Ionian endemics in individual island floras ranged
from 0.0 to 5.0% and 0.0 to 1.1%, respectively.

Species-area relationships

The ISAR power function model explained a higher
proportion of variance for N than TE (Supplementary mate-
rial Appendix 1 Table A2). Nevertheless, TE increased with
70 increasing area more steeply than N.

Predictors of species richness

Simple linear regressions of species richness metrics were
75 applied in order to compare the performance of the explana-
tory variables (Supplementary material Appendix 1 Table A1).
The analyses confirmed A as the most powerful single explan-
atory variable of island plant species richness, while E, T
and G also explained a large proportion of the variance for
80 almost all species richness metrics (Supplementary material
Appendix 1 Table A2).

Predictive modelling

Alpha-diversity

Area (A) and P were retained in the optimal model for N,
while A together with T, D_m, D_i and HD were retained in the
optimal model for TE (Table 2).
90

Generalized Dissimilarity Modelling

Our GDM models explained 66.9% and 46.2% of the
deviance for the native and endemic taxa, respectively. The
most important gradient for both native and endemic taxa
95 was area, followed by mean annual temperature and geo-
graphical distance, respectively (Fig. 2–3). The fitted func-
tions describing the turnover rate and magnitude along
each gradient were not linear; the turnover rate varied with
position along gradients and was greatest at low levels of
100 spatial extent for both metrics and at the high end of the
temperature gradient (Fig. 3). More specifically, the Ionian

Table 2. Summary statistics for predictive models of N and TE, regarding alpha-diversity with A, D_m, D_i, HD, T and P as proposed by the best
subset regression. All variables were log₁₀-transformed. The adjusted R² (total variance explained by each model), the p-value, Akaike's
45 information criterion (AIC), the estimate (Est.), the significance (p), the Bonferroni-corrected p-values (p_{Bon}), as well as the relative importance
(r_w) of each significant predictor of the best subset regression are reported. Abbreviations as in Table 1.

	Est.	t	p	p _{Bon}	AIC	R ² _{adj}	r _w	F	p-value	
N					–31.90	0.94	–	136	< 0.001	110
Intercept	–2.81	–1.42		–	–		–			
A	0.34	14.28	0.00	0.00	–		86.2			
P	1.66	2.50	0.03	0.05	–		14.8			
TE					6.41	0.84	–	17.4	< 0.001	115
Intercept	24.32	1.78		–	–		–			
A	0.64	5.89	0.00	0.00	–		56.8			
T	–10.99	–1.78	0.10	0.50	–		21.1			
D _i	0.32	1.16	0.27	1.00	–		14.0			
HD	–0.17	–1.76	0.11	0.55	–		7.7			
60 D _m	0.12	1.29	0.22	1.00	–		0.4			121




Figure 2. The relative importance of each predictor variable in driving beta-diversity. Abbreviations as in Table 1. G_D =Geographical distance between the Ionian islands.

plant communities (native or endemic) differed significantly even at areas ca 30 km² and their composition continued to change with a rapid, but less abrupt trend until the 400 km² threshold (Fig. 3). Isolation metrics (geographical distance or distance from mainland) had a more profound effect on the endemic plant communities (Fig. 3). Furthermore, climate played an important role in shaping native and endemic plant assemblages in the Ionian archipelago. The endemic plant communities changed rapidly until a plateau was reached at ca 1000 mm of mean annual precipitation (Fig. 3). Finally, a turning point was observed at ~17°C for the native plant assemblages, where they started to differ more swiftly (Fig. 3).

Locating plant diversity hot-spots

Using the α -index (Table 3), Ereikoussa and Kalamos presented values well above average regarding native taxa richness (2.06 and 1.37 respectively), whereas Petalas displayed the lowest value (−2.29). Two islands could be regarded as floristically impoverished (very low α -values): Dragonera and Petalas. For TE, Atokos and Kalamos have remarkably high values (1.81 and 1.69 respectively), while Dragonera has the lowest value (−1.67). Ereikoussa is a diversity hotspot among the Ionian islands, Antipaxoi and Atokos are endemism hotspots, while Kalamos is both a diversity and an endemism hotspot (Fig. 4).

Discussion

Patterns of alpha-diversity

Species–area relationships

The increase in species richness with area is one of the few law-like regularities in ecology (Rosenzweig 1995, Storch 2016, Whittaker et al. 2017). Area is the most powerful single explanatory variable of species richness in both oceanic and continental island systems (Panitsa et al. 2006, Whittaker and Fernández-Palacios 2007, Triantis et al. 2008, 2012, Kagiampaki et al. 2011, Kouglioumoutzis and Tiniakou 2014). In the present study, species richness and area are strongly related for the diversity metrics used, namely the native taxa (N) and the total number of endemics (TE). Our power function model has an overall predictive capacity of $R^2=0.93$ and is consistent with the R^2 -value reported for the ISAR power function model of 488 islands across the globe (Kreft et al. 2008). It is also in line with the R^2 value reported for the Echinades islets group ($R^2=0.84$, Iliadou et al. 2014b) and for some Ionian islands ($R^2=0.89$, Iliadou et al. 2014a).

The z-value is an important parameter that reflects the increase of species richness with increasing surface area. High z-values are associated with high isolation, and consequently




Figure 3. GDM-fitted I-splines (partial regression fits) for variables significantly associated with native (green line) and endemic (blue line) plant beta-diversity in the Ionian archipelago. The maximum height reached by each curve indicates the total amount of turnover associated with that variable, holding all other variables constant. The shape of each function indicates the variation of turnover rate along the gradient.

low immigration rates (Triantis et al. 2012, Matthews et al. 2016). In our study, the ISAR z value ($z=0.37$) fell within the range of 0.2–0.45 proposed by Whittaker and Fernández-Palacios (2007) for insular systems, as in the case of other Greek island complexes. In particular, the z -value for the east Aegean islands is 0.326 (Panitsa et al. 2010), for the south Aegean islands is 0.39 (Kagiampaki et al. 2011) and for the

Cyclades is 0.43 (Kougioumoutzis and Tiniakou 2014). The z -values of off-shore island groups (i.e. Ionian and east Aegean islands) are smaller compared to that of more isolated island complexes (i.e. Cyclades, South Aegean islands), which can be attributed to their palaeogeographical history (Wang et al. 2010, Hu et al. 2011, Ding et al. 2013). Consequently, shallow slopes characterize island groups with minimal isolation, such as the Ionian islands, in which a rescue effect could be observed and therefore explains the higher species richness on smaller islands (Whittaker et al. 2017). Finally, according to our results, TE has the highest z -value. The recent formation and low isolation of the Ionian islands probably promote the colonization of TE in our study system.

Table 3. Values of the α -index for the total native (N), the Greek endemic taxa (TE) and the Ionian endemic taxa (E). Abbreviations as in Table 1.

Island	Alpha-N	Alpha-TE	Alpha-IoE
Antipaxoi	-0.16	1.42	2.02
Atokos	-0.14	1.81	0.01
Dragonera	-1.44	-1.67	-1.31
Ereikoussa	2.06	-0.05	0.20
Ithaki	0.76	0.57	0.88
Kalamos	1.37	1.69	1.18
Kefalonia	-0.93	-0.16	0.12
Kerkyra	0.12	-1.24	-1.92
Lefkada	0.07	0.57	0.38
Mathraki	0.58	-0.62	0.39
Meganisi	-0.10	0.53	0.19
Othonoi	-0.29	-0.28	0.17
Oxeia	0.29	0.33	0.04
Paxoi	0.86	-0.35	0.05
Petalas	-2.29	-1.24	-2.15
Provati	-0.31	-0.90	-0.55
Zakynthos	-0.59	-0.61	0.13

Factors affecting plant species richness

Area and precipitation emerged as the variables most positively influencing plant species richness in the Ionian islands (Table 2). The importance of water availability (and precipitation as its proxy) in ecosystems ranks precipitation among the most significant climatic variables (Kalimeris et al. 2012). It has already been documented in several studies (Kreft and Jetz 2007, Sommer et al. 2010, Kougioumoutzis and Tiniakou 2014) that water availability imposes a limiting factor regarding plant species richness. High species richness likely results from high resource availability and low selective pressure for specialization, suitable for common species (Stohlgren et al. 2005, Weigelt et al. 2013). The Ionian islands are characterized by a humid climate (Krigas et al. 2010) and




Figure 4. (A) Plot of residuals of total endemics (y-axis) versus native taxa (x-axis) and (B) plot of residuals of the Ionian endemics (y-axis) versus native taxa (x-axis), as estimated from the ISAR power function model for Dataset-1.

dense vegetation, which is primarily dominated by common species (Iliadou et al. 2014a), features shared with the continental species pool of western mainland Greece (Kolios and Kalimeris 2017). As islands receive colonizers from climatically similar areas (Weigelt and Kreft 2013), colonization of the Ionian islands is expected to have been unhindered, increasing their plant species richness.

Factors affecting endemic plant species richness

Our results support a strong negative effect of mean annual temperature on TE (Table 2). Furthermore, there is a strong negative correlation between mean annual precipitation and mean annual temperature in the Ionian islands. According to Sommer et al. (2010), in humid regions, such as Ionian islands, with positive water balance, there is a clear positive relationship between the regional capacity for plant species richness and temperature. In a global model for contemporary plant species richness, the relationship between temperature and water availability appeared to be a major limiting factor of the species number a certain region can maintain (Kreft and Jetz 2007, Sommer et al. 2010). In areas with humid conditions, a positive relationship between species richness and temperature was found, i.e. the warmer the temperature, the higher the corresponding species richness (Zachos and Habel 2011). However, the degree of endemism often decreases as floristic richness increases, possibly due to interspecific competition (Médail and Verlaque 1997). The Ionian islands are characterized by a dense vegetation cover, limiting endemic plant taxa to particular habitats with low disturbance and high stress level (Trigas et al. 2008, Georgiou

and Delipetrou 2010 and references therein), such as rocky habitats, calcareous cliffs and slopes. Thus, it seems that temperature has an indirect strong negative effect on endemic plant species richness.

Another possible explanation of the strong negative effect of temperature on the endemic plant species richness comes from its relation to elevation (excluded from the analyses as collinear with A – Supplementary material Appendix 1 Table A1). Decreasing temperature with increasing elevation is widely accepted as a general pattern (Barry 2008). Moreover, endemic species richness usually peaks at higher elevations (Trigas et al. 2013). Topographic complexity may cause gene-flow barriers among diverging populations, supporting reproductive isolation and hence local differentiation (Gillespie and Roderick 2014). Topography-driven isolation increases speciation rates in mountainous areas, resulting in a globally consistent pattern of higher endemism at higher elevations (Steinbauer et al. 2016). The large and mountainous islands of Kefalonia and Lefkada, together with Zakynthos, are the richest islands in terms of Ionian endemics. They are also the only islands that host single island endemics in the Ionian archipelago. As topographic isolation has been related to increased endemism, the low elevation of most Ionian islands seems to be an additional cause of low endemism in the Ionian archipelago (the other being the recent isolation).

Geographical distances (D_i and D_m) have a positive effect on TE. During the Pleistocene glacial and interglacial periods, connections among many Ionian islands and adjacent mainland were established or broken in a chained sequence

of events (Shackleton 1982), while the final formation of the Ionian islands took place during the Holocene (ca 9–8 Kya). The positive effect of geographical distances on endemism indicates that the established inter-island and island-mainland land-bridges during Pleistocene low sea level stages have filtered the distribution of at least some plant lineages.

Human population density (HD) appeared to have a negative effect on TE. Global human activity is affecting species richness, in particular through land-use changes and pollution (Pautasso 2007). The negative effect on TE possibly reflects the effect of the continuous human presence over several millennia in the Ionian islands (Triantis and Mylonas 2009), which are less resistant and resilient to disturbance than mainland areas, a phenomenon also apparent in the central Aegean archipelago (Kougioumoutzis and Tiniakou 2014). Habitat loss as a result of human activities (e.g. tourism, agriculture, animal husbandry) constitutes a major threat for the endemic plants (Trigas et al. 2012).

Beta-diversity patterns

Mediterranean plant communities are characterized by high species turnover along sharp environmental gradients (Molina-Venegas et al. 2016 and references therein). Globally, beta-diversity patterns are best predicted by elevation rather than biotic transitions (Peixoto et al. 2017 and references therein), while extreme temperature (and precipitation) variation occurring over short geographical distances gives rise to increasing beta-diversity (McKnight et al. 2007, Melo et al. 2009, König et al. 2017). This aligns with our results, since in the Ionian archipelago, beta-diversity patterns are primarily driven by area – which is highly collinear with elevation – and temperature (Fig. 2–3). Local island environmental conditions create rather strong filters, thus reducing the set of potential colonizers from the adjacent mainland and hence result in the biotic homogenization of island plant communities (König et al. 2017). The aforementioned patterns are less affected by geographical distance (inter-island or island-mainland distance), suggesting that dispersal filtering plays a minor role in shaping native island plant assemblages in the Ionian. The recent isolation of the Ionian islands and the short distances that separate them from mainland Greece also support this aspect. Nevertheless, dispersal limitation, as expressed by the isolation metrics (D_i and D_m), seems to play an important role in driving endemic beta-diversity patterns in the Ionian archipelago (Fig. 2–3).

Plant diversity hotspots

A promising approach for the long-term maintenance of biodiversity is to identify biodiversity hotspots (sensu Myers et al. 2000). The most frequently applied criterion by conservationists is endemism (Cañadas et al. 2014). In Greece, most endemic plant taxa are chasmophytes, which are mainly found on steep calcareous cliffs and in crevices

(Georgiou and Delipetrou 2010, Panitsa and Kontopanou 2017). There is a great affinity of endemic chasmophytic plant taxa between the Ionian islands and the neighboring phytogeographical regions of mainland Greece with west-facing coastal areas (Panitsa and Kontopanouh 2017) and endemic plant taxa in the Ionian are expected to be found on the larger and more heterogeneous islands (Iliadou et al. 2014a), such as Kefalonia, Kerkyra, Zakynthos and Lefkada, the higher and most topographically complex islands of the Ionian archipelago. However, the island of Kalamos emerged as a plant diversity hotspot in our study area. More specifically, Kalamos has high α -values for both the total native and the endemic plant taxa. This is not surprising, considering 1) Kalamos' uneven terrain (dominated by 2 mountain ranges with elevation of 677 m and 745 m) mostly covered by calcareous slopes (Baliouis 2015) and 2) its unique phytogeographical position since it is located in close proximity to the most rugged landscape of western Greece.

Our hotspot analysis highlights the important role of small islands in the conservation aspects of the Ionian islands. According to MacArthur and Wilson (1967), below a threshold island area, island species richness apparently varies independently of area. Verifying this 'small island effect', small islands exhibited both lowest and highest diversity values for all diversity metrics and IoE in the Ionian archipelago. Our results indicate four small Ionian islands (i.e. Kalamos, Ereikoussa, Antipaxoi and Atokos) as diversity and/or endemism hotspots. These islands belong to the NATURA 2000 network of protected areas. Consequently, small islands with exceptionally high plant diversity could keep a key role in conservation management plans in the Ionian archipelago.

Conclusions

The minimal isolation/recent formation and the close proximity of the Ionian islands to the mainland source have resulted in islands that still behave as parts of a continuous land mass. The rather reduced dispersal filtering (due to the recent formation of the Ionian islands) have led to islands with homogenized plant communities and to island floras principally comprising common species, also explaining the low proportion of endemics. Finally, our hotspot analysis revealed four small islands with exceptionally high plant diversity, with Kalamos emerging as the most important diversity and endemism hotspot. Our results underline the need to integrate more protected areas into the existing Ionian Management Agencies, in order to achieve an effective management of the priority sites for plant conservation in the Ionian islands.

Funding – This study was supported by grants from the General Secretariat for Research and Technology (GSRT) and the Hellenic Foundation for Research and Innovation (HFRI) (grant no. 1585 to A.-T. Valli).

0 References

- Accordi, G. et al. 2014. Microfacies analysis of deep-water breccia clasts: a tool for interpreting shallow- vs deep-ramp Paleogene Sedimentation in Cephalonia and Zakynthos (Ionian islands, Greece). – *Facies* 60: 445–466.
- Arrhenius, O. 1921. Species and area. – *J. Ecol.* 9: 95–99.
- Balioussis, E. 2015. Flora and vegetation of the island of Kalamos (Ionian Sea, Greece) – Floristic analysis and phytogeographical aspects. – *Hacquetia* 14: 307–318.
- Barry, R. G. 2008. Mountain weather and climate. – 3rd ed., Cambridge Univ. Press.
- Baselga, A. and Orme, C. D. L. 2012. Betapart: an R package for the study of beta diversity. – *Methods Ecol. Evol.* 3: 808–812.
- Blondel, J. et al. 2010. The Mediterranean basin – biological diversity in space and time. – Oxford Univ. Press.
- Bornovas, I. and Rondogianni-Tsiambaou, T. 1983. Geological map of Greece. Scale 1:500 000, 2nd ed. – Inst. of Geology and Mineral Exploration, Division of General Geology and Economic Geology, Athens, Greece.
- Brunet, A. K. and Medellín, R. A. 2001. The species-area relationships in bat assemblages of tropical caves. – *J. Mammal.* 82: 1114–1122.
- Burnham, K. P. and Anderson, D. R. 2002. Model selection and multimodel inference: a practical-theoretical approach, 2nd ed. – Springer.
- Cañadas, E. M. et al. 2014. Hotspots within hotspots: endemic plant richness, environmental drivers, and implications for conservation. – *Biol. Conserv.* 170: 282–291.
- Caujapé-Castells, J. et al. 2010. Conservation of oceanic island floras: present and future global challenges. – *Perspect. Plant Ecol.* 12: 107–129.
- Denslow, J. S. 2001. The ecology of insular biota. – *Trends Ecol. Evol.* 16: 423–424.
- Dimopoulos, P. et al. 2013. Vascular plants of Greece: an annotated checklist. – *Englera* 31, Botanischer Garten und Botanisches Museum Berlin-Dahlem, Berlin Germany; Hellenic Botanical Society, Athens, Greece.
- Dimopoulos, P. et al. 2016. Vascular plants of Greece: an annotated checklist. Supplement. – *Willdenowia* 46: 301–347.
- Ding, Z. F. et al. 2013. Patterns of bird functional diversity on land-bridge island fragments. – *J. Anim. Ecol.* 82:781–790.
- Fairbanks, R. O. 1989. A 17 000 year glacio-eustatic sea level record: influence of glacial melting rates on the Younger Dryas event and deep-ocean circulation. – *Nature* 342: 637–642.
- Fattorini, S. 2006. Detecting biodiversity hotspots by species-area relationships: a case study of Mediterranean beetles. – *Conserv. Biol.* 20: 1169–1180.
- Ferentinos, G. et al. 2012. Early seafaring activity in the southern Ionian islands, Mediterranean Sea. – *J. Archaeol. Sci.* 39: 2167–2176.
- Ferrier, S. et al. 2007. Using generalized dissimilarity modelling to analyse and predict patterns of beta diversity in regional biodiversity assessment. – *Divers. Distrib.* 13: 252–264.
- Fitzpatrick, M. C. et al. 2013. Environmental and historical imprints on beta diversity: insights from variation in rates of species turnover along gradients. – *Proc. R. Soc. B* 280: 20131201.
- Flora Ionica Working Group 2016–. Flora Ionica – an inventory of ferns and flowering plants of the Ionian islands (Greece). Available online at <<https://floraionica.univie.ac.at>> (accessed June 2018).
- Georghiou, K. and Delipetrou, P. 2010. Patterns and traits of the endemic plants of Greece. – *Bot. J. Linn. Soc.* 162: 130–422.
- Gillespie, R. G. and Roderick, G. K. 2014. Evolution: geology and climate drive diversification. – *Nature* 509: 297–298.
- Hellenic Statistical Authority 2011. Announcement of the temporary results of the population census 2011. Piraeus, in Greek.
- Hijmans, R. J. et al. 2005. Very high resolution interpolated climate surfaces for global land areas. – *Int. J. Climatol.* 25: 1965–1978.
- Hobohm, C. 2000. Plant species diversity and endemism on islands and archipelagos, with special reference to the Macaronesian islands. – *Flora* 195: 9–24.
- Hobohm, C. 2003. Characterization and ranking of biodiversity hotspots: centers of species richness and endemism. – *Biodivers. Conserv.* 12: 279–287.
- Hu, G. et al. 2011. Determinants of plant species richness and patterns of nestedness in fragmented landscapes: evidence from land-bridge islands. – *Landsc. Ecol.* 26: 1405–1417.
- Iliadou, E. et al. 2014a. Comparing the two Greek archipelagos plant species diversity and endemism patterns highlight the importance of isolation and precipitation as biodiversity drivers. – *J. Biol. Res.* 21: 16.
- Iliadou, E. et al. 2014b. Flora and factors affecting species diversity in protected ‘Natura 2000’ sites of the Ionian area: the Echinades islet group (Greece). – *Willdenowia* 44: 121–136.
- Johnson, J. S. et al. 2017. Landscape genomic insights into the historic migration of mountain hemlock in response to Holocene climate change. – *Am. J. Bot.* 104: 439–450.
- Kagiampaki, A. et al. 2011. Factors affecting plant species richness and endemism in the South Aegean (Greece). – *J. Biol. Res.* 16: 282–295.
- Kalimeris, A. et al. 2012. Long-term precipitation variability in the Ionian islands, Greece (central Mediterranean): climatic signal analysis and future projections. – *Theor. Appl. Climatol.* 109: 51–72.
- Kallimanis, A. S. et al. 2010. Determinants for total and endemic species richness in a continental archipelago. – *Biodivers. Conserv.* 19: 1225–1235.
- Kolios, S. and Kalimeris, A. 2017. Annual and monthly precipitation variability in the Ionian islands using 16-year TRMM data. – In: Karacostas, T. et al. (eds), *Perspectives on atmospheric sciences*. Springer Atmospheric Sciences, Springer, pp. 491–496.
- König, C. et al. 2017. Dissecting global turnover in vascular plants. – *Global Ecol. Biogeogr.* 26: 228–242.
- Kougioumoutzis, K. and Tiniakou, A. 2014. Ecological factors driving plant species diversity in the South Aegean Volcanic Arc and other central Aegean islands. – *Plant Ecol. Divers.* 8: 173–186.
- Kougioumoutzis, K. et al. 2014. Network biogeographical analysis of the central Aegean archipelago. – *J. Biogeogr.* 41: 1848–1858.
- Kougioumoutzis, K. et al. 2017. Network biogeography of a complex island system: the Aegean Archipelago revisited. – *J. Biogeogr.* 44: 651–660.
- Kreft, H. and Jetz W. 2007. Global patterns and determinants of vascular plant diversity. – *Proc. Natl Acad. Sci. USA* 104: 5925–5930.
- Kreft, H. et al. 2008. Global diversity of island floras from a macroecological perspective. – *Ecol. Lett.* 11: 116–127.
- Krigas, N. et al. 2010. Conservation of important plants from the Ionian islands at the Balkan Botanic Garden of Kroussia, N Greece: using GIS to link the in situ collection data with plant

- 0 propagation and ex situ cultivation. – *Biodivers. Conserv.* 19: 3583–3603.
- Lumley, T. 2009. Using Fortran code by Alan Miller, leaps: regression subset selection. – R package, ver. 2.3., <<https://CRAN.R-project.org/package=leaps>>.
- 5 MacArthur, R. H. and Wilson, E. O. 1967. The theory of island biogeography. – Princeton Univ. Press.
- Manion, G. et al. 2018. gdm: generalized dissimilarity modeling. – R package ver. 1.3.8., <<https://cran.r547project.org/package=gdm>>.
- 10 Matthews, T. J. et al. 2016. On the form of species–area relationships in habitat islands and true islands. – *Global Ecol. Biogeogr.* 25: 847–858.
- McKnight, M. W. et al. 2007. Putting beta-diversity on the map: broad-scale congruence and coincidence in the extremes. – *PLoS Biol.* 5: 2424–2432.
- 15 Melo, A. S. et al. 2009. Environmental drivers of beta-diversity patterns in New World birds and mammals. – *Ecography* 32: 226–236.
- Molina-Venegas, R. et al. 2016. How soil and elevation shape local plant biodiversity in a Mediterranean hotspot. – *Biodivers. Conserv.* 25: 1133–1149.
- 20 Médail, F. 2013. The unique nature of Mediterranean island floras and the future of plant conservation. – In: Pons, E. C. et al. (eds), 2nd Botanical Conference in Menorca. Proceedings and abstracts. Islands and plants: preservation and understanding of flora on Mediterranean islands. Maó, Inst. Menorquid' Estudis, Consell Insular de Menorca, pp. 325–350.
- 25 Médail, F. 2017. The specific vulnerability of plant biodiversity and vegetation on Mediterranean islands in the face of global change. – *Reg. Environ. Change* 17: 1775–1790.
- 30 Médail, F. and Myers, N. 2004. Mediterranean basin. – In: Mittermeier, R. A. et al. (eds), Hotspots revisited. Earth's biologically richest and most endangered terrestrial ecoregions. CEMEX, Mexico City, pp. 144–147.
- 35 Médail, F. and Quézel, P. 1999. Biodiversity hotspots in the Mediterranean basin: setting global conservation priorities. – *Conserv. Biol.* 13: 1510–1513.
- Médail, F. and Verlaque, R. 1997. Ecological characteristics and rarity of endemic plants from southeast France and Corsica: implications for biodiversity conservation. – *Biol. Conserv.* 80: 269–281.
- 40 Myers, N. et al. 2000. Biodiversity hotspots for conservation priorities. – *Nature* 403: 853–858.
- Panitsa, M. and Iliadou, E. 2013. Flora and phytogeography of the Ionian islands (Greece). – In: Pons, E. C. et al. (eds), Second Botanical Conference in Menorca. Proceedings and abstracts. Islands and plants: preservation and understanding of flora on Mediterranean islands. Maó, Inst. Menorquid' Estudis, Consell Insular de Menorca, pp. 243–260.
- 45 Panitsa, M. and Kontopanou, A. 2017. Diversity of chasmophytes in the vascular flora of Greece: floristic analysis and phytogeographical patterns. – *Bot. Serb.* 41: 199–211.
- Panitsa, M. et al. 2006. Patterns of species richness on very small islands: the plants of the Aegean Archipelago. – *J. Biogeogr.* 33: 1223–1234.
- 50 Panitsa, M. et al. 2008. Turnover of plants on smallislets of the eastern Aegean Sea within two decades. – *J. Biogeogr.* 35: 1049–1061.
- Panitsa, M. et al. 2010. Factors affecting plant species richness and endemism on land-bridge islands – an example from the East Aegean archipelago. – *Acta Oecol.* 36: 431–437.
- 60 Pautasso, M. 2007. Scale dependence of the correlation between human population presence and vertebrate and plant species richness. – *Ecol. Lett.* 10: 16–24.
- Peixoto, F. P. et al. 2017. Geographical patterns of phylogenetic beta-diversity components in terrestrial mammals. – *Global Ecol. Biogeogr.* 26: 573–583.
- 65 Perissoratis, C. and Conispoliatis, N. 2003. The impacts of sea-level changes during latest Pleistocene and Holocene times on the morphology of the Ionian and Aegean seas (SE Alpine Europe). – *Mar. Geol.* 196: 145–156.
- 70 Rechinger, K. H. 1965. Der Endemismus in der griechischen Flora. – *Revue Roumaine de Biologie, Série Botanique* 10: 135–138.
- Rosenzweig, M. L. 1995. Species diversity in space and time. – Cambridge Univ. Press.
- Sfenthourakis, S. and Panitsa, M. 2012. From plots to islands: species diversity at different scales. – *J. Biogeogr.* 39: 750–759.
- 75 Sfenthourakis, S. and Triantis, K. A. 2017. The Aegean archipelago: a natural laboratory of evolution, ecology and civilizations. – *J. Biol. Res.* 24: 1–13.
- Shackleton, N. J. 1982. The deep-sea sediment record of climate variability. – *Prog. Oceanogr.* 11: 199–218.
- 80 Shackleton, N. J. 1987. Oxygen isotopes, ice volume and sea level. – *Quat. Sci. Rev.* 6: 183–190.
- Sommer, J. H. et al. 2010. Projected impacts of climate change on regional capacities for global plant species richness. – *Proc. R. Soc. B* 277: 2271–2280.
- 85 Steinbauer, M. J. et al. 2016. Topography-driven isolation, speciation and a global increase of endemism with elevation. – *Global Ecol. Biogeogr.* 25: 1097–1107.
- Stohlgren, T. J. et al. 2005. Plant species invasions along the latitudinal gradient in the United States. – *Ecology* 86: 2298–2309.
- 90 Storch, D. 2016. The theory of the nested species-area relationship: Geometric foundations of biodiversity scaling. – *J. Veg. Sci.* 27: 880–891.
- Strid, A. 1996. Phytogeographia Aegaea and the Flora Hellenica Database. – *Ann. Nat. Hist. Mus. Wien* 98: 279–289.
- 95 Strid, A. and Tan, K. 1997. Flora Hellenica. Vol. 1. – Koeltz Scientific Books, Königstein.
- Thompson, J. D. et al. 2005. Ecological differentiation of Mediterranean endemic plants. – *Taxon* 54: 967–976.
- 100 Triantis, K. A. and Mylonas, M. 2009. Greek islands, biology. – In: Gillespie, R. and Glague, D. A. (eds), Encyclopedia of islands. Univ. of California Press, pp. 388–392.
- Triantis, K. A. et al. 2003. A model for the species-area-habitat-relationship. – *J. Biogeogr.* 30: 19–27.
- 105 Triantis, K. A. et al. 2005. Species richness, environmental heterogeneity and area: a case study based on land snails in Skyros archipelago (Aegean Sea, Greece). – *J. Biogeogr.* 32: 1727–1735.
- Triantis, K. A. et al. 2008. Evolutionary species-area curves as revealed by single-island endemics: insights for the inter-provincial species-area relationship. – *Ecography* 31: 401–407.
- 110 Triantis, K. A. et al. 2012. The island species–area relationship: biology and statistics. – *J. Biogeogr.* 39: 215–231.
- Trigas, P. et al. 2008. Vascular plant species diversity, biogeography and vulnerability in the Aegean islands as exemplified by Evvia island (W Aegean, Greece). – *Fresen. Environ. Bull.* 17: 48–57.
- 115 Trigas, P. et al. 2012. Distribution patterns and conservation perspectives of the endemic flora of Peloponnese (Greece). – *Folia Geobot.* 47: 421–439.
- 120

0	Trigas, P. et al. 2013. Elevational gradient of vascular plant species richness and endemism in crete – the effect of post-isolation mountain uplift on a continental island system. – PLoS One 8: e59425.	61
5	Turrill, W. B. 1929. The plant life of the Balkan peninsula. – Clarendon Press.	65
	Vogiatzakis, I. N. et al. 2008. Mediterranean island landscapes: natural and cultural approaches. Landscape series. Vol 9. – Springer.	
10	Wang, Y. P. et al. 2010. Nestedness for different reasons: the distributions of birds, lizards and small mammals on islands of an inundated Lake. – Divers. Distrib. 16: 862–873.	70
15	Weigelt, P. and Kreft, H. 2013. Quantifying island isolation – insights from global patterns of insular plant species richness. – Ecography 36: 417–429.	75
	Weigelt, P. et al. 2013. Bioclimatic and physical characterization of the world's islands. – Proc. Natl Acad. Sci. USA 110: 15307–15312.	
	Whittaker, R. J. and Fernández-Palacios, J. M. 2007. Island biogeography: ecology, evolution and conservation, 2nd ed. – Oxford Univ. Press.	
	Whittaker, R. J. et al. 2017. Island biogeography: taking the long view of nature's laboratories. – Science 357: eaam8326.	
	Wickham, H. 2009. ggplot2: elegant graphics for data analysis, 2nd ed. – Springer Science.	
	Zachos, F. E. and Habel, J. C. 2011. Biodiversity hotspots: distribution and protection of conservation priority areas. – Springer.	
	Zavitsanou, A. et al. 2015. Palaeogeographic reconstruction of the Inner Ionian Sea during Late Pleistocene low sea level stands: preliminary results. – 11th Panhellenic Symposium on Oceanography and Fisheries, Mytilene, Lesbos island, Greece.	
20	Supplementary material (Appendix NJB-02156 at <www.nordicjbotany.org/appendix/njb-02156>). Appendix 1.	80
25		85
30		90
35		95
40		100
45		105
50		110
55		115
60		121

Author Queries

JOB NUMBER: 2156

JOURNAL: OIK_NJB

- Q1 There is a mismatch in author names between the reference citation and the list for the reference 'Georgiou and Delipetrou 2010'. Please check.
- Q2 There is a mismatch in author names between the reference citation and the list for the reference 'Panitsa and Kontopanouh 2017'. Please check.