

U N	Goal	3. Ensure healthy lives and promote well-being for all at all ages
	Target	3.4 By 2030, reduce by one third premature mortality from non-communicable diseases through prevention and treatment and promote mental health and well-being
	Indicator	3.4.2 Suicide mortality rate

I. National indicator

<Type 1>

Indicator	Suicide mortality rate
Definition	The number of suicide deaths per 100,000 population(X60-X84) * Classification under the seventh Korean Standard Classification of Diseases(KCD-7).
Calculation method	$\text{Suicide mortality rate} = \frac{\text{Suicide deaths}}{\text{Mid-year resident population}} \times 100,000$
Unit	Deaths per 100,000 population

II. National indicator's source

Data sources	<ul style="list-style-type: none"> ■ Source: <i>Cause of Death Statistics</i> ■ Collection method: Compiling data from death reports filed with city, district, and municipal offices, baby and fetal death reports from crematories, and data from the 'Complementary Survey on Causes of Death' of medical institutions
Calendar	<ul style="list-style-type: none"> ■ Frequency: Annually (Jan 1 - Dec 31 of the reference year) ■ Data release: September in the following year
Organizations	Vital Statistics Division(Tel. 042-481-2252), Statistics Korea
Related International Agency	WHO

III. Comparison with UN SDG indicator

① Indicator		② Definition		③ Data value	
Same	Different	Same	Different	Same	Different
Note	<ul style="list-style-type: none"> ■ Deviations from national statistics due to statistical correction by the WHO(i.e. unidentified deaths(Y10-Y34, Y872) classified as suicide, or incomplete information for death registration) 				
Global indicator link	<ul style="list-style-type: none"> ■ Metadata: https://unstats.un.org/sdgs/metadata/files/Metadata-03-04-02.pdf ■ Data: https://unstats.un.org/sdgs/indicators/database/ 				