

Wuhan-coronavirus homologue map

Kou Amano

Background and Object

Background: At the beginning of 2020, global risk of infection of coronavirus is spreaded. The pandemic started in 2019 and governments announced a state of emergency. In Japan, the government adopted PCR as a diagnosis method of the infection. But the selection of prmers infrence the accuracy greatly.

Object: Therefor, I provide a "map" of homological region of coronavirus to other viruses and animals to help the primer design.

Data

Vertebrates:

● Bat (NW_017738920.1 .. NW_017802358.1; 63439frgs)	● Dog (NC_006583.3 .. NC_006621.3; 39chrs)	● Rabbit (NC_013669.1 .. NC_013690.1; 22chrs)
● Beluga (NW_022097992.1 .. NW_022103895.1; 5904frgs)	● Ferret (NW_004569142.1 .. NW_004576923.1; 7782frgs)	● Pig (NC_010443.5 .. NC_010462.3; 20chrs)
● Camel (NC_044511.1 .. NC_044547.1; 37chrs)	● Human (NC_000001.4 .. NC_000024.3; 24chrs)	● Rock Pigeon (NW_004973171.1 .. NW_004988092.1; 14922frgs)
● Cat (NC_018723.3 .. NC_018741.3; 19chrs)	● Mouse (NC_000067.6 .. NC_000087.7; 21chrs)	● Turkey (NC_015011.2 .. NC_015042.2; 32chrs)

Viruses:

Method

blast

self-blast

window-fourrier

The plot

