# Transaktionssysteme – Übungsblatt 2

Gruppe 1, Team Zoidberg

Koppera Thomas 51366

Kinseher Josef

Treuberg Matthias 51164

#### Aufgabe 1

a)  $S := w_0(x)w_0(y)w_0(z)r_1(y)r_2(y)w_1(y)r_3(z)r_2(x)w_2(x)w_1(x)c_1c_2c_3$ 

$$H_{S}(w_{0}(x)) = f_{0}^{x}()$$

$$H_{S}(w_{0}(y)) = f_{0}^{y}()$$

$$H_{S}(w_{0}(z)) = f_{0}^{z}()$$

$$H_{S}(r_{1}(y)) = H_{S}(w_{0}(y)) = f_{0}^{y}()$$

$$H_{S}(r_{2}(y)) = H_{S}(w_{0}(y)) = f_{0}^{y}()$$

$$H_{S}(w_{1}(y)) = H_{S}(r_{1}(y), r_{2}(y)) = f_{1}^{y}(f_{0}^{y}(), f_{0}^{y}())$$

$$H_{S}(r_{3}(z)) = H_{S}(w_{0}(z)) = f_{0}^{z}()$$

$$H_{S}(w_{3}(z)) = H_{S}(r_{3}(z)) = f_{3}^{z}(f_{0}^{z}())$$

$$H_{S}(r_{2}(x)) = H_{S}(w_{0}(x)) = f_{0}^{x}()$$

$$H_{S}(w_{2}(x)) = H_{S}(r_{2}(x)) = f_{2}^{x}(f_{0}^{x}())$$

$$H_{S}(w_{1}(x)) = H_{S}(r_{2}(x)) = f_{1}^{x}(f_{0}^{x}())$$

b)

### Aufgabe 2

 $S := w_0(x)w_0(y)w_0(z)c_0r_1(x)r_2(x)w_2(y)r_3(x)c_1r_2(z)w_3(z)c_3w_2(z)c_2$ 

a)

$$\begin{split} H_S(w_0(x)) &= f_0^x() \\ H_S(w_0(y)) &= f_0^y() \\ H_S(w_0(z)) &= f_0^z() \\ H_S(r_1(x)) &= H_S(w_0(x)) = f_0^x() \\ H_S(r_2(x)) &= H_S(w_0(x)) = f_0^x() \\ H_S(w_2(y)) &= f_2^y() \\ H_S(r_3(x)) &= H_S(w_0(x)) = f_0^x() \\ H_S(r_2(z)) &= H_S(w_0(z)) = f_0^z() \\ H_S(w_3(z)) &= H_S(r_2(z)) = f_3^z(f_0^z()) \\ H_S(w_2(z)) &= H_S(r_2(z)) = f_2^z(f_0^z()) \end{split}$$

b)

$$H[S](x) = H_S(w_0(x)) = f_0^x()$$

$$H[S](y) = H_S(w_2(y)) = f_2^y()$$

$$H[S](z) = H_S(w_2(z)) = f_2^z(f_0^z())$$

- c)  $S' := w_0(x)w_0(y)w_0(z)c_0w_2(y)c_1r_2(z)c_3w_2(z)c_2$
- d) Nein da  $op(S) \neq op(S')$

## Aufgabe 3

a)  $H_S(w_3(z)) = f_3^z(f_0^z())$ 

## Aufgabe 4

- a)
- b)