
Euscorpius

Occasional Publications in Scorpiology



Scorpions of the Horn of Africa (Arachnida, Scorpiones). Part XIV. *Hottentotta somalicus* sp. n. (Buthidae) from Somalia

František Kovařík

February 2018 - No. 256

Euscorpius

Occasional Publications in Scorpiology

EDITOR: Victor Fet, Marshall University, 'fet@marshall.edu' ASSOCIATE EDITOR: Michael E. Soleglad, 'soleglad@znet.com'

Euscorpius is the first research publication completely devoted to scorpions (Arachnida: Scorpiones). **Euscorpius** takes advantage of the rapidly evolving medium of quick online publication, at the same time maintaining high research standards for the burgeoning field of scorpion science (scorpiology). **Euscorpius** is an expedient and viable medium for the publication of serious papers in scorpiology, including (but not limited to): systematics, evolution, ecology, biogeography, and general biology of scorpions. Review papers, descriptions of new taxa, faunistic surveys, lists of museum collections, and book reviews are welcome.

Derivatio Nominis

The name *Euscorpius* Thorell, 1876 refers to the most common genus of scorpions in the Mediterranean region and southern Europe (family Euscorpiidae).

Euscorpius is located at: http://www.science.marshall.edu/fet/Euscorpius

(Marshall University, Huntington, West Virginia 25755-2510, USA)

ICZN COMPLIANCE OF ELECTRONIC PUBLICATIONS:

Electronic ("e-only") publications are fully compliant with ICZN (<u>International Code of Zoological Nomenclature</u>) (i.e. for the purposes of new names and new nomenclatural acts) when properly archived and registered. All **Euscorpius** issues starting from No. 156 (2013) are archived in two electronic archives:

- **Biotaxa**, http://biotaxa.org/Euscorpius (ICZN-approved and ZooBank-enabled)
- Marshall Digital Scholar, http://mds.marshall.edu/euscorpius/. (This website also archives all *Euscorpius* issues previously published on CD-ROMs.)

Between 2000 and 2013, ICZN did not accept online texts as "published work" (Article 9.8). At this time, *Euscorpius* was produced in two identical versions: online (*ISSN 1536-9307*) and CD-ROM (*ISSN 1536-9293*) (laser disk) in archive-quality, read-only format. Both versions had the identical date of publication, as well as identical page and figure numbers. Only copies distributed on a CD-ROM from *Euscorpius* in 2001-2012 represent published work in compliance with the ICZN, i.e. for the purposes of new names and new nomenclatural acts.

In September 2012, ICZN Article 8. *What constitutes published work*, has been amended and allowed for electronic publications, disallowing publication on optical discs. From January 2013, *Euscorpius* discontinued CD-ROM production; only online electronic version (*ISSN 1536-9307*) is published. For further details on the new ICZN amendment, see http://www.pensoft.net/journals/zookeys/article/3944/.

Scorpions of the Horn of Africa (Arachnida, Scorpiones). Part XIV. *Hottentotta somalicus* sp. n. (Buthidae) from Somalia

František Kovařík

P. O. Box 27, CZ-145 01 Praha 45, Czech Republic; www.scorpio.cz

http://zoobank.org/urn:lsid:zoobank.org:pub:86482F7C-F1F7-4667-B099-ED496D6E053C

Summary

Hottentotta somalicus **sp. n.** from Somalia is described and fully complemented with color photos. Morphologically it is similar to *H. polystictus* (Pocock, 1896). These two species have very narrow metasomal segments (1.63–1.73 in both sexes versus 1.31–1.61 in both sexes of other *Hottentotta* species from the Horn of Africa). *H. polystictus* and *H. somalicus* **sp. n.** occur in separate areas (Somaliland versus Somalia) and can be differentiated by color.

Introduction

In 2011–2017, the author has had an opportunity to participate in expeditions to the Horn of Africa, study scorpions at 113 localities in Ethiopa, Eritrea, and Somaliland, and published several articles on this fauna. During the expeditions he collected a lot of *Hottentotta* specimens which were summarized in a revison published by Kovařík & Mazuch in 2015. Here, we describe another species, *Hottentotta somalicus* sp. n. from the vicinity of Mogadishu, which belongs to the *Hottentotta polystictus* complex.

Methods, Material & Abbreviations

Nomenclature and measurements follow Stahnke (1971), Kovařík (2009), and Kovařík & Ojanguren Affilastro (2013), except for trichobothriotaxy (Vachon, 1974). Short, stout spiniform macrosetae are termed spinules.

We intentionally use here the name Somaliland (Hargeysa) for the northern territory corresponding to the former British colony (British Somaliland), which we distinguish from Somalia (Mogadishu). Somaliland has its own currency, a functional government with representation in several countries. Specimens used for this study were collected and imported with permitions of Amoud and Hargeisa Universities and Ministry of the Environment of the Republic of Somaliland.

All studied specimens are preserved in 80% ethanol in the author's collection, FKCP (František Kovařík,

private collection, Prague, Czech Republic). *Morphometrics*: D, depth; L, length; W, width.

Systematics

Family Buthidae C. L. Koch, 1837

Hottentotta Birula, 1908 (Figs. 1–24, Table 1)

Buthus (Hottentotta) Birula, 1908: 141.

Hottentotta: Fet & Lowe, 2000: 133–144: Kovařík, 2007: 1–107, figs. 1–153; Kovařík & Mazuch, 2015: 1–37, figs. 1–158.

- = *Dasyscorpio* Pallary, 1938: 279–280 (syn. by Vachon, 1949: 146 (1952: 232)).
- = *Buthotus* Vachon, 1949: 143 (1952: 229–241, figs. 313–330) (syn. by Francke, 1985: 4).
- *Mesobuthus*: Tikader & Bastawade, 1983: 185–242, figs. 504–700.
- = Buthotus (Balfourianus) Vachon, 1979: 236, fig. 8 (syn. by Kovařík, 2007: 2).
- = Hottentotta (Deccanobuthus) Lourenço, 2000: 192–195, figs. 1–7 (syn. by Kovařík, 2007: 2).

Type species. Scorpio hottentotta Fabricius, 1787.

DIAGNOSIS. Medium to large buthids, adults 30–130 mm. Sternum type 1 (Soleglad & Fet, 2003), triangular in shape. Pedipalps orthobothriotaxic, type A β (Vachon, 1974, 1975), femur trichobothrium d_2 dorsal, patella d_3

dorsal of dorsomedian carina. Chelal trichobothrium db usually located between est and et, or may be on level with trichobothrium est, rarely between est and esb. Trichobothrium eb clearly on fixed finger of pedipalp. Pectines with fulcra. Dentate margin of pedipalp-chela movable finger with distinct granules divided into 11-16 linear rows and (4)5-7 terminal granules. Chelicerae with typical buthid dentition, fixed finger armed with two denticles on ventral surface. Tergites I-VI granular, with three carinae, tergite VII with 5 carinae. Carapace with distinct carinae, entire dorsal surface nearly planate. First sternite with two granulated lateral stridulatory areas, which however may be reduced in some species (e. g. in *H. pachyurus* and *H. trilineatus*). Metasoma elongate, segment I with 10 carinae, segments II-IV with 8-10 carinae. Ventrolateral carinae of fifth metasomal segment with all granules more or less equal in size and never lobate. Telson bulbous, lumpy and granulated, without subaculear tooth. Legs III and IV with well developed tibial spurs, first and second tarsomeres with paired ventral setae.

Hottentotta polystictus (Pocock, 1896) (Figs. 9–13)

Buthus polystictus Pocock, 1896: 178–179, pl. XI, fig. 1.
Hottentotta polystictus: Kovařík & Ojanguren, 2013: 171–172, 318, 338–339, figs. 1069–1072, 1206–1216 (complete reference list until 2013); Kovařík & Mazuch, 2015: 23, figs. 112–131, table 4).

TYPE LOCALITY AND TYPE REPOSITORY. Somaliland, Goolis Mts., inland of Berbera; BMNH (The Natural History Museum, London, United Kingdom).

ETHIOPIAN MATERIAL EXAMINED. Ethiopia, NE of Dire Dawa, on road to Djibouti, 09°37'59"N 41°52'43"E, 1124 m a.s.l., 30.I.2015, 1° , leg. T. Mazuch.

SOMALILAND MATERIAL EXAMINED. Somaliland, Hamas, between Hargeisa and Berbera, 10°02.267'N 44°47. 299'E, 650 m a.s.l., XI.2010, 1♂1♀, leg. T. Mazuch; 70 km from Berbera to Hargeysa, XI.2010, 13, leg. T. Mazuch; Laas Gel, 50 km NE Hargeisa, 09°46'16.6"N 44°27'07.2"E, 1090 m a.s.l., 7.VII.2011, 1♂2♀3juvs., leg. F. Kovařík; beetwen Berbera and Sheikh, 10°05' 49.9"N 45°11'40.1"E, 628 m a.s.l., 10.VII.2011, 1♀im., leg. F. Kovařík; Sheikh, Goolis Mts., 09°56'23"N 45° 11'14.2"E, 1439 m a.s.l., 11.VII.2011, 2♀2juvs., leg. F. Kovařík; beetween Burao and Laas Caanood, 09°11' 18.4"N 45°54'24"E, 871 m a.s.l., 11.VII.2011, 1♀2juvs., leg. F. Kovařík; Sheikh, 09°57'25.9"N 45°09'52.2"E, 1492 m a.s.l., 12.VII.2011, 1♂7♀, leg. F. Kovařík; 15 km N of Sheikh, Goolis Mts., 09°32'27.7"N 45°31' 38.9"E, 1056 m a.s.l, 6.II.2017, 1♀1juv., leg. F. Kovařík; Laas Gel, 50 km NE Hargeisa, 09°46'47"N 44°26'43"E, 1043 m a.s.l., 7.II.2017, 1♀, leg. F. Kovařík; Laas Gel, 50 km NE Hargeisa, 09°46'47"N 44°26'43"E, 1043 m a.s.l., 28.-30.VIII.2017, 2♂3♀7 juvs., leg. F. Kovařík; between Berbera and Burao, 10°02'12"N 44°47'21"E, 60 m a.s.l., 30.VIII.2017, 1♂im., leg. F. Kovařík; Sheikh, Goolis Mts., 09°56'38"N 45°10'59"E, 1418 m a.s.l., 6.IX.2017, 1♂9♀8juvs., leg. F. Kovařík; Borama, Amoud University campus, 09°56'49"N 43°13'23"E, 1394 m a.s.l., 9-13.IX.2017, 1♂, leg. F. Kovařík.

DIAGNOSIS. Total length 40-60 mm, some males may be only 35 mm long. Trichobothrium db on fixed finger of pedipalp situated between trichobothria et and est. Sexual dimorphism not pronounced; manus of pedipalp of approximately same width in both sexes, but males have fingers of pedipalps slightly twisted. Pectinal teeth number 22-27 in males, 18-22 in females. Chelicerae yellow, anterior part could be reticulated in darker specimens. Pedipalps sparsely hirsute. Metasoma with only a few hairs. Color uniformly yellowish brown. Dorsal surfaces of pedipalps and ventral surfaces of metasoma with numerous dark spots. Mesosoma yellowish to reddish brown, with black spots. Femur of pedipalp with 5 carinae that may be incomplete. Patella with 8 carinae, of which some are smooth, without granules and obsolete. Chela lacks carinae but is usually granulate. Movable fingers of pedipalps with 12–14 rows of granules and 5 or 6 terminal granules. Seventh sternite bears 4 well marked carinae, usually granulated. First to third metasomal segments with 10 carinae; fourth with 8 or 10 carinae; fifth with 5 carinae. All carinae granulated, dorsal carinae bear larger terminal granules. Metasoma very narrow. First metasomal segment of adults usually longer than wide or as long as wide, second metasomal segment always longer than wide. Fourth metasomal segment length/width ratio 1.70-1.73 in both sexes. Telson very bulbous. Telson length/depth ratio 2.45-2.48 in females.


Hottentotta somalicus sp. n. (Figs. 1-8, 14-24, Table 1) http://zoobank.org/urn:lsid:zoobank.org:act:C9A7E9 D8-E654-4D6B-B7DD-C2935ABBC76B

TYPE LOCALITY AND TYPE DEPOSITORY. Somalia, near Mogadishu; FKCP.

TYPE MATERIAL EXAMINED. Somalia, near Mogadishu, IX.2011, $1 \circlearrowleft$ (holotype) $2 \circlearrowleft$ (paratypes), FKCP.

ETYMOLOGY. Named after the country of occurence.


DIAGNOSIS. Total length 40 (male) to 65 (female) mm. Trichobothrium *db* on fixed finger of pedipalp is situated between trichobothria *et* and *est*. Sexual dimorphism not



Figures 1–4: *Hottentotta somalicus* **sp. n. Figures 1–2.** Male holotype in dorsal (1) and ventral (2) views. **Figures 3–4:** Female paratype in dorsal (3) and ventral (4) aspects. Scale bars: 10 mm.

pronounced; manus of pedipalp of approximately same width in both sexes, but males have fingers of pedipalps slightly twisted. Pectinal teeth number 23 in males, 19–


21 in females. Chelicerae yellow. Pedipalps sparsely hirsute. Metasoma with only a few hairs. Color uniformly yellowish brown. Femur of pedipalp with 4 or 5



Figures 5–10: Figures 5–8: *Hottentotta somalicus* sp. n. Figures 5–6. Male holotype, carapace and tergites I–III (5), coxosternal area and sternites (6). Figures 7–8. Female paratype, carapace and tergites I–III (7), coxosternal area and sternites III–IV (8). Figures 9–10: *Hottentotta polystictus*, female from Ethiopia, NE of Dire Dawa, on road to Djibouti, 09°37'59"N 41°52'43"E, 1124 m a.s.l., chelicerae, carapace and tergites I–III (9); male from Somaliland, Sheikh, 09°57'25.9"N 45°09'52.2"E, 1492 m a.s.l., coxosternal area and sternites (10).

carinae. Patella with 8 carinae, of which some are smooth, without granules and obsolete. Chela lacks carinae but is usually finely granulate. Movable fingers of pedipalps with 13–14 rows of granules and 5 or 6 terminal granules. Sternite WII bears 4 well marked

granulated carinae. First to third metasomal segments with 10 carinae; fourth with 8 or 10 carinae; fifth with 5 carinae. All carinae granulated, dorsal carinae bear larger terminal granules. Metasoma very narrow. All metasomal segment of adults longer than wide. Fourth



Figures 11–16: Metasoma and telson lateral (11, 14), ventral (12, 15), and dorsal (13, 16). **Figures 11–13:** *Hottentotta polystictus*, male from Somaliland, Sheikh, 09°57'25.9"N 45°09'52.2"E, 1492 m a.s.l. **Figures 14–16:** *Hottentotta somalicus* **sp. n.**, male holotype. Scale bar: 10 mm.


metasomal segment length/width ratio 1.63–1.77 in both sexes. Telson bulbous. Telson length/depth ratio 2.41–2.49 in females.

DESCRIPTION. Total length 40 (male) to 65 (female) mm. The habitus is shown in Figs. 1–4. For position and distribution of trichobothria of pedipalps see Figs. 17–21, 23. Note, four internal femoral trichobothria are present but not visible in these figures. Sexual dimorphism

minor, there is no difference in length and width of metasomal segments.

Coloration (Figs. 1–4). The base color is uniformly yellow to yellowish brown, dark spots absent or extremely reduced.

Carapace and mesosoma (Figs. 5–8). The carapace is carinate and unevenly covered by granules of varying size; much of the granulation is fine, but some granules



Figures 17–24: Figures 17–23: *Hottentotta somalicus* sp. n., male holotype, pedipalp chela dorsal (17), externodorsal (18) and ventrointernal (19), pedipalp patella dorsal (20), external (21) and ventral (22), and trochanter and femur dorsoexternal (23). **Figure 24**. Female paratype, pedipalp chela dorsal (24). Trichobothrial pattern is indicated in Figures 17–21 and 23.

Dimensions (MM)		H. somalicus sp. n. ♂ holotype	H. somalicus sp. n. ♀ paratype
Carapace	L/W	4.35 / 4.50	6.88 / 7.90
Mesosoma	L	11.10	18.60
Tergite VII	L/W	3.03 / 4.40	4.70 / 8.45
Metasoma & telson	L	24.78	34.80
Segment I	L/W/D	3.15 / 2.83 / 2.60	4.35 / 4.25 / 3.80
Segment II	L/W/D	3.55 / 2.55 / 2.40	4.90 / 3.80 / 3.60
Segment III	L/W/D	3.70 / 2.48 / 2.46	5.05 / 3.65 / 3.50
Segment IV	L/W/D	4.25 / 2.40 / 2.35	5.65 / 3.45 / 3.65
Segment V	L/W/D	5.33 / 2.47 / 2.30	7.60 / 3.50 / 3.30
Telson	L/W/D	4.80 / 2.06 / 2.05	7.96 / 3.20 / 3.30
Pedipalp	L	16.38	24.00
Femur	L/W	3.80 / 1.30	5.60 / 2.05
Patella	L/W	4.78 / 2.03	7.10 / 3.06
Chela	L	7.80	11.30
Manus	L/W/D	2.70 / 2.05 / 1.98	3.60 / 3.10 / 2.88
Movable finger	L	5.10	7.70
Total	L	40.23	60.28

Table 1: Comparative measurements of adults of *Hottentotta somalicus* **sp. n.** Abbreviations: length (L), width (W, in carapace it corresponds to posterior width), depth (D).

are larger and distinctly rounded. Tergites I–VI bear three carinae and are granulated, with some intercarinal granules small and others larger and rounded. Tergite VII is pentacarinate. The pectinal tooth count is 23 in

male and 19–21 in females. The pectinal marginal tips extend to about the third quarter of the fourth sternite in males and about a quarter of the fourth sternite in females. The pectines have three marginal lamellae and

eight to nine middle lamellae. The lamellae bear numerous dark long setae, each fulcrum with three to five setae. All sternites are smooth and sparsely hirsute. The seventh sternite bears four granulate carinae. The other sternites bear two furrows.

METASOMA AND TELSON (Figs. 14–16). All metasomal segments are only very sparsely hirsute. All metasomal segments are longer then wide. Segments I–III bear 10 carinae, segment IV bears 8 or 10 carinae and segment V bears five carinae, three or five ventral and two dorsal. All carinae are granulated. The dorsal surface of all segments is smooth to very finely granulated. Other surfaces are granulated. The telson is only sparsely hirsute, bulbous, lumpy and granulated.

PEDIPALPS (Figs. 17–24). The pedipalps are sparsely hirsute and weakly granulated. The femur bears four or fifth granulated carinae. The patella bears eight granulated carinae. The chela is without carinae. The movable fingers of pedipalps bear 13–14 rows of granules and five or six terminal granules.

LEGS. The tarsomeres bear two rows of macrosetae on the ventral surface and numerous macrosetae on the other surfaces; bristle combs absent. Femur bears only several macrosetae. Femur coarsely granulose, femur and patella with carinae well developed. Tibial spurs present and long on third and fourth legs and absent in the other legs.

Measurements. See Table 1.

AFFINITIES. The described features distinguish H. somalicus sp. n. from all other species of the genus. The morphology makes the new species close to H. polystictus. Females of most Hottentotta from the Horn of Africa have telson length/depth ratio 2.06-2.36 and females of *H. polystictus* and *H. somalicus* **sp. n.** usually have telson length/depth ratio 2.41-2.49. These two "sibling" species also have very narrow metasomal segments (1.63-1.73 in both sexes versus 1.31-1.61 in both sexes of other Hottentotta species of Horn of Africa). H. polystictus and H. somalicus sp. n. occur in separate areas (fig. 158 in Kovařík & Mazuch, 2015: 35) and can be differentiated by color. H. somalicus sp. n. is uniformly yellow to yellowish brown with dark spots absent or extremely reduced (Figs. 1–8, 14–16) while H. polystictus has dark spots strongly developed (Figs. 9-12). Because this character might not be stable, it is important that the DNA analysis also indicates divergence between these two species (Šťáhlavský et al., in preparation).

References

BIRULA, A. A. 1908. Ergebnisse der mit Subvention aus der Erbschaft Treitl unternommenen zoologischen Forschungsreise Dr. F. Werner's nach dem ägyptischen Sudan und Nord-Uganda. XIV.

- Scorpiones und Solifugae. Sitzungsberichte der Kaiserlich-Königlichen Akademie der Wissenschaften, Wien, 117/2 (1): 121–152.
- FET, V. & G. LOWE. 2000. Family Buthidae C. L. Koch, 1837. Pp. 54–286 in Fet, V., W. D. Sissom, G. Lowe & M. E. Braunwalder. *Catalog of the Scorpions of the World (1758–1998)*. New York: The New York Entomological Society, 689 pp.
- FRANCKE, O. F. 1985. Conspectus genericus Scorpionorum 1758–1982 (Arachnida: Scorpiones). Occasional Papers of the Museum, Texas Tech University, 98: 1–32.
- KOVAŘÍK, F. 2007. A revision of the genus *Hottentotta* Birula, 1908, with descriptions of four new species (Scorpiones, Buthidae). *Euscorpius*, 58: 1–107.
- KOVAŘÍK, F. 2009. Illustrated catalog of scorpions. Part I. Introductory remarks; keys to families and genera; subfamily Scorpioninae with keys to Heterometrus and Pandinus species. Prague: Clairon Production, 170 pp.
- KOVAŘÍK, F. & T. MAZUCH. 2015. Scorpions of Ethiopia (Arachnida: Scorpiones). Part III. Genus *Hottentotta* Birula, 1908 (Buthidae), with description of three new species. *Euscorpius*, 202: 1–37
- KOVAŘÍK, F. & A.A. OJANGUREN AFFILASTRO. 2013. *Illustrated catalog of scorpions. Part II. Both-riuridae; Chaerilidae; Buthidae I. Genera* Compsobuthus, Hottentotta, Isometrus, Lychas, *and* Sassanidotus. Prague: Clairon Production, 400 pp.
- LOURENÇO, W. R. 2000. The genus *Hottentotta* Birula, 1908, with the description of a new subgenus and species from India (Scorpiones, Buthidae). *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, 13(162): 191–195.
- PALLARY, P. 1938. Sur des scorpions de la Berbérie, de la Syrie et du Congo. *Archives de l'Institut Pasteur d'Algérie*, 16(3): 279–282.
- POCOCK, R. I. 1896. Report upon the scorpions, spiders, centipedes, and millipedes obtained by Mr. and Mrs. E. Lord Philips in the Goolis Mountains inland of Berbera, N. Somaliland. *Annals and Magazine of Natural History*, 6(18): 178–186.
- SOLEGLAD, M. E. & V. FET 2003. The scorpion sternum: structure and phylogeny (Scorpiones: Orthosterni). *Euscorpius*, 5: 1–34.

- STAHNKE, H. L. 1971. Scorpion nomenclature and mensuration. *Entomological News*, 81: 297–316.
- TIKADER, B. K. & D. B. BASTAWADE. 1983. Scorpions (Scorpionida: Arachnida). In *The Fauna of India*, Vol. 3. (Edited by the Director). Calcutta: Zoological Survey of India, 671pp.
- VACHON, M. 1952. Études sur les scorpions. *Institut Pasteur d'Algérie, Alger*, 1–482. (published 1948–1951 in *Archives de l'Institut Pasteur d'Algérie,* 1948, 26: 25–90, 162–208, 288–316, 441–481. 1949, 27: 66–100, 134–169, 281–288, 334–396. 1950, 28: 152–216, 383–413. 1951, 29: 46–104).
- VACHON, M. 1974. Études des caractères utilisés pour classer les familles et les genres des scorpions (Arachnides). 1. La trichobothriotaxie en arachnologie. Sigles trichobothriaux et types de trichobothriotaxie chez les Scorpions. *Bulletin du Muséum national d'Histoire naturelle*, 3e série, 140 (Zoologie, 104): 857–958.
- VACHON, M. 1979. Notes on the types of scorpions in the British Museum (Natural History), London Buthus socotrensis Pocock, 1889 (Family: Buthidae). Bulletin of the British Museum, Natural History (Zoology) 36(4): 233–237.