

CS2020A Discrete Mathematics

TUTORIAL 6 SUBMISSION

Submitted By

Kalakuntla Parjanya	112401018
Prachurjya Pratim Goswami	102401023
Chavva Srinivasa Saketh	112401009
Madeti Tarini	112401020
Vedant Singh	142401041

Problem 1. Convert the following statements in first order predicate logic to English and identify the famous theorem or conjecture they refer to. Domain of discourse is \mathbb{N}^+ , the set of positive natural numbers, P denotes the unary predicate “is prime”, $+$ denotes the binary function addition (in infix notation) and $>$ denotes the binary predicate “is greater than” (also in infix notation).

- a) $\forall x \forall y \neg (x^2 = 2y^2)$
- b) $\forall x \exists y ((y > x) \wedge P(y))$
- c) $\forall x \exists y ((y > x) \wedge (y \leq 2x) \wedge P(y))$
- d) $\forall x \exists y ((y > x) \wedge P(y) \wedge P(y + 2))$
- e) $\forall (x > 1) \exists y \exists z ((y + z = 2x) \wedge P(y) \wedge P(z))$
 $\equiv \forall x [(x > 1) \rightarrow \exists y \exists z ((y + z = 2x) \wedge P(y) \wedge P(z))]$

Solution.

- a) For all x and y belonging to positive numbers, x^2 cannot be equal to $2y^2$.
Theorem: $\sqrt{2}$ is irrational
- b) For all x , there exists some prime number y , such that y is greater than x .
Theorem: There exist infinite prime numbers.
- c) For all x , there exists some prime number y , such that y is greater than x , but less than $2x$.
Theorem: Bertrand's Postulate
- d) For all x , there exists some prime number y , such that $(y + 2)$ is a prime and y is greater than x .
Theorem: Twin Prime Conjecture
- e) For all x greater than 1, there exists some prime numbers y and z , such that $2x$ can be written as a sum of y and z .
Theorem: Goldbach's Conjecture

Problem 2. This is a “loose” definition of an algebraic structure called field.

A field is a set S with two operations (binary functions) ‘+’ (called addition) and ‘ \times ’ (called multiplication) which satisfies the following properties.

- Addition is commutative and associative.
- There exists an additive identity in S .
- Every member of S has an additive inverse.
- Multiplication is commutative and associative.
- There exists a multiplicative identity in S which is different from the additive identity.
- Every member of S , which is not the additive identity, has a multiplicative inverse.
- Multiplication distributes over addition.

Write a first order formula for each of the seven properties above.

Solution.

Here, $\mathbb{D} = S$

- $\forall x \forall y \forall z [(x + y = y + x) \wedge (x + (y + z) = (x + y) + z)]$ ✓
- $\exists y \forall x (x + y = y + x = x)$ ✓
- $\exists O \forall x \exists y ((x + O = O + x = x) \wedge (x + y = y + x = O))$ ✓
- $\forall x \forall y \forall z [(xy = yx) \wedge (x(yz) = (xy)z)]$ ✓
- $\forall x \exists y ((xy = yx = x) \wedge \neg(x + y = y + x = x))$ ✓
- $\forall x \exists y \exists z (\neg(x + y = y + x = x) \rightarrow (yz = zy = 1))$ ✗ $\rightarrow \forall x (x \neq \text{additive identity}) \rightarrow \exists y (xy = 1)$
- $\forall x \forall y \forall z (x(y + z) = xy + xz)$ ✓

Problem 3. Write a first order formula for the following predicates over natural numbers. You can use the predicates defined earlier in the later ones.

- a) x divides y . (Ans: $D(x, y) = \exists k(y = kx)$)
- b) x is prime. (Ans: $P(x) = \forall y(D(y, x) \rightarrow (y = 1) \vee (y = x))$).
- c) x is even.
- d) x is a perfect square.
- e) x has exactly three distinct prime factors.
- f) x and y are relatively prime.
- g) x and y have the same set of prime factors.

Solution.

- a) $D(x, y) = \forall x \forall y \exists k (y = kx)$ ✓
 - b) $P(x) = \forall x \forall y (D(x, y) \rightarrow ((y = 1) \vee (y = x)))$ ✓
 - c) $E(x) = \forall x (D(2, x))$ ✓
 - d) $S(x) = \forall x \exists y (x = y^2)$ ✓
 - e) $T(x) = \exists p \exists q \exists r \forall s [(P(p) \wedge P(q) \wedge P(r) \wedge P(s)) \wedge (D(p, x) \wedge D(q, x) \wedge D(r, x)) \wedge \neg(s = p) \wedge \neg(s = q) \wedge \neg(s = r)] \rightarrow (\neg D(s, x))$ ✓
 - f) $R(x, y) = \forall x \forall y \exists k [(P(x) \wedge P(y)) \vee (D(k, y) \wedge D(k, x) \rightarrow (k = 1))]$ ✓
 - g) $F(x) = \forall p \forall x \forall y [P(p) \wedge (D(p, x) \leftrightarrow D(p, y))]$ ✓
- only this is sufficient.

Problem 4. Convert the following statements into predicate logic formulae.

- a) If a prime number divides the product of two integers, then it divides at least one of them.
- b) Fermat's Last Theorem: No three positive integers a, b, c satisfy the equation $a^n + b^n = c^n$ for any integer $n > 2$.
- c) Every non-empty subset of natural numbers has a least element.
- d) Principle of Mathematical Induction

Solution.

- a) $\forall x \forall y \forall p (P(p) \wedge D(p, xy) \rightarrow (D(p, x) \vee D(p, y)))$ ✓
- b) $\forall a \forall b \forall c \forall n ((n > 2) \rightarrow \neg(a^n + b^n = c^n))$ ✓
- c) $\forall S (S \subseteq \mathbb{N} \wedge S \neq \emptyset \rightarrow \exists m \in S \forall x \in S m \leq x)$ ✓
- d) $\forall P (P(0) \wedge \forall n \in \mathbb{N} (P(n) \rightarrow P(n+1)) \rightarrow \forall n \in \mathbb{N} P(n))$ ✓

Problem 5. Let the domain of discourse $D = \{1, \dots, 10\}$. Let P be a 3-ary predicate defined on D . P can be represented as a binary 3D-Matrix M (a tensor), where $M(i, j, k) = 1$ if $P(i, j, k) = \text{True}$ and $M(i, j, k) = 0$ if $P(i, j, k) = \text{False}$. The statement “ M has an all-ones plane perpendicular to its first axis” is the first order sentence $\exists x \forall y \forall z P(x, y, z)$ (where x, y, z are domain variables). Write down similar first order sentences for the following statements.

- a) M has an all-zeros plane perpendicular to its second axis
- b) M has an all-ones line perpendicular to two of its three axes
- c) M has a zero in each plane perpendicular to its third axis
- d) M has at least two ones
- e) M has at least two ones in each plane perpendicular to its third axis
- f) Each plane perpendicular to the first axis in M represents a symmetric binary relation.

Solution.

- a) $\exists y \forall x \forall z \neg P(x, y, z)$
- b) $\exists x \exists y \forall z P(x, y, z) \vee \exists x \exists z \forall y P(x, y, z) \vee \exists y \exists z \forall x P(x, y, z)$
- c) $\forall z \exists x \exists y \neg P(x, y, z)$
- d) $\exists x_1 \exists y_1 \exists z_1 \exists x_2 \exists y_2 \exists z_2 \left(P(x_1, y_1, z_1) \wedge P(x_2, y_2, z_2) \wedge \neg(x_1 = x_2 \wedge y_1 = y_2 \wedge z_1 = z_2) \right)$
- e) $\forall z \exists x_1 \exists y_1 \exists x_2 \exists y_2 \left(P(x_1, y_1, z) \wedge P(x_2, y_2, z) \wedge \neg(x_1 = x_2 \wedge y_1 = y_2) \right)$
- f) $\forall x \forall y \forall z \left(P(x, y, z) \leftrightarrow P(x, z, y) \right)$