

SHEN COMIX

# Introduction to Optimization

K. R. Sahasranand

Data Science

sahasranand@iitpkd.ac.in

Surprise online test tomorrow at 6:00 am

Surprise online test tomorrow at 6:00 am



## Surprise online test tomorrow at 6:00 am

- Syllabus – modules I and II
- Time to prepare ~ 20 hours



## Surprise online test tomorrow at 6:00 am

- Syllabus – modules I and II
- Time to prepare ~ 20 hours
- Will have to drink 3 cups of coffee **every hour** for module I




## Surprise online test tomorrow at 6:00 am

- Syllabus – modules I and II
- Time to prepare ~ 20 hours
- Will have to drink 3 cups of coffee **every hour** for module I
- Will have to drink 4 cups of coffee **every hour** for module II



# Surprise online test tomorrow at 6:00 am

- Syllabus – modules I and II
- Time to prepare ~ 20 hours
- Will have to drink 3 cups of coffee **every hour** for module I
- Will have to drink 4 cups of coffee **every hour** for module II




Nobody:

Mom: *"Don't drink too much coffee..."*

# Surprise online test tomorrow at 6:00 am

- Syllabus – modules I and II
- Time to prepare ~ 20 hours
- Will have to drink 3 cups of coffee **every hour** for module I
- Will have to drink 4 cups of coffee **every hour** for module II




Nobody:

Mom: *"Don't drink too much coffee...  
But, since this test counts towards  
your CGPA, you may drink up to 72  
cups."*



# Surprise online test tomorrow at 6:00 am

- Syllabus – modules I and II
- Time to prepare ~ 20 hours
- Will have to drink 3 cups of coffee **every hour** for module I
- Will have to drink 4 cups of coffee **every hour** for module II




Nobody:

Mom: *"Don't drink too much coffee...  
But, since this test counts towards  
your CGPA, you may drink up to 72  
cups."*

- You score 4 points for every hour spent on module I  
and 5 points for every hour spent on module II

# Surprise online test tomorrow at 6:00 am

- Syllabus – modules I and II
- Time to prepare ~ 20 hours
- Will have to drink 3 cups of coffee **every hour** for module I
- Will have to drink 4 cups of coffee **every hour** for module II



Nobody:

Mom: *"Don't drink too much coffee...  
But, since this test counts towards  
your CGPA, you may drink up to 72  
cups."*

- You score 4 points for every hour spent on module I  
and 5 points for every hour spent on module II

How many hours for each module?

## Another example

maximize  $x_1 + x_2$

subject to  $x_2 - x_1 \leq 1$

$$x_1 + 6x_2 \leq 15$$

$$4x_1 - x_2 \leq 10$$

$$x_1 \geq 0, \quad x_2 \geq 0.$$

(continue on board)

## Another example

$$\begin{aligned} &\text{maximize } x_1 + x_2 \\ &\text{subject to } x_2 - x_1 \leq 1 \\ &\quad \quad \quad x_1 + 6x_2 \leq 15 \\ &\quad \quad \quad 4x_1 - x_2 \leq 10 \\ &\quad \quad \quad x_1 \geq 0, \ x_2 \geq 0. \end{aligned}$$

(continue on board)

To-do:

- show vector  $c$  and feasible region

## Another example

maximize  $x_1 + x_2$

subject to  $x_2 - x_1 \leq 1$

$$x_1 + 6x_2 \leq 15$$

$$4x_1 - x_2 \leq 10$$

$$x_1 \geq 0, \quad x_2 \geq 0.$$

(continue on board)

To-do:

- show vector  $c$  and feasible region

*unique solution*

## Another example

$$\begin{aligned} &\text{maximize } x_1 + x_2 \\ &\text{subject to } x_2 - x_1 \leq 1 \\ &\quad \quad \quad x_1 + 6x_2 \leq 15 \\ &\quad \quad \quad 4x_1 - x_2 \leq 10 \\ &\quad \quad \quad x_1 \geq 0, \quad x_2 \geq 0. \end{aligned}$$

(continue on board)

To-do:

- show vector  $c$  and feasible region
- replace  $c = \begin{bmatrix} 1 & 1 \end{bmatrix}^\top$  with  $\begin{bmatrix} \frac{1}{6} & 1 \end{bmatrix}^\top$

*unique solution*

## Another example

$$\begin{aligned} &\text{maximize } x_1 + x_2 \\ &\text{subject to } x_2 - x_1 \leq 1 \\ &\quad \quad \quad x_1 + 6x_2 \leq 15 \\ &\quad \quad \quad 4x_1 - x_2 \leq 10 \\ &\quad \quad \quad x_1 \geq 0, \quad x_2 \geq 0. \end{aligned}$$

(continue on board)

To-do:

- show vector  $c$  and feasible region

*unique solution*

- replace  $c = \begin{bmatrix} 1 & 1 \end{bmatrix}^\top$  with  $\begin{bmatrix} \frac{1}{6} & 1 \end{bmatrix}^\top$

*infinitely many solutions*

## Another example

$$\begin{aligned} &\text{maximize } x_1 + x_2 \\ &\text{subject to } x_2 - x_1 \leq 1 \\ &\quad \quad \quad x_1 + 6x_2 \leq 15 \\ &\quad \quad \quad 4x_1 - x_2 \leq 10 \\ &\quad \quad \quad x_1 \geq 0, \quad x_2 \geq 0. \end{aligned}$$

(continue on board)

To-do:

- show vector  $c$  and feasible region *unique solution*
- replace  $c = \begin{bmatrix} 1 & 1 \end{bmatrix}^\top$  with  $\begin{bmatrix} \frac{1}{6} & 1 \end{bmatrix}^\top$  *infinitely many solutions*
- reverse 1<sup>st</sup> and 3<sup>rd</sup>



## Another example

$$\begin{aligned} &\text{maximize } x_1 + x_2 \\ &\text{subject to } x_2 - x_1 \leq 1 \\ &\quad \quad \quad x_1 + 6x_2 \leq 15 \\ &\quad \quad \quad 4x_1 - x_2 \leq 10 \\ &\quad \quad \quad x_1 \geq 0, \quad x_2 \geq 0. \end{aligned}$$

(continue on board)

To-do:

- show vector  $c$  and feasible region *unique solution*
- replace  $c = \begin{bmatrix} 1 & 1 \end{bmatrix}^\top$  with  $\begin{bmatrix} \frac{1}{6} & 1 \end{bmatrix}^\top$  *infinitely many solutions*
- reverse 1<sup>st</sup> and 3<sup>rd</sup> *infeasible*

## Another example

$$\begin{aligned} &\text{maximize } x_1 + x_2 \\ &\text{subject to } x_2 - x_1 \leq 1 \\ &\quad \quad \quad x_1 + 6x_2 \leq 15 \\ &\quad \quad \quad 4x_1 - x_2 \leq 10 \\ &\quad \quad \quad x_1 \geq 0, \quad x_2 \geq 0. \end{aligned}$$

(continue on board)

To-do:

- show vector  $c$  and feasible region *unique solution*
- replace  $c = \begin{bmatrix} 1 & 1 \end{bmatrix}^\top$  with  $\begin{bmatrix} \frac{1}{6} & 1 \end{bmatrix}^\top$  *infinitely many solutions*
- reverse 1<sup>st</sup> and 3<sup>rd</sup> *infeasible*
- remove 2<sup>nd</sup> and 3<sup>rd</sup>

## Another example

$$\begin{aligned} &\text{maximize } x_1 + x_2 \\ &\text{subject to } x_2 - x_1 \leq 1 \\ &\quad \quad \quad x_1 + 6x_2 \leq 15 \\ &\quad \quad \quad 4x_1 - x_2 \leq 10 \\ &\quad \quad \quad x_1 \geq 0, \quad x_2 \geq 0. \end{aligned}$$


(continue on board)

To-do:

- show vector  $c$  and feasible region *unique solution*
- replace  $c = \begin{bmatrix} 1 & 1 \end{bmatrix}^\top$  with  $\begin{bmatrix} \frac{1}{6} & 1 \end{bmatrix}^\top$  *infinitely many solutions*
- reverse 1<sup>st</sup> and 3<sup>rd</sup> *infeasible*
- remove 2<sup>nd</sup> and 3<sup>rd</sup> *unbounded*


# Why it suffices to look at *corner points*?

$\max f(z)$  subject to some linear constraints where  $f(z) = c^\top z$




# Why it suffices to look at *corner points*?

$\max f(z)$  subject to some linear constraints where  $f(z) = c^\top z$



# Why it suffices to look at *corner points*?


$\max f(z)$  subject to some linear constraints where  $f(z) = c^\top z$



$$z^{(\alpha)} = \alpha z^{(a)} + (1 - \alpha) z^{(b)} \quad ; \quad 0 \leq \alpha \leq 1.$$

# Why it suffices to look at *corner points*?

$\max f(z)$  subject to some linear constraints where  $f(z) = c^\top z$



$$z^{(\alpha)} = \alpha z^{(a)} + (1 - \alpha) z^{(b)} \quad ; \quad 0 \leq \alpha \leq 1.$$


That is,

$$x_1^{(\alpha)} = \alpha x_1^{(a)} + (1 - \alpha) x_1^{(b)}$$

$$x_2^{(\alpha)} = \alpha x_2^{(a)} + (1 - \alpha) x_2^{(b)}$$

## Why it suffices to look at *corner points*?

$\max f(z)$  subject to some linear constraints where  $f(z) = c^\top z$



$$x_1^{(\alpha)} = \alpha x_1^{(a)} + (1 - \alpha)x_1^{(b)}$$

$$x_2^{(\alpha)} = \alpha x_2^{(a)} + (1 - \alpha)x_2^{(b)}$$

$$; 0 \leq \alpha \leq 1.$$

---


Objective function value at  $z^{(\alpha)}$ :

$$f(z^{(\alpha)}) = c^\top z^{(\alpha)}$$



# Why it suffices to look at *corner points*?

$\max f(z)$  subject to some linear constraints where  $f(z) = c^\top z$



$$x_1^{(\alpha)} = \alpha x_1^{(a)} + (1 - \alpha)x_1^{(b)}$$

$$x_2^{(\alpha)} = \alpha x_2^{(a)} + (1 - \alpha)x_2^{(b)}$$

$$; 0 \leq \alpha \leq 1.$$

---


Objective function value at  $z^{(\alpha)}$ :

$$f(z^{(\alpha)}) = c^\top z^{(\alpha)}$$

$$= c_1 x_1^{(\alpha)} + c_2 x_2^{(\alpha)}$$

## Why it suffices to look at *corner points*?

$\max f(z)$  subject to some linear constraints where  $f(z) = c^\top z$



$$x_1^{(\alpha)} = \alpha x_1^{(a)} + (1 - \alpha)x_1^{(b)}$$

$$x_2^{(\alpha)} = \alpha x_2^{(a)} + (1 - \alpha)x_2^{(b)}$$

$$; \quad 0 \leq \alpha \leq 1.$$

---

Objective function value at  $z^{(\alpha)}$ :


$$f(z^{(\alpha)}) = c^\top z^{(\alpha)}$$

$$= c_1 x_1^{(\alpha)} + c_2 x_2^{(\alpha)}$$

$$= c_1 \left( \alpha x_1^{(a)} + (1 - \alpha)x_1^{(b)} \right) + c_2 \left( \alpha x_2^{(a)} + (1 - \alpha)x_2^{(b)} \right)$$

## Why it suffices to look at *corner points*?

$\max f(z)$  subject to some linear constraints where  $f(z) = c^\top z$



$$x_1^{(\alpha)} = \alpha x_1^{(a)} + (1 - \alpha)x_1^{(b)}$$

$$x_2^{(\alpha)} = \alpha x_2^{(a)} + (1 - \alpha)x_2^{(b)}$$

$$; 0 \leq \alpha \leq 1.$$

---

Objective function value at  $z^{(\alpha)}$ :

$$f(z^{(\alpha)}) = c^\top z^{(\alpha)}$$


$$= c_1 x_1^{(\alpha)} + c_2 x_2^{(\alpha)}$$

$$= c_1 \left( \alpha x_1^{(a)} + (1 - \alpha)x_1^{(b)} \right) + c_2 \left( \alpha x_2^{(a)} + (1 - \alpha)x_2^{(b)} \right)$$

$$= \alpha \left( c_1 x_1^{(a)} + c_2 x_2^{(a)} \right) + (1 - \alpha) \left( c_1 x_1^{(b)} + c_2 x_2^{(b)} \right)$$

## Why it suffices to look at *corner points*?

$\max f(z)$  subject to some linear constraints where  $f(z) = c^\top z$



$$x_1^{(\alpha)} = \alpha x_1^{(a)} + (1 - \alpha)x_1^{(b)}$$

$$x_2^{(\alpha)} = \alpha x_2^{(a)} + (1 - \alpha)x_2^{(b)}$$

$$; 0 \leq \alpha \leq 1.$$

---

Objective function value at  $z^{(\alpha)}$ :

$$f(z^{(\alpha)}) = c^\top z^{(\alpha)}$$

$$= c_1 x_1^{(\alpha)} + c_2 x_2^{(\alpha)}$$

$$= c_1 \left( \alpha x_1^{(a)} + (1 - \alpha)x_1^{(b)} \right) + c_2 \left( \alpha x_2^{(a)} + (1 - \alpha)x_2^{(b)} \right)$$

$$= \alpha \left( c_1 x_1^{(a)} + c_2 x_2^{(a)} \right) + (1 - \alpha) \left( c_1 x_1^{(b)} + c_2 x_2^{(b)} \right)$$

$$= \alpha f(z^{(a)}) + (1 - \alpha)f(z^{(b)})$$

# Top 10 algorithms from the 20<sup>th</sup> century

*Computing in Science & Engineering*, American Institute of Physics and the IEEE Computer Society, Jan 2000.



# Simplex algorithm – bird's eye view

*Algorithms*, Dasgupta-Papadimitriou-Vazirani [online]

Let  $v$  be any vertex of the feasible region.

while  $\exists$  a neighbor  $v'$  of  $v$  with better objective value:

    set  $v = v'$

# Simplex algorithm – bird's eye view

*Algorithms*, Dasgupta-Papadimitriou-Vazirani [online]

Let  $v$  be any vertex of the feasible region.

while  $\exists$  a neighbor  $v'$  of  $v$  with better objective value:  
    set  $v = v'$

On each iteration, the algorithm has **two** tasks:

# Simplex algorithm – bird's eye view

*Algorithms*, Dasgupta-Papadimitriou-Vazirani [online]

Let  $v$  be any vertex of the feasible region.

while  $\exists$  a neighbor  $v'$  of  $v$  with better objective value:

    set  $v = v'$

On each iteration, the algorithm has **two** tasks:

1. Check whether the current vertex is optimal (and if so, halt).



# Simplex algorithm – bird's eye view

*Algorithms*, Dasgupta-Papadimitriou-Vazirani [online]

Let  $v$  be any vertex of the feasible region.

while  $\exists$  a neighbor  $v'$  of  $v$  with better objective value:

    set  $v = v'$

On each iteration, the algorithm has **two** tasks:

1. Check whether the current vertex is optimal (and if so, halt).
2. Determine where to move next.

# Simplex algorithm – bird's eye view

*Algorithms*, Dasgupta-Papadimitriou-Vazirani [online]

Let  $v$  be any vertex of the feasible region.

while  $\exists$  a neighbor  $v'$  of  $v$  with better objective value:

    set  $v = v'$

On each iteration, the algorithm has **two** tasks:

1. Check whether the current vertex is optimal (and if so, halt).
2. Determine where to move next.

Both tasks are easy if the vertex is at the origin!

# Simplex algorithm – bird's eye view

*Algorithms*, Dasgupta-Papadimitriou-Vazirani [online]

Let  $v$  be any vertex of the feasible region.

while  $\exists$  a neighbor  $v'$  of  $v$  with better objective value:

    set  $v = v'$

On each iteration, the algorithm has **two** tasks:

1. Check whether the current vertex is optimal (and if so, halt).
2. Determine where to move next.

Both tasks are easy if the vertex is at the origin!

But what if our current vertex  $u$  is elsewhere?

# Simplex algorithm – bird's eye view

*Algorithms*, Dasgupta-Papadimitriou-Vazirani [online]

Let  $v$  be any vertex of the feasible region.

while  $\exists$  a neighbor  $v'$  of  $v$  with better objective value:

set  $v = v'$

On each iteration, the algorithm has **two** tasks:

1. Check whether the current vertex is optimal (and if so, halt).
2. Determine where to move next.

Both tasks are easy if the vertex is at the origin!

But what if our current vertex  $u$  is elsewhere?



*Transform  $u$  into the origin, by shifting the coordinate system from the usual  $(x_1, x_2, \dots, x_n)$  to the “local view” from  $u$ .*

# The two tasks at the origin

*"Both tasks are easy if the vertex is at the origin!"*

On each iteration, the algorithm has **two** tasks:

1. Check whether the current vertex is optimal (and if so, halt).
2. Determine where to move next.

Consider the LP:

$$\text{maximize } c^\top x \text{ subject to } Ax \geq b, x \geq 0 \quad ; \quad x \in \mathbb{R}^n.$$

# The two tasks at the origin

*"Both tasks are easy if the vertex is at the origin!"*

On each iteration, the algorithm has **two** tasks:

1. Check whether the current vertex is optimal (and if so, halt).
2. Determine where to move next.

Consider the LP:

$$\text{maximize } c^\top x \text{ subject to } Ax \geq b, x \geq 0 \quad ; \quad x \in \mathbb{R}^n.$$

Origin is feasible  $\implies$  Origin is a vertex

since it is the unique point at which the  $n$  inequalities

$$x_1 \geq 0, \quad x_2 \geq 0, \quad \dots, \quad x_n \geq 0 \quad \text{are tight.}$$

# The two tasks at the origin

*"Both tasks are easy if the vertex is at the origin!"*

On each iteration, the algorithm has **two** tasks:

1. Check whether the current vertex is optimal (and if so, halt).
2. Determine where to move next.

Consider the LP:

$$\text{maximize } c^\top x \text{ subject to } Ax \geq b, x \geq 0 \quad ; x \in \mathbb{R}^n.$$

Origin is feasible  $\implies$  Origin is a vertex

since it is the unique point at which the  $n$  inequalities

$$x_1 \geq 0, x_2 \geq 0, \dots, x_n \geq 0 \quad \text{are tight.}$$

The origin is optimal if and only if all  $c_i \leq 0$ .

# The two tasks at the origin

*"Both tasks are easy if the vertex is at the origin!"*

On each iteration, the algorithm has **two** tasks:

1. Check whether the current vertex is optimal (and if so, halt).
2. Determine where to move next.

Consider the LP:

$$\text{maximize } c^T x \text{ subject to } Ax \geq b, x \geq 0 \quad ; \quad x \in \mathbb{R}^n.$$

Origin is feasible  $\implies$  Origin is a vertex

since it is the unique point at which the  $n$  inequalities

$$x_1 \geq 0, \quad x_2 \geq 0, \quad \dots, \quad x_n \geq 0 \quad \text{are tight.}$$

The origin is optimal if and only if all  $c_i \leq 0$ .

Increase some  $x_i$  for which  $c_i > 0$ .

Increase by how much? Until we hit some other constraint.



# Solve using simplex algorithm

*“always operate at the **origin**”*

max  $2x_1 + 5x_2$  subject to

$$2x_1 - x_2 \leq 4 \quad (1)$$

$$x_1 + 2x_2 \leq 9 \quad (2)$$

$$-x_1 + x_2 \leq 3 \quad (3)$$

$$x_1 \geq 0 \quad (4)$$

$$x_2 \geq 0 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(5)

**Objective value:**

max  $2x_1 + 5x_2$  subject to

$$2x_1 - x_2 \leq 4 \quad (1)$$

$$x_1 + 2x_2 \leq 9 \quad (2)$$

$$-x_1 + x_2 \leq 3 \quad (3)$$

$$x_1 \geq 0 \quad (4)$$

$$x_2 \geq 0 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(5)

**Objective value:** 0

max  $2x_1 + 5x_2$  subject to

$$2x_1 - x_2 \leq 4 \quad (1)$$

$$x_1 + 2x_2 \leq 9 \quad (2)$$

$$-x_1 + x_2 \leq 3 \quad (3)$$

$$x_1 \geq 0 \quad (4)$$

$$x_2 \geq 0 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(5)

**Objective value:** 0

**Move:**

max  $2x_1 + 5x_2$  subject to

$$2x_1 - x_2 \leq 4 \quad (1)$$

$$x_1 + 2x_2 \leq 9 \quad (2)$$

$$-x_1 + x_2 \leq 3 \quad (3)$$

$$x_1 \geq 0 \quad (4)$$

$$x_2 \geq 0 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(5)

**Objective value:** 0

**Move:** increase  $x_2$

max  $2x_1 + 5x_2$  subject to

$$2x_1 - x_2 \leq 4 \quad (1)$$

$$x_1 + 2x_2 \leq 9 \quad (2)$$

$$-x_1 + x_2 \leq 3 \quad (3)$$

$$x_1 \geq 0 \quad (4)$$

$$x_2 \geq 0 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(5)

**Objective value:** 0

**Move:** increase  $x_2$

(5) is released, (3) is tightened

max  $2x_1 + 5x_2$  subject to

$$2x_1 - x_2 \leq 4 \quad (1)$$

$$x_1 + 2x_2 \leq 9 \quad (2)$$

$$-x_1 + x_2 \leq 3 \quad (3)$$

$$x_1 \geq 0 \quad (4)$$

$$x_2 \geq 0 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(5)

**Objective value:** 0

**Move:** increase  $x_2$

STOP at  $x_2 = ?$

(5) is released, (3) is tightened
-----------------------------------

max  $2x_1 + 5x_2$  subject to

$$2x_1 - x_2 \leq 4 \quad (1)$$

$$x_1 + 2x_2 \leq 9 \quad (2)$$

$$-x_1 + x_2 \leq 3 \quad (3)$$

$$x_1 \geq 0 \quad (4)$$

$$x_2 \geq 0 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(5)

**Objective value:** 0

**Move:** increase  $x_2$

STOP at  $x_2 = 3$

max  $2x_1 + 5x_2$  subject to

$$2x_1 - x_2 \leq 4 \quad (1)$$

$$x_1 + 2x_2 \leq 9 \quad (2)$$

$$-x_1 + x_2 \leq 3 \quad (3)$$

$$x_1 \geq 0 \quad (4)$$

$$x_2 \geq 0 \quad (5)$$



# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:**  $(4), (5)$

**Objective value:** 0

**Move:** increase  $x_2$

STOP at  $x_2 = 3$

<b>New vertex:</b> $(4), (3)$
-------------------------------

max  $2x_1 + 5x_2$  subject to

$$2x_1 - x_2 \leq 4 \quad (1)$$

$$x_1 + 2x_2 \leq 9 \quad (2)$$

$$-x_1 + x_2 \leq 3 \quad (3)$$

$$x_1 \geq 0 \quad (4)$$

$$x_2 \geq 0 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(5)

**Objective value:** 0

**Move:** increase  $x_2$

STOP at  $x_2 = 3$

**New vertex:** (4),(3)

$$y_1 = x_1$$

$$y_2 = 3 + x_1 - x_2$$

max  $2x_1 + 5x_2$  subject to

$$2x_1 - x_2 \leq 4 \quad (1)$$

$$x_1 + 2x_2 \leq 9 \quad (2)$$

$$-x_1 + x_2 \leq 3 \quad (3)$$

$$x_1 \geq 0 \quad (4)$$

$$x_2 \geq 0 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(3)

**Objective value:**

max  $15 + 7y_1 - 5y_2$  subject to

$$y_1 + y_2 \leq 7 \quad (1)$$

$$3y_1 - 2y_2 \leq 3 \quad (2)$$

$$y_2 \geq 0 \quad (3)$$

$$y_1 \geq 0 \quad (4)$$

$$-y_1 + y_2 \leq 3 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(3)

**Objective value:** 15

max  $15 + 7y_1 - 5y_2$  subject to

$$y_1 + y_2 \leq 7 \quad (1)$$

$$3y_1 - 2y_2 \leq 3 \quad (2)$$

$$y_2 \geq 0 \quad (3)$$

$$y_1 \geq 0 \quad (4)$$

$$-y_1 + y_2 \leq 3 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(3)

**Objective value:** 15

**Move:**

max  $15 + 7y_1 - 5y_2$  subject to

$$y_1 + y_2 \leq 7 \quad (1)$$

$$3y_1 - 2y_2 \leq 3 \quad (2)$$

$$y_2 \geq 0 \quad (3)$$

$$y_1 \geq 0 \quad (4)$$

$$-y_1 + y_2 \leq 3 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(3)

**Objective value:** 15

**Move:** increase  $y_1$

max  $15 + 7y_1 - 5y_2$  subject to

$$y_1 + y_2 \leq 7 \quad (1)$$

$$3y_1 - 2y_2 \leq 3 \quad (2)$$

$$y_2 \geq 0 \quad (3)$$

$$y_1 \geq 0 \quad (4)$$

$$-y_1 + y_2 \leq 3 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(3)

**Objective value:** 15

**Move:** increase  $y_1$

(4) is released, (2) is tightened

max  $15 + 7y_1 - 5y_2$  subject to

$$y_1 + y_2 \leq 7 \quad (1)$$

$$3y_1 - 2y_2 \leq 3 \quad (2)$$

$$y_2 \geq 0 \quad (3)$$

$$y_1 \geq 0 \quad (4)$$

$$-y_1 + y_2 \leq 3 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(3)

**Objective value:** 15

**Move:** increase  $y_1$

STOP at  $y_1 = ?$

(4) is released, (2) is tightened
-----------------------------------

max  $15 + 7y_1 - 5y_2$  subject to

$$y_1 + y_2 \leq 7 \quad (1)$$

$$3y_1 - 2y_2 \leq 3 \quad (2)$$

$$y_2 \geq 0 \quad (3)$$

$$y_1 \geq 0 \quad (4)$$

$$-y_1 + y_2 \leq 3 \quad (5)$$



# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (4),(3)

**Objective value:** 15

**Move:** increase  $y_1$

STOP at  $y_1 = 1$

max  $15 + 7y_1 - 5y_2$  subject to

$$y_1 + y_2 \leq 7 \quad (1)$$

$$3y_1 - 2y_2 \leq 3 \quad (2)$$

$$y_2 \geq 0 \quad (3)$$

$$y_1 \geq 0 \quad (4)$$

$$-y_1 + y_2 \leq 3 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:**  $(4), (3)$

**Objective value:** 15

**Move:** increase  $y_1$

STOP at  $y_1 = 1$

**New vertex:**  $(2), (3)$

max  $15 + 7y_1 - 5y_2$  subject to

$$y_1 + y_2 \leq 7 \quad (1)$$

$$3y_1 - 2y_2 \leq 3 \quad (2)$$

$$y_2 \geq 0 \quad (3)$$

$$y_1 \geq 0 \quad (4)$$

$$-y_1 + y_2 \leq 3 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:**  $(4), (3)$

**Objective value:**  $15$

**Move:** increase  $y_1$

STOP at  $y_1 = 1$

**New vertex:**  $(2), (3)$

$$z_1 = 3 - 3y_1 + 2y_2$$

$$z_2 = y_2$$

max  $15 + 7y_1 - 5y_2$  subject to

$$y_1 + y_2 \leq 7 \quad (1)$$

$$3y_1 - 2y_2 \leq 3 \quad (2)$$

$$y_2 \geq 0 \quad (3)$$

$$y_1 \geq 0 \quad (4)$$

$$-y_1 + y_2 \leq 3 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (2),(3)

**Objective value:**

$$\max 22 - \frac{7}{3}z_1 - \frac{1}{3}z_2 \text{ subject to}$$

$$-\frac{1}{3}z_1 + \frac{5}{3}z_2 \leq 6 \quad (1)$$

$$z_1 \geq 0 \quad (2)$$

$$z_2 \geq 0 \quad (3)$$

$$\frac{1}{3}z_1 - \frac{2}{3}z_2 \leq 1 \quad (4)$$

$$\frac{1}{3}z_1 + \frac{1}{3}z_2 \leq 4 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (2),(3)

**Objective value:** 22

$$\max 22 - \frac{7}{3}z_1 - \frac{1}{3}z_2 \text{ subject to}$$

$$-\frac{1}{3}z_1 + \frac{5}{3}z_2 \leq 6 \quad (1)$$

$$z_1 \geq 0 \quad (2)$$

$$z_2 \geq 0 \quad (3)$$

$$\frac{1}{3}z_1 - \frac{2}{3}z_2 \leq 1 \quad (4)$$

$$\frac{1}{3}z_1 + \frac{1}{3}z_2 \leq 4 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (2),(3)

**Objective value:** 22

**Move:**

$$\max 22 - \frac{7}{3}z_1 - \frac{1}{3}z_2 \text{ subject to}$$

$$-\frac{1}{3}z_1 + \frac{5}{3}z_2 \leq 6 \quad (1)$$

$$z_1 \geq 0 \quad (2)$$

$$z_2 \geq 0 \quad (3)$$

$$\frac{1}{3}z_1 - \frac{2}{3}z_2 \leq 1 \quad (4)$$

$$\frac{1}{3}z_1 + \frac{1}{3}z_2 \leq 4 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

Current vertex:

Objective value: 22

Move:



max  $22 - \frac{1}{3}z_1 - \frac{1}{3}z_2$  subject to

$$-\frac{1}{3}z_1 + \frac{5}{3}z_2 \leq 6 \quad (1)$$

$$z_1 \geq 0 \quad (2)$$

$$z_2 \geq 0 \quad (3)$$

$$\frac{1}{3}z_1 - \frac{2}{3}z_2 \leq 1 \quad (4)$$

$$\frac{1}{3}z_1 + \frac{1}{3}z_2 \leq 4 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

Current vertex: **(2),(3)**

Objective value: **22**

Move: **origin is optimal!**

(all  $c_i < 0$ )

$$\max \quad 22 - \frac{7}{3}z_1 - \frac{1}{3}z_2 \quad \text{subject to}$$

$$-\frac{1}{3}z_1 + \frac{5}{3}z_2 \leq 6 \quad (1)$$

$$z_1 \geq 0 \quad (2)$$

$$z_2 \geq 0 \quad (3)$$

$$\frac{1}{3}z_1 - \frac{2}{3}z_2 \leq 1 \quad (4)$$

$$\frac{1}{3}z_1 + \frac{1}{3}z_2 \leq 4 \quad (5)$$



# Solve using simplex algorithm

*"always operate at the **origin**"*

Current vertex: **(2),(3)**

Objective value: 22

Move: **origin is optimal!**

(all  $c_i < 0$ )

$$\max \quad 22 - \frac{7}{3}z_1 - \frac{1}{3}z_2 \quad \text{subject to}$$

$$-\frac{1}{3}z_1 + \frac{5}{3}z_2 \leq 6 \quad (1)$$

$$z_1 \geq 0 \quad (2)$$

$$z_2 \geq 0 \quad (3)$$

$$\frac{1}{3}z_1 - \frac{2}{3}z_2 \leq 1 \quad (4)$$

$$\frac{1}{3}z_1 + \frac{1}{3}z_2 \leq 4 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

**Current vertex:** (2),(3)

**Objective value:** 22

**Move:** origin is optimal!  
(all  $c_i < 0$ )

Solve (2),(3) in the original LP:

$$\max 22 - \frac{7}{3}z_1 - \frac{1}{3}z_2 \text{ subject to}$$

$$-\frac{1}{3}z_1 + \frac{5}{3}z_2 \leq 6 \quad (1)$$

$$z_1 \geq 0 \quad (2)$$

$$z_2 \geq 0 \quad (3)$$

$$\frac{1}{3}z_1 - \frac{2}{3}z_2 \leq 1 \quad (4)$$

$$\frac{1}{3}z_1 + \frac{1}{3}z_2 \leq 4 \quad (5)$$

# Solve using simplex algorithm

*"always operate at the **origin**"*

Current vertex: **(2),(3)**

Objective value: 22

Move: **origin is optimal!**  
(all  $c_i < 0$ )

Solve **(2),(3)** in the original LP:

$(x_1, x_2) = (1, 4)$

$$\max 22 - \frac{7}{3}z_1 - \frac{1}{3}z_2 \text{ subject to}$$

$$-\frac{1}{3}z_1 + \frac{5}{3}z_2 \leq 6 \quad (1)$$

$$z_1 \geq 0 \quad (2)$$


$$z_2 \geq 0 \quad (3)$$

$$\frac{1}{3}z_1 - \frac{2}{3}z_2 \leq 1 \quad (4)$$

$$\frac{1}{3}z_1 + \frac{1}{3}z_2 \leq 4 \quad (5)$$

# Back to the first example

solve using simplex



$$\begin{aligned} &\text{maximize } 4x_1 + 5x_2 \\ &\text{subject to } x_1 + x_2 \leq 20 \\ &\quad \quad \quad 3x_1 + 4x_2 \leq 72 \\ &\quad \quad \quad x_1 \geq 0 \\ &\quad \quad \quad x_2 \geq 0 \end{aligned}$$




(complete on board)

# Another interesting example


Unbounded LP

$$\begin{array}{ll}\text{maximize } x_1 & \\ \text{subject to } x_1 - x_2 \leq 1 & (1) \\ -x_1 + x_2 \leq 2 & (2) \\ x_1 \geq 0 & (3) \\ x_2 \geq 0 & (4)\end{array}$$



# Another interesting example

Unbounded LP



maximize  $x_1$

subject to  $x_1 - x_2 \leq 1$  (1)

$-x_1 + x_2 \leq 2$  (2)

$x_1 \geq 0$  (3)


$x_2 \geq 0$  (4)

**Current vertex:** (3),(4)

**Objective value:** 0

# Another interesting example

Unbounded LP



maximize  $x_1$

subject to  $x_1 - x_2 \leq 1$  (1)

$-x_1 + x_2 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)


**Current vertex:** (3),(4)

**Objective value:** 0

**Move:** increase  $x_1$

# Another interesting example

Unbounded LP



maximize  $x_1$

subject to  $x_1 - x_2 \leq 1$  (1)

$-x_1 + x_2 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

**Current vertex:** (3),(4)

**Objective value:** 0


**Move:** increase  $x_1$

STOP at  $x_1 = 1$



# Another interesting example

Unbounded LP



maximize  $x_1$

subject to  $x_1 - x_2 \leq 1$  (1)

$-x_1 + x_2 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

**Current vertex:** (3),(4)

**Objective value:** 0


**Move:** increase  $x_1$

STOP at  $x_1 = 1$

**New vertex:** (1),(4)

# Another interesting example

Unbounded LP



maximize  $x_1$

subject to  $x_1 - x_2 \leq 1$  (1)

$-x_1 + x_2 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_1$

STOP at  $x_1 = 1$


New vertex: (1),(4)

$$y_2 = x_2$$

$$y_1 = 1 - x_1 + x_2$$

# Another interesting example

Unbounded LP



maximize  $x_1$

subject to  $x_1 - x_2 \leq 1$  (1)

$-x_1 + x_2 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_1$

STOP at  $x_1 = 1$

New vertex: (1),(4)

$$y_2 = x_2$$

$$y_1 = 1 - x_1 + x_2$$

---

maximize  $1 - y_1 + y_2$

subject to  $y_1 \geq 0$  (1)

$y_1 \leq 3$  (2)


$y_1 - y_2 \leq 1$  (3)

$y_2 \geq 0$  (4)

Current vertex: (1),(4)

# Another interesting example

Unbounded LP



maximize  $x_1$

subject to  $x_1 - x_2 \leq 1$  (1)

$-x_1 + x_2 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_1$

STOP at  $x_1 = 1$

New vertex: (1),(4)

$$y_2 = x_2$$

$$y_1 = 1 - x_1 + x_2$$

---

maximize  $1 - y_1 + y_2$

subject to  $y_1 \geq 0$  (1)

$y_1 \leq 3$  (2)

$y_1 - y_2 \leq 1$  (3)


$y_2 \geq 0$  (4)

Current vertex: (1),(4)

Objective value: 1

# Another interesting example

Unbounded LP



maximize  $x_1$

subject to  $x_1 - x_2 \leq 1$  (1)

$-x_1 + x_2 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_1$

STOP at  $x_1 = 1$

New vertex: (1),(4)

$$y_2 = x_2$$

$$y_1 = 1 - x_1 + x_2$$

---

maximize  $1 - y_1 + y_2$

subject to  $y_1 \geq 0$  (1)

$y_1 \leq 3$  (2)

$y_1 - y_2 \leq 1$  (3)

$y_2 \geq 0$  (4)


Current vertex: (1),(4)

Objective value: 1

Move: increase  $y_2$

# Another interesting example

Unbounded LP



maximize  $x_1$

subject to  $x_1 - x_2 \leq 1$  (1)

$-x_1 + x_2 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_1$

STOP at  $x_1 = 1$

New vertex: (1),(4)

$$y_2 = x_2$$

$$y_1 = 1 - x_1 + x_2$$

---

maximize  $1 - y_1 + y_2$

subject to  $y_1 \geq 0$  (1)

$y_1 \leq 3$  (2)

$y_1 - y_2 \leq 1$  (3)

$y_2 \geq 0$  (4)

Current vertex: (1),(4)


Objective value: 1

Move: increase  $y_2$

STOP at  $y_2 = ?$

# Another interesting example

Unbounded LP



maximize  $x_1$

subject to  $x_1 - x_2 \leq 1$  (1)

$-x_1 + x_2 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_1$

STOP at  $x_1 = 1$

New vertex: (1),(4)

$y_2 = x_2$

$y_1 = 1 - x_1 + x_2$

---

maximize  $1 - y_1 + y_2$

subject to  $y_1 \geq 0$  (1)

$y_1 \leq 3$  (2)

$y_1 - y_2 \leq 1$  (3)

$y_2 \geq 0$  (4)

Current vertex: (1),(4)

Objective value: 1

Move: increase  $y_2$

STOP at  $y_2 = ?$

LP is unbounded

# One last example

## Degeneracy

maximize  $x_2$

subject to  $-x_1 + x_2 \leq 0$  (1)

$x_1 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

**Current vertex:** (3),(4)



# One last example

## Degeneracy

maximize  $x_2$

subject to  $-x_1 + x_2 \leq 0$  (1)

$x_1 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

Current vertex: (3),(4)

Objective value: 0

# One last example

## Degeneracy

maximize  $x_2$

subject to  $-x_1 + x_2 \leq 0$  (1)

$x_1 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)

**Current vertex:** (3),(4)

**Objective value:** 0

**Move:** increase  $x_2$

# One last example

## Degeneracy

$$\begin{array}{ll}\text{maximize } x_2 & \\ \text{subject to } -x_1 + x_2 \leq 0 & (1) \\ x_1 \leq 2 & (2) \\ x_1 \geq 0 & (3) \\ x_2 \geq 0 & (4)\end{array}$$

Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_2$



# One last example

## Degeneracy

$$\begin{array}{ll}\text{maximize } x_2 & \\ \text{subject to } -x_1 + x_2 \leq 0 & (1) \\ x_1 \leq 2 & (2) \\ x_1 \geq 0 & (3) \\ x_2 \geq 0 & (4)\end{array}$$

Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_2$



CAN'T!

# One last example

## Degeneracy

$$\begin{array}{ll}\text{maximize } x_2 & \\ \text{subject to } -x_1 + x_2 \leq 0 & (1) \\ x_1 \leq 2 & (2) \\ x_1 \geq 0 & (3) \\ x_2 \geq 0 & (4)\end{array}$$

Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_2$



CAN'T!

# One last example

## Degeneracy

maximize  $x_2$

subject to  $-x_1 + x_2 \leq 0$  (1)

$x_1 \leq 2$  (2)

$x_1 \geq 0$  (3)

$x_2 \geq 0$  (4)


Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_2$



CAN'T!



# One last example

## Degeneracy

- maximize  $x_2$
- subject to  $-x_1 + x_2 \leq 0$  (1)
- $x_1 \leq 2$  (2)
- $x_1 \geq 0$  (3)
- $x_2 \geq 0$  (4)


Current vertex: (3),(4)

Objective value: 0

Move: increase  $x_2$



CAN'T!



Move: increase  $x_1 \rightarrow$

# LP in standard form

simplex tableau demands . . .

$$\begin{aligned} & \text{minimize } c^\top x \\ & \text{subject to } Ax = b, \\ & \quad x \geq 0. \end{aligned}$$



## LP in standard form

simplex tableau demands . . .

$$\begin{aligned} &\text{minimize } c^\top x \\ &\text{subject to } Ax = b, \\ &\quad x \geq 0. \end{aligned}$$

Suppose we are given the inequality constraint

$$x_1 \leq 7.$$

## LP in standard form

simplex tableau demands . . .

$$\begin{aligned} &\text{minimize } c^\top x \\ &\text{subject to } Ax = b, \\ &\quad x \geq 0. \end{aligned}$$

Suppose we are given the inequality constraint

$$x_1 \leq 7.$$

Convert to an **equality constraint** – introduce a **slack variable**:

$$\begin{aligned} x_1 + s_1 &= 7 \\ s_1 &\geq 0. \end{aligned}$$