



Introduction to Optimization

K. R. Sahasranand

Data Science

sahasranand@iitpkd.ac.in

Full-rank factorization

a matrix of rank r = a matrix of full column rank r \times a matrix of full row rank r .

Full-rank factorization

a matrix of rank r = a matrix of full column rank r \times a matrix of full row rank r .

Helper lemma – Let $A \in \mathbb{R}^{m \times n}$, $\text{rank } A = r \leq \min\{m, n\}$. Then, there exist matrices $B \in \mathbb{R}^{m \times r}$ and $C \in \mathbb{R}^{r \times n}$ such that

$$A = BC$$

with

$$\text{rank } A = \text{rank } B = \text{rank } C = r.$$

Full-rank factorization

a matrix of rank r = a matrix of full column rank r \times a matrix of full row rank r .

Helper lemma – Let $A \in \mathbb{R}^{m \times n}$, $\text{rank } A = r \leq \min\{m, n\}$. Then, there exist matrices $B \in \mathbb{R}^{m \times r}$ and $C \in \mathbb{R}^{r \times n}$ such that

$$A = BC$$

with

$$\text{rank } A = \text{rank } B = \text{rank } C = r.$$

Proof – WLOG, a_1, \dots, a_r (columns of A) be linearly independent

Full-rank factorization

a matrix of rank r = a matrix of full column rank r \times a matrix of full row rank r .

Helper lemma – Let $A \in \mathbb{R}^{m \times n}$, $\text{rank } A = r \leq \min\{m, n\}$. Then, there exist matrices $B \in \mathbb{R}^{m \times r}$ and $C \in \mathbb{R}^{r \times n}$ such that

$$A = BC$$

with

$$\text{rank } A = \text{rank } B = \text{rank } C = r.$$

Proof – WLOG, a_1, \dots, a_r (columns of A) be linearly independent

$$B = \begin{bmatrix} | & | & | \\ a_1 & \cdots & a_r \\ | & | & | \end{bmatrix}$$

Full-rank factorization

a matrix of rank r = a matrix of full column rank r \times a matrix of full row rank r .

Helper lemma – Let $A \in \mathbb{R}^{m \times n}$, $\text{rank } A = r \leq \min\{m, n\}$. Then, there exist matrices $B \in \mathbb{R}^{m \times r}$ and $C \in \mathbb{R}^{r \times n}$ such that

$$A = BC$$

with

$$\text{rank } A = \text{rank } B = \text{rank } C = r.$$

Proof – WLOG, a_1, \dots, a_r (columns of A) be linearly independent

$$B = \begin{bmatrix} | & & | \\ a_1 & \cdots & a_r \\ | & & | \end{bmatrix} \quad ; \quad C = \begin{bmatrix} 1 & \cdots & 0 & c_{1,r+1} & \cdots & c_{1,n} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 1 & c_{m,r+1} & \cdots & c_{m,n} \end{bmatrix}$$

An example

$$A = \begin{bmatrix} 2 & 1 & -2 & 5 \\ 1 & 0 & -3 & 2 \\ 3 & -1 & -13 & 5 \end{bmatrix}$$

rank $A = ?$

An example

$$A = \begin{bmatrix} 2 & 1 & -2 & 5 \\ 1 & 0 & -3 & 2 \\ 3 & -1 & -13 & 5 \end{bmatrix}$$

$$\text{rank } A = 2$$

An example

$$A = \begin{bmatrix} 2 & 1 & -2 & 5 \\ 1 & 0 & -3 & 2 \\ 3 & -1 & -13 & 5 \end{bmatrix}$$

$$\text{rank } A = 2$$

$$B = \begin{bmatrix} 2 & 1 \\ 1 & 0 \\ 3 & -1 \end{bmatrix}$$

An example

$$A = \begin{bmatrix} 2 & 1 & -2 & 5 \\ 1 & 0 & -3 & 2 \\ 3 & -1 & -13 & 5 \end{bmatrix}$$

$$\text{rank } A = 2$$

$$B = \begin{bmatrix} 2 & 1 \\ 1 & 0 \\ 3 & -1 \end{bmatrix} \quad ; \quad C = \begin{bmatrix} 1 & 0 & -3 & 2 \\ 0 & 1 & 4 & 1 \end{bmatrix}$$

Generalized inverse

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times m}$ is called a **generalized inverse** of A if

$$AA^\dagger A = A$$

and there exist matrices $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ such that

$$A^\dagger = UA^\top \quad \text{and} \quad A^\dagger = A^\top V.$$

Generalized inverse

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times n}$ is called a **generalized inverse** of A if

$$AA^\dagger A = A$$

and there exist matrices $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ such that

$$A^\dagger = UA^\top \quad \text{and} \quad A^\dagger = A^\top V.$$

Tall – $A \in \mathbb{R}^{m \times n}, m \geq n, \text{rank } A = n$: $A^\dagger = (A^\top A)^{-1}A^\top$

Generalized inverse

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times m}$ is called a **generalized inverse** of A if

$$AA^\dagger A = A$$

and there exist matrices $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ such that

$$A^\dagger = UA^\top \quad \text{and} \quad A^\dagger = A^\top V.$$

Tall – $A \in \mathbb{R}^{m \times n}, m \geq n, \text{rank } A = n$: $A^\dagger = (A^\top A)^{-1} A^\top$

Fat – $A \in \mathbb{R}^{m \times n}, m \leq n, \text{rank } A = m$: $A^\dagger = A^\top (AA^\top)^{-1}$

Generalized inverse

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times n}$ is called a **generalized inverse** of A if

$$AA^\dagger A = A$$

and there exist matrices $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ such that

$$A^\dagger = UA^\top \quad \text{and} \quad A^\dagger = A^\top V.$$

Tall – $A \in \mathbb{R}^{m \times n}, m \geq n, \text{rank } A = n$: $A^\dagger = (A^\top A)^{-1} A^\top$

Fat – $A \in \mathbb{R}^{m \times n}, m \leq n, \text{rank } A = m$: $A^\dagger = A^\top (AA^\top)^{-1}$

What are U and V here?

Generalized inverse

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times n}$ is called a **generalized inverse** of A if

$$AA^\dagger A = A$$

and there exist matrices $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ such that

$$A^\dagger = UA^\top \quad \text{and} \quad A^\dagger = A^\top V.$$

Tall – $A \in \mathbb{R}^{m \times n}, m \geq n, \text{rank } A = n$: $A^\dagger = (A^\top A)^{-1} A^\top$

Fat – $A \in \mathbb{R}^{m \times n}, m \leq n, \text{rank } A = m$: $A^\dagger = A^\top (AA^\top)^{-1}$

What are U and V here?

If A^\dagger exists, it is *unique*.

Proof?

Generalized inverse

General form

Theorem – Let $A \in \mathbb{R}^{m \times n}$, $\text{rank } A = r \leq \min\{m, n\}$ have a full rank factorization

$$A = BC$$

with $B \in \mathbb{R}^{m \times r}$, $C \in \mathbb{R}^{r \times n}$, and

$$\text{rank } A = \text{rank } B = \text{rank } C = r.$$

Then,

$$A^\dagger = C^\dagger B^\dagger.$$

Generalized inverse

General form

Theorem – Let $A \in \mathbb{R}^{m \times n}$, $\text{rank } A = r \leq \min\{m, n\}$ have a full rank factorization

$$A = BC$$

with $B \in \mathbb{R}^{m \times r}$, $C \in \mathbb{R}^{r \times n}$, and

$$\text{rank } A = \text{rank } B = \text{rank } C = r.$$

Then,

$$A^\dagger = C^\dagger B^\dagger.$$

Proof –

1. Verify that $AC^\dagger B^\dagger A = A$

Generalized inverse

General form

Theorem – Let $A \in \mathbb{R}^{m \times n}$, $\text{rank } A = r \leq \min\{m, n\}$ have a full rank factorization

$$A = BC$$

with $B \in \mathbb{R}^{m \times r}$, $C \in \mathbb{R}^{r \times n}$, and

$$\text{rank } A = \text{rank } B = \text{rank } C = r.$$

Then,

$$A^\dagger = C^\dagger B^\dagger.$$

Proof –

1. Verify that $AC^\dagger B^\dagger A = A$
2. Let

$$U = C^\top (CC^\top)^{-1} (B^\top B)^{-1} (CC^\top)^{-1} C$$

$$V = B (B^\top B)^{-1} (CC^\top)^{-1} (B^\top B)^{-1} B^\top$$

Generalized inverse

General form

Theorem – Let $A \in \mathbb{R}^{m \times n}$, $\text{rank } A = r \leq \min\{m, n\}$ have a full rank factorization

$$A = BC$$

with $B \in \mathbb{R}^{m \times r}$, $C \in \mathbb{R}^{r \times n}$, and

$$\text{rank } A = \text{rank } B = \text{rank } C = r.$$

Then,

$$A^\dagger = C^\dagger B^\dagger.$$

Proof –

1. Verify that $AC^\dagger B^\dagger A = A$
2. Let

$$U = C^\top (CC^\top)^{-1} (B^\top B)^{-1} (CC^\top)^{-1} C$$

$$V = B (B^\top B)^{-1} (CC^\top)^{-1} (B^\top B)^{-1} B^\top$$

3. Verify that $C^\dagger B^\dagger = UA^\top$ and $A^\dagger = A^\top V$

Generalized inverse

$$A^\dagger = C^\dagger B^\dagger \quad ; \quad B^\dagger = (B^\top B)^{-1} B^\top \quad ; \quad C^\dagger = C^\top (C C^\top)^{-1}$$

$$A^\dagger = C^\top (C C^\top)^{-1} (B^\top B)^{-1} B^\top$$

Generalized inverse

$$A^\dagger = C^\dagger B^\dagger \quad ; \quad B^\dagger = (B^\top B)^{-1} B^\top \quad ; \quad C^\dagger = C^\top (CC^\top)^{-1}$$

$$\begin{aligned} A^\dagger &= C^\top (CC^\top)^{-1} (B^\top B)^{-1} B^\top \\ &= C^\top (B^\top A C^\top)^{-1} B^\top \end{aligned}$$

Generalized inverse

$$A^\dagger = C^\dagger B^\dagger \quad ; \quad B^\dagger = (B^\top B)^{-1} B^\top \quad ; \quad C^\dagger = C^\top (CC^\top)^{-1}$$

$$\begin{aligned} A^\dagger &= C^\top (CC^\top)^{-1} (B^\top B)^{-1} B^\top \\ &= C^\top (B^\top A C^\top)^{-1} B^\top \end{aligned}$$

Verify using $A = BC$

Generalized inverse

$$A^\dagger = C^\dagger B^\dagger \quad ; \quad B^\dagger = (B^\top B)^{-1} B^\top \quad ; \quad C^\dagger = C^\top (CC^\top)^{-1}$$

$$\begin{aligned} A^\dagger &= C^\top (CC^\top)^{-1} (B^\top B)^{-1} B^\top \\ &= C^\top (B^\top A C^\top)^{-1} B^\top \end{aligned}$$

Verify using $A = BC$

Caution – $A^\dagger = C^\dagger B^\dagger$ *not valid* if $A = BC$ is *not* a full rank factorization.

See Example 12.11 in textbook.

Equivalent definition

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times m}$ is called a **generalized inverse** of A if

$$AA^\dagger A = A$$

and there exist matrices $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ such that

$$A^\dagger = UA^\top \quad \text{and} \quad A^\dagger = A^\top V.$$

Equivalent definition

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times m}$ is called a **generalized inverse** of A if

$$AA^\dagger A = A$$

and there exist matrices $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ such that

$$A^\dagger = UA^\top \quad \text{and} \quad A^\dagger = A^\top V.$$

equivalently, the **Penrose definition**:

Equivalent definition

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times m}$ is called a **generalized inverse** of A if

$$AA^\dagger A = A$$

and there exist matrices $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ such that

$$A^\dagger = UA^\top \quad \text{and} \quad A^\dagger = A^\top V.$$

equivalently, the **Penrose definition**:

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times m}$ is called a **generalized inverse** of A if

1. $AA^\dagger A = A$
2. $A^\dagger AA^\dagger = A^\dagger$
3. $(AA^\dagger)^\top = AA^\dagger$
4. $(A^\dagger A)^\top = A^\dagger A$.

Equivalent definition

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times m}$ is called a **generalized inverse** of A if

$$AA^\dagger A = A$$

and there exist matrices $U \in \mathbb{R}^{n \times n}$ and $V \in \mathbb{R}^{m \times m}$ such that

$$A^\dagger = UA^\top \quad \text{and} \quad A^\dagger = A^\top V.$$

equivalently, the **Penrose definition**:

Definition – Given a matrix $A \in \mathbb{R}^{m \times n}$, a matrix $A^\dagger \in \mathbb{R}^{n \times m}$ is called a **generalized inverse** of A if

1. $AA^\dagger A = A$
2. $A^\dagger AA^\dagger = A^\dagger$
3. $(AA^\dagger)^\top = AA^\dagger$
4. $(A^\dagger A)^\top = A^\dagger A$.

(Verify the equivalence)

Properties of A^\dagger

Verify:

$$(A^\top)^\dagger = (A^\dagger)^\top$$

$$(A^\dagger)^\dagger = A$$

Properties of A^\dagger

Verify:

$$(A^\top)^\dagger = (A^\dagger)^\top$$

$$(A^\dagger)^\dagger = A$$

However,

$$(BC)^\dagger \neq C^\dagger B^\dagger \quad \text{in general}^*$$

(construct an example)

*Note that if it were true that

$$(BC)^\dagger = C^\dagger B^\dagger \quad \text{for all } B, C,$$

then we **need not have** proved that $C^\dagger B^\dagger$ is the generalized inverse of A since it would have followed from

$$A = BC.$$

Least squares, minimum norm

Theorem – Consider a system of linear equations

$$Ax = b, \quad A \in \mathbb{R}^{m \times n}, \quad \text{rank } A = r.$$

The vector

$$x^* = A^\dagger b$$

minimizes $\|Ax - b\|^2$ on \mathbb{R}^n .

Furthermore, among all vectors in \mathbb{R}^n that minimize $\|Ax - b\|^2$, the vector

$$x^* = A^\dagger b$$

is the unique vector with minimal norm.

(proof on board)