

```
In [1]: import numpy as np
import matplotlib.pyplot as plt
from matplotlib.ticker import AutoMinorLocator
import sklearn as sk
from scipy import stats as st
from scipy.optimize import minimize
import pandas as pd
import os
```

```
In [2]: np.set_printoptions(formatter={'float': lambda x: "{0:0.2e}".format(x)}) #to
```

Functions

```
In [3]: #for plotting distributions
def plot_distrib(data, bins = 100, title = ""):
    plt.hist(data, bins=bins, density=True)
    plt.title(title)
    plt.ylabel("Probability density")
```

```
In [4]: #to calculate 1σ uncertainty bands and medians
def median_and_error(data):
    med, low, high = np.percentile(data, [50, 15.87, 84.13])
    median_and_error = np.array([med, low - med, high - med])

    return median_and_error
```

```
In [5]: #Fits a skew-normal distribution to a given set of percentiles.
def fit_skewnorm(percentiles, values): #the percentiles and corresponding va

    #set initial guesses as if it was a gaussian
    median = np.interp(0.5, percentiles, values)
    p10 = np.interp(0.1, percentiles, values)
    p90 = np.interp(0.9, percentiles, values)
    init_scale = (p90 - p10) / 2.56 if (p90 - p10) > 0 else 1 #for a gaussi

    initial_guesses = [0., median, init_scale]

    bounds = [(-20, 20), (None, None), (1e-6, None)]  #(a, loc, scale) -> kee

    #Error function to minimize
    def error_function(params):
        a, loc, scale = params
        theoretical_values = st.skewnorm.ppf(percentiles, a=a, loc=loc, scal
        return np.sum((theoretical_values - values) ** 2) #sum of squared er

    result = minimize(
        error_function,
        initial_guesses,
        method='SLSQP',
        bounds=bounds,
        options={'ftol': 1e-9, 'disp': False}
```

```
)

return result.x #(a, loc, scale)
```

Isolation Time Figure

```
In [6]: k = 2.3 #best
```

```
In [7]: #production ratio
prIaNb = 1.58*10**-3
prIaSm = 0.347

EssNb = (3.2 * 10**-5, 0.3 * 10**-5)
EssSm = (8.28 * 10**-3, 0.44 * 10**-3)

#mean-life (half-life values from NUBASE)
tauNb = np.array([34.7, 2.4]) / np.log(2.)
tauSm = np.array([92, 2.6]) / np.log(2.)
tau = np.array([tauNb, tauSm])

#age of the galaxy
TGal = 8400
```

```
In [8]: #τ for table
print(tau)
```

```
[5.01e+01 3.46e+00]
[1.33e+02 3.75e+00]
```

Fit skewnormal distribution to percentiles

Assuming assuming that a skew-normal distribution is a sufficient approximation for the assymetric distributions of the ISS ratios.

Percentiles from [Côté et al. \(2019\)](#), γ is taken to be 10 Myr.

```
In [9]: percentiles = [0.16, 0.50, 0.84]

statNb = [3.13-0.73, 3.13, 3.13+0.80]
statSm = [9.99-0.84, 9.99, 9.99+0.87]

paramNb = fit_skewnorm(percentiles, statNb)
paramSm = fit_skewnorm(percentiles, statSm)


distNb = st.skewnorm(paramNb[0], paramNb[1], paramNb[2])
distSm = st.skewnorm(paramSm[0], paramSm[1], paramSm[2])

#normalise
paramNb[1:] = paramNb[1:] / distNb.mean()
paramSm[1:] = paramSm[1:] / distSm.mean()
```

```
#create normalised distribution
distNb = st.skewnorm(paramNb[0], paramNb[1], paramNb[2])
distSm = st.skewnorm(paramSm[0], paramSm[1], paramSm[2])
```

```
In [10]: #visualising the fitted distributions
x = np.linspace(0,5,1000)
plt.plot(x,distNb.pdf(x=x),marker='none',label='Nb',color=plt.get_cmap('Blue'))
plt.plot(x,distSm.pdf(x=x),marker='none',label='Sm',color='r')
plt.legend()
```

Out[10]: <matplotlib.legend.Legend at 0x7a8730e9fe90>



Calculate and plot isolation times for Typela production

Then I draw Monte Carlo distributions of the ESS and ISM ratios, and mean-lives. τ and ESS is assumed to be Gaussian, ISM comes from the fitted skew-normal distributions.

```
In [11]: mcNum = 10**7 #number of samples for MC
```

```
In [12]: #Sampling distributions for the  $\tau$  values
ratTauNb = np.clip(np.random.normal(loc = tauNb[0], scale = tauNb[1], size=mcNum), 0, 1)
ratTauSm = np.clip(np.random.normal(loc = tauSm[0], scale = tauSm[1], size=mcNum), 0, 1)
```

```
In [13]: #The ISM ratios
IsmNb = k * prIaNb * ratTauNb / TGal
```

```
IsmSm = k * prIaSm * ratTauSm / TGal
```

```
In [14]: #Sampling distributions for the ISM values
ratIsmNb = np.clip(IsmNb * distNb.rvs(mNum), 1e-30, np.inf)
ratIsmSm = np.clip(IsmSm * distSm.rvs(mNum), 1e-30, np.inf)
```

```
In [15]: #Sampling normal distribution for the ESS values
ratEssNb = np.clip(np.random.normal(loc = EssNb[0], scale = EssNb[1], size=n
ratEssSm = np.clip(np.random.normal(loc = EssSm[0], scale = EssSm[1], size=n
```

```
In [16]: #Calculating isolation time
isoTNb = -np.log(ratEssNb / ratIsmNb) * ratTauNb
isoTSm = -np.log(ratEssSm / ratIsmSm) * ratTauSm
```

```
In [17]: #downsample for visualisation (this is for optimisation)
isoTSampleNb = sk.utils.resample(isoTNb, n_samples=1000000)
isoTSampleSm = sk.utils.resample(isoTSm, n_samples=1000000)
```

```
In [18]: #create the approximate isolation time distributions
isoTSampleNbKern = sk.neighbors.KernelDensity(kernel='gaussian', bandwidth=1
isoTSampleNbKern.fit(isoTSampleNb.reshape(-1, 1))
isoTSampleSmKern = sk.neighbors.KernelDensity(kernel='gaussian', bandwidth=1
isoTSampleSmKern.fit(isoTSampleSm.reshape(-1, 1))
```

```
Out[18]: KernelDensity
```

Parameters

```
In [19]: x = np.linspace(-120, 120, 100).reshape(-1, 1)
```

```
In [20]: #evaluate the distribution at 100 points
isoTSampleNbDensity = np.exp(isoTSampleNbKern.score_samples(x))
isoTSampleSmDensity = np.exp(isoTSampleSmKern.score_samples(x))
```

```
In [21]: #Creating figure
blue = plt.get_cmap('Blues')(2/4)

plt.plot(x, isoTSampleNbDensity, color=blue, label='${92}$Nb')
plt.fill_between(x.flatten(), isoTSampleNbDensity, 0, color=blue, alpha=0.3)

plt.plot(x, isoTSampleSmDensity, color='red', label='${146}$Sm')
plt.fill_between(x.flatten(), isoTSampleSmDensity, 0, color='red', alpha=0.3)

plt.xlabel('Isolation time (Myr)')
plt.ylabel('Probability density')
plt.legend(loc='upper right')


#for tickmarks
plt.gca().xaxis.set_minor_locator(AutoMinorLocator())
plt.gca().yaxis.set_minor_locator(AutoMinorLocator())
plt.gca().tick_params(axis='x', which='both', top=True)
plt.gca().tick_params(axis='y', which='both', right=True)
```

```
plt.legend(loc='upper right')

#for the gray box
plt.axvspan(xmin=x.flatten()[0], xmax=0, color='gray', alpha=0.3, zorder=0)
plt.text(0.05, 0.95, "Isolation times < 0 \n(grey) are unphysical", transform=plt.gca().transAxes)

plt.xlim(x[0], x[-1])

plt.savefig("../figures/"+ "isoT_Ia.png", bbox_inches='tight')
plt.show()
```



Calculate and plot isolation times for CCSN production

```
In [22]: # Import production ratios of different CCSN models
data = pd.read_csv("../data/production_ratios.txt", sep='\\s+')
data = data.set_index("model")
```

```
In [23]: shock = np.array(data['initial_shock'])
mtot = np.array(data['MTOT'])
mrem = np.array(data['MREM'])
msurf = np.array(data['MSURF'])
modelSet = np.array(data['Set'])

prCCSnNb = np.array(data['Nb-92/Mo-92'])
prCCSnSm = np.array(data['Sm-146/Sm-144'])
```

```

In [24]: mcNum = 10**7 #number of samples for MC

In [25]: #Sampling distributions for the  $\tau$  values
ratTauNb = np.clip(np.random.normal(loc=tauNb[0], scale=tauNb[1], size=mcNum), 1e-30, np.inf)
ratTauSm = np.clip(np.random.normal(loc=tauSm[0], scale=tauSm[1], size=mcNum), 1e-30, np.inf)

In [26]: #The ISM ratios
IsmNbCCSn = k * prCCSnNb[:, np.newaxis] * ratTauNb / TGal
IsmSmCCSn = k * prCCSnSm[:, np.newaxis] * ratTauSm / TGal

In [27]: ratIsmbNbCCSn = np.clip(IsmNbCCSn * distNb.rvs(mcNum), 1e-30, np.inf)
ratIsmbSmCCSn = np.clip(IsmSmCCSn * distSm.rvs(mcNum), 1e-30, np.inf)

isoTNbCCSn = -np.log(ratEssNb / ratIsmbNbCCSn) * ratTauNb
isoTSmCCSn = -np.log(ratEssSm / ratIsmbSmCCSn) * ratTauSm

In [28]: isoTSampleNbCCSn = []
isoTSampleSmCCSn = []

for i in range(np.size(prCCSnNb)):
    isoTSampleNbCCSn.append(sk.utils.resample(isoTNbCCSn[i], n_samples=10000))
    isoTSampleSmCCSn.append(sk.utils.resample(isoTSmCCSn[i], n_samples=10000))

In [29]: isoTNbKernCCSn = []
isoTSmKernCCSn = []

for i in range(np.size(prCCSnNb)):
    isoTNbKernCCSn.append(sk.neighbors.KernelDensity(kernel='gaussian', bandwidth=1).fit(isoTSampleNbCCSn[i].reshape(-1, 1)))
    isoTSmKernCCSn.append(sk.neighbors.KernelDensity(kernel='gaussian', bandwidth=1).fit(isoTSampleSmCCSn[i].reshape(-1, 1)))

In [30]: #create the ranges for plotting the data (limits selected based on previous
limits = np.array([[-150,50],[-150,50],[-150,110],[-150,110],[-380,110]])
x = np.linspace(limits[:,0], limits[:,1], 100).T

In [31]: #evaluate the distributions at 100 points
isoTNbDensityBest_CCSn = []
isoTSmDensityBest_CCSn = []

for i in range(np.size(prCCSnNb)):
    isoTNbDensityBest_CCSn.append(np.exp(isoTNbKernCCSn[i].score_samples(x[i])))
    isoTSmDensityBest_CCSn.append(np.exp(isoTSmKernCCSn[i].score_samples(x[i])))

In [32]: blue = plt.get_cmap('Blues')(2/4)

for i in range(np.size(prCCSnNb)):

    plt.plot(x[i], isoTNbDensityBest_CCSn[i], color=blue, label='${92}$Nb')
    plt.fill_between(x[i], isoTNbDensityBest_CCSn[i], 0, color=blue, alpha=0.5)

    #for tickmarks
    plt.gca().xaxis.set_minor_locator(AutoMinorLocator())

```

```

plt.gca().yaxis.set_minor_locator(AutoMinorLocator())
plt.gca().tick_params(axis='x', which='both', top=True)
plt.gca().tick_params(axis='y', which='both', right=True)

plt.plot(x[i], isoTSmDensityBest_CCSn[i], color="red", label='${146}$Sm')
plt.fill_between(x[i].flatten(), isoTSmDensityBest_CCSn[i], 0, color="red")

plt.xlabel('Isolation time (Myr)')
plt.ylabel('Probability density')
plt.legend(loc='upper right')


#for the gray box
plt.axvspan(xmin=x[i][0], xmax=0, color='gray', alpha=0.3, zorder=0)
#plt.text(0.05, 0.95, "Isolation times < 0 \n(grey) are unphysical", tra



plt.xlim(x[i][0], x[i][-1])



plt.title(modelSet[i]+", "+str(mtot[i])+"M$_{\odot}$", "+str(shock[i])+"$\\cdot$")

plt.savefig("../figures/"+modelSet[i]+"_M"+str(int(mtot[i]))+"_V"+str(sh
plt.show()

```







1 σ uncertainty bands for ISS ratios

```
In [33]: #median, -, +
for i in range(np.size(prCCSnNb)):
    print(modelSet[i]+", "+str(mtot[i])+"M $\odot$ , "+str(shock[i])+" · 109 cm/s")
    print("Nb: ",median_and_error(ratIsmNbCCSn[i]))
    print("Sm: ",median_and_error(ratIsmSmCCSn[i]))
    print()
```

Roberti, 20.0M $\odot$, 5.0 · 10⁹ cm/s
 Nb: [7.75e-06 -1.94e-06 2.01e-06]
 Sm: [4.36e-03 -3.92e-04 3.98e-04]

Roberti, 20.0M $\odot$, 4.0 · 10⁹ cm/s
 Nb: [7.66e-06 -1.91e-06 1.98e-06]
 Sm: [5.07e-03 -4.55e-04 4.62e-04]

Roberti, 20.0M $\odot$, 3.0 · 10⁹ cm/s
 Nb: [5.87e-05 -1.47e-05 1.52e-05]
 Sm: [4.39e-03 -3.94e-04 4.00e-04]

Roberti, 20.0M $\odot$, 2.0 · 10⁹ cm/s
 Nb: [4.95e-05 -1.24e-05 1.28e-05]
 Sm: [4.55e-03 -4.09e-04 4.15e-04]

Ritter, 20.0M $\odot$, 2.0 · 10⁹ cm/s
 Nb: [1.58e-05 -3.94e-06 4.08e-06]
 Sm: [8.11e-04 -7.28e-05 7.40e-05]

1 σ uncertainty bands for the Isolation Times

```
In [34]: #median, -, +
for i in range(np.size(prCCSnNb)):
    print(modelSet[i]+", "+str(mtot[i])+"M $\odot$ , "+str(shock[i])+" · 109 cm/s")
    print("Nb: ",median_and_error(isoTNbCCSn[i]))
    print("Sm: ",median_and_error(isoTSmCCSn[i]))
    print()
```

Roberti, $20.0M_{\odot}$, $5.0 \cdot 10^9$ cm/s
 Nb: [-7.05e+01 -1.48e+01 1.20e+01]
 Sm: [-8.49e+01 -1.38e+01 1.33e+01]

Roberti, $20.0M_{\odot}$, $4.0 \cdot 10^9$ cm/s
 Nb: [-7.12e+01 -1.48e+01 1.20e+01]
 Sm: [-6.51e+01 -1.38e+01 1.33e+01]

Roberti, $20.0M_{\odot}$, $3.0 \cdot 10^9$ cm/s
 Nb: [3.02e+01 -1.52e+01 1.37e+01]
 Sm: [-8.42e+01 -1.38e+01 1.33e+01]

Roberti, $20.0M_{\odot}$, $2.0 \cdot 10^9$ cm/s
 Nb: [2.17e+01 -1.51e+01 1.34e+01]
 Sm: [-7.94e+01 -1.38e+01 1.33e+01]

Ritter, $20.0M_{\odot}$, $2.0 \cdot 10^9$ cm/s
 Nb: [-3.52e+01 -1.46e+01 1.21e+01]
 Sm: [-3.08e+02 -1.47e+01 1.40e+01]

References:

- Leckenby, G., Sidhu, R. S., Chen, R. J., Szányi, B., Dillmann, I., Gernhäuser, R., Glorius, J., Griffin, C., Litvinov, Y. A., Lugaro, M., Sanjari, S., & Yagüe López, A. (2024). Measurement of the bound-state beta decay of $^{205}\text{Tl}(81+)$: analysis scripts and figures (1.0). Zenodo. <https://doi.org/10.5281/zenodo.11560338>
- Côté, Benoit et al. (Dec. 2019). "Stochastic Chemical Evolution of Radioactive Isotopes with a Monte Carlo Approach". In: ApJ 887.2, 213, p. 213. doi: [10.3847/1538-4357/ab5a88](https://doi.org/10.3847/1538-4357/ab5a88). arXiv: [1911.01457](https://arxiv.org/abs/1911.01457) [astro-ph.GA].