

Studentpad

MHT-CET-XI CHEMISTRY 2022-23

Time : 150 Min

Chem : Full Portion Paper

Marks : 50

Hints and Solutions

01) Ans: **A)** adsorbent

02) Ans: **C)** $\text{Li}^+ > \text{Na}^+ > \text{K}^+ > \text{Rb}^+ > \text{Cs}^+$

Sol: Smaller the cation, greater is the degree of its hydration.

As the degree of hydration decreases from

Li^+ to Cs^+ , the hydration energy of alkali metals

also decreases from Li to Cs^+ .

Therefore the sequence in

$\text{Li}^+ > \text{Na}^+ > \text{K}^+ > \text{Rb}^+ > \text{Cs}^+$

03) Ans: **C)** Stability of hydrides of alkali metals decreases from CsH to LiH

Sol: Stability of hydrides decrease from LiH to CsH .

04) Ans: **A)** 3

Sol: $[\text{H}^+] \sqrt{\text{Ka.C}} = \sqrt{1 \times 10^{-5} \times 0.1} = \sqrt{10^{-6}}$

$[\text{H}^+] = 10^{-3} \text{ mol / dm}^3 \therefore \text{pH} = 3$

05) Ans: **D)** TI

06) Ans: **B)** biological fluids

Sol: Monovalent sodium and potassium ions and divalent magnesium and calcium ions are found in large proportions in biological fluids.

These ions perform important biological functions like maintenance of ion balance and nerve impulse conduction.

07) Ans: **C)** Salting

Sol: Salting is the most common chemical method to preserve fishes.

08) Ans: **B)** 32 hours

Sol: $1 - \frac{1}{2} - \frac{1}{4} (0.25 \text{ gm})$ i.e. 2 half lives = 16 hr

$\therefore \text{H.L.P.} = 8 \text{ hrs}$

$48 \text{ gm} - 24 - 12 - 6 - 3 \text{ gm}$

4 half lives i.e. $8 \times 4 = 32 \text{ hrs}$ needed.

09) Ans: **A)** H_2^{2+} , He_2

Sol: H_2^+ : Bond order = 0

He_2 : Bond order = $\frac{2-2}{2} = 0$

So, both H_2^{2+} and He_2 do not exist.

10) Ans: **D)** (Absolute error/measurement being taken)

Sol: Relative error (RE) - when used as a measure of precision - is the ratio of the absolute error of a

measurement to the measurement being taken.

In other words, this type of error is relative to the size of the item being measured. RE is expressed as a percentage and has no units.

11) Ans: **C)** -9E

Sol: $E_3 = -E = -13.6 \times \frac{Z^2}{3^2}$

$\Rightarrow -E = E_0 \times \frac{Z^2}{9} [E_0 = -13.6]$

$\Rightarrow E_0 = \frac{-9E}{Z^2} \dots (i)$

According to question

$E_1 = -13.6 \times \frac{Z^2}{1^2} = E_0 \times Z^2 = \frac{-9E}{Z^2} \times Z^2 \Rightarrow E_1 = -9E$

12) Ans: **B)** neither has electron pair available for donation nor can accommodate electron since all shells of nitrogen are fully occupied

Sol: $(\text{CH}_3)_4\text{N}^+$ has no electron pair available for donation to act as nucleophile.

Valence shell of nitrogen to act as nucleophile.

Valence shell of nitrogen is completely filled and has no scope for acceptance of extra electrons to act as electrophile.

13) Ans: **B)** β rays

Sol: β rays.

14) Ans: **B)** n-pentane > 2-methylbutane > 2,2-dimethylpropane

Sol: Branching of the chain makes the molecule more compact and thereby decreases the surface area. and the intermolecular forces become small in magnitude on account of branching.

Consequently, the boiling points of branched chain alkanes are less than the straight chain isomers.

15) Ans: **C)** $\text{C} < \text{N} < \text{F} < \text{O}$

Sol: The second I.P. means removal of electron from a cation

$\text{C}^{-1} (5e) = 1s^2 2s^2 2p^1$

$\text{N}^{-1} (6e) = 1s^2 2s^2 2p^2$

$\text{O}^{-1} (7e) = 1s^2 2s^2 2p^3$

$\text{F}^{-1} (8e) = 1s^2 2s^2 2p^4$

$\therefore \text{C} < \text{N} < \text{F} < \text{O}$.

16) Ans: **C)** $P_1 T_2 = P_2 T_1$

Sol: Gay-lussac's law state that,

$P \propto T$ (At constant volume)


$$\therefore \frac{P_1}{T_1} = \frac{P_2}{T_2}, \therefore P_1 T_2 = P_2 T_1$$

17) Ans: C) two molecular orbital.


Sol: They will form one bonding M.O. and one anti-bonding M.O.

18) Ans: A) Dehydrohalogenation of an alkyl halide

Sol: Dehydrohalogenation of an alkyl halide gives alkene not alkane.



Dehydrohalogenation is a β -elimination and it is governed by Saytzeff rule according to which a more substituted alkene is obtained as major product.



19) Ans: D) 0.0693 year^{-1}

Sol: 0.0693 year^{-1}

20) Ans: D) +6 and +6


Sol: Peroxomonosulphuric acid $\text{H}-\text{O}-\text{O}-\overset{\text{O}}{\underset{\text{O}}{\parallel}}\text{S}-\text{OH}$

$$1+x+2(-1)+2(-2)+(-2)+1=0; x=+6$$

Peroxodisulphuric acid $\text{OH}-\overset{\text{O}}{\underset{\text{O}}{\parallel}}\text{S}-\text{O}-\text{O}-\overset{\text{O}}{\underset{\text{O}}{\parallel}}\text{S}-\text{OH}$

$$1+(-2)+x+2(-2)+2(-1)+x+2(-2)+(-2)+1=0; x=+6$$

21) Ans: B) 35 dm^3



1 volume 3.5 volume

2 dm^3 7 dm^3

For complete combustion of 2 dm^3 of ethane 7 dm^3 of oxygen is required.

\therefore air contains 20% of oxygen

$\therefore 20 \text{ dm}^3$ of oxygen $\equiv 100 \text{ dm}^3$ of air

$$7 \text{ dm}^3 \text{ of oxygen} = \frac{7 \times 100}{20} = 35 \text{ dm}^3 \text{ of air}$$

Volume of air required is 35 dm^3

22) Ans: C) electronic configuration

23) Ans: C) 16

Sol: The expression is given as,

$$w=4 \text{ gm}, R=8.314 \text{ JK}^{-1} \text{ mol}^{-1}$$

$$V=5.60 \times 10^{-3} \text{ m}^3 \quad T=546 \text{ K}$$

$$P=2.026 \times 10^{-5} \text{ N/m}^2$$

We know that, $Pv=nRT$

$$\frac{wRT}{\text{Mol.wt}} \left(\because n = \frac{wt}{\text{mol.wt}} \right)$$

$$\text{mol.wt.} = \frac{wRT}{PV} = \frac{4 \times 8.314 \times 546}{2.026 \times 10^{-5} \times 5.603 \times 10^{-3}}$$


$$= \frac{18158}{1135} = 16$$

24) Ans: A) Chloramphenicol

Sol: Chloramphenicol is broad spectrum antibiotic, which is used in the treatment of typhoid, dysentery, acute fever.

25) Ans: D) distillation

Sol: The difference between the boiling point of oxygen and Nitrogen is frequently large to be separated by distillation.



Sol:

1° carbon - A carbon atom attached to one carbon atom only is termed as 1° carbon or primary carbon.

2° carbon - A carbon atom attached to two carbon atoms is called 2° carbon or secondary carbon.

3° carbon - A carbon atom attached to three carbon atoms is called 3° carbon or tertiary carbon.

4° carbon - A carbon atom attached to four carbon atoms is called 4° carbon or quaternary carbon.

27) Ans: D) poor shielding effect of d-electrons of Ga atoms

Sol: Shielding of d electrons is poor than s as well as p electrons.

28) Ans: A) CH_3CN


Sol: In CH_3CN , the bond order between C and N is 3, thus its bond length is minimum.

29) Ans: C) 1


30) Ans: A) 3-Methylhexane







Sol: As per the IUPAC rule.

31) Ans: A) 5,6,3,3



Step 1 : $\text{I}^- \rightarrow \text{I}_2$ (oxidation)





By comparing, a=5, b=6, c=3, d=3.

32) Ans: B) critical temperature

Sol: The highest temperature at which liquefaction of gas first occurs is termed as critical temperature.

33) Ans: D) hydrogen bonding

Sol: It (boric acid) has BO_3 units which are bonded by hydrogen bonding.

34) Ans: A) 10 ml

Sol: $N_1V_1 = N_2V_2$

$$0.001 \times V_1 = 0.0001 \times 900$$


$$V_1 = \frac{9 \times 10^{-2}}{10^{-3}} = 90 \text{ ml}$$

The original volume=90=10 ml

35) Ans: A) 3



Writing oxidation states, we have



To balance oxidation state of Cr on both sides, add 3e^- on R.H.S.

36) Ans: B) 200-500

37) Ans: A) 2.4 N_A


38) Ans: C) II > IV > I > III

Sol: Stability of carbocations is enhanced by

(i) presence of conjugation.

(ii) presence of group having +I effect.

Therefore, the correct order is





39) Ans: B) Ionic, polar covalent



Sol: In compound AC electronegativity difference is 2 (i.e. 3-1) hence bond is ionic.

In compound BC electronegativity difference is 0.5 (=3-2.5) evidently the bond is polar covalent.

40) Ans: B) Be



All alkali metals decompose water with the evolution of hydrogen gas.



Ca, Sr, Ba and Ra decompose cold water readily with evolution of hydrogen.

Mg decomposes boiling water but Be is not attacked by water even at high temperatures as its oxidation potential is lower than the other members.

41) Ans: A) tertiary > secondary > primary

42) Ans: D) hydrogen bonding interactions.

Sol: Water is denser than ice due to hydrogen bonding interaction and structure of ice.


43) Ans: C) sp^2 hybridized

44) Ans: C) Sorption

45) Ans: C) 25-100

Sol: Crystalline phases are prepared at low temperature range such as 25-100 degree Celsius whereas much high temperatures would be required for a normal solid state synthesis. Crystals usually form at low temperatures but because of the release of heat of fusion during crystallization, the entropy of the universe increases so the second law of thermodynamics is not altered.

46) Ans: B) nuclear fission



Neutrons thus formed bring about further nuclear fission. Hence it results into chain reaction producing large amount of energy is released.

47) Ans: B) conformational isomerism


Sol: The different arrangements of atoms in space that result from the free rotation of groups about C-C bond axis are called conformational isomers and the phenomenon is known as conformational isomerism.

48) Ans: B) There is loss of electrons

49) Ans: B) $3.0 \times 10^{-19} \text{ J}$

$$\text{Sol: } E = \frac{hc}{\lambda} = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{660 \times 10^{-9}} = 3 \times 10^{-19} \text{ J}$$

50) Ans: A) 53.78



$$2(36 + 1 + 31 + 64) = 62 + 80 = 264 = 142$$

$$\% \text{P}_2\text{O}_5 = \frac{\text{Wt. of P}_2\text{O}_5}{\text{Wt. of s}_9\text{H}} \times 100 = \frac{142}{264} \times 100 = 53.78\%$$