

The genome of the Pacific acorn barnacle provides insights into the evolution of extremely large populations

Angel G. Rivera-Colón¹, Scott T. Small¹, and Andrew D. Kern¹

¹University of Oregon, Institute of Ecology and Evolution

September 24, 2025

Abstract

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

Introduction

Nulla ac nisl. Nullam urna nulla, ullamcorper in, interdum sit amet, gravida ut, risus. Aenean ac enim. In luctus. Phasellus eu quam vitae turpis viverra pellentesque. Duis feugiat felis ut enim. Phasellus pharetra, sem id porttitor sodales, magna nunc aliquet nibh, nec blandit nisl mauris at pede. Suspendisse risus risus, lobortis eget, semper at, imperdiet sit amet, quam. Quisque scelerisque dapibus nibh. Nam enim. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Nunc ut metus. Ut metus justo, auctor at, ultrices eu, sagittis ut, purus. Aliquam aliquam.

Results

Genome assembly

We assembled a chromosome-level reference genome for the Pacific acorn barnacle using PacBio Hi-Fi long reads and Hi-C chromosome conformation capture. Our initial contig-level assembly, generated using `hifiasm` [Cheng2021; Cheng2022] with default parameters, generated sequences with a total length of 1.64 Gbp, composed of 4,212 contigs, and a contig N50 of 734 kbp. We also assessed gene-completeness using BUSCOs [Manni2021; Tegenfeldt2024], as implemented in `compleseam` [Huang&Li2023], resulting in a completeness of 95.66% ([S:45.31%; D:50.35%; F:0.79%; M:3.55%; N:1,013]) for the `arthropoda_odb10` single-copy ortholog set. The statistics for this initial assembly, with twice the expected genome size and over 50% duplicated BUSCOs, suggested the presence of a large fraction of haplotig sequences, reflecting the high heterozygosity expected for the species.

Following optimization, both in the integration of additional data types and in the increased stringency of the purging of haplotig and contaminant sequences (see Methods), we generated a contig-level assembly composed of 1,342 sequences, with a total length of 1.05 Gbp, a contig N50 of 1.35 Mbp, and a BUSCO completeness of 93.88% ([S:89.04%; D:4.84%; F:1.18%; M:4.94%; N:1,013], `arthropoda_odb10`). The contig-level assembly was then scaffolded with Hi-C data using the `YAHS` software [Zhou2022]. Our scaffolding process generated an assembly with a total length of 1.05 Gbp, composed of 650 scaffolds, a scaffold N50 of 50.96 Mbp, and a BUSCO completeness of 93.98% ([S:89.24%; D:4.74%; F:0.89%; M:5.13%; N:1,013], `arthropoda_odb10`).

This scaffold-level assembly was further processed by performing manual curation of the Hi-C contact map and self-correction using the software **Inspector** [Chen2021]. This curated assembly was composed of 592 fragments, a total sequence length of 1.04 Gbp, a scaffold N50 of 51.41 Mbp, and a per-base quality value (QV) of 30.49. When assessing completeness, the curated assembly was 94.08% ([S:89.44%; D:4.64%; F:0.89%; M:5.03%; N:1,013]) and 92.77% ([S:87.77%; D:5.40%; F:2.93%; M:4.30%; N:1,536]) gene-complete for the **arthropoda_odb10** and **crustacea_odb12** BUSCO datasets, respectively. Of the 1.04 Gbp of total sequence, 906.6 Mbp (86.74%) was contained in 58 scaffolds larger than 1 Mbp. Moreover, the 16 largest sequences (larger than 8.5 Mbp) contain 841.2 Mbp (80.61%) of the total sequence, and likely represent 16 putative chromosomes. This karyotype number, N=16, is in agreement with that observed for the genus *Balanus* [Austin1958] and other acorn barnacle species in the family Balanidae [Han2024].

Genome annotation

Once the assembly and scaffolding was completed, we annotated and masked repeat sequences in the *B. glandula* reference assembly using the **EarlGrey** software [Baril2024]. This analysis identified 698.6 Mbp (66.94%) of the assembly as repetitive elements; however, the majority of these, 404.5 Mbp (38.76% of the assembly), remain unclassified among the major repeat element types. This large proportion of unclassified repeat elements is likely the product of the very small number of representative barnacle sequences among curated repeat reference libraries, reflecting barnacle’s status as a genomically understudied group. Among the classified repeats, DNA and LINE elements were the two most abundant repeats classes, comprising 8.45% and 9.16% of the total assembly, respectively.

Following repeat annotation and masking of the reference genome, we annotated protein-coding genes using short-read RNAseq and long-read PacBio Iso-Seq sequencing, combining and curating annotations generated across the two data types. This process resulted in 20,234 protein-coding genes annotated in the genome. The resulting annotation was 93.58% ([S:79.76%; D:13.82%; F:1.78%; M:4.64%; N:1,013]) and 84.77% ([S:72.66%; D:12.11%; F:4.10%; M:11.13%; N:1,536]) gene-complete when compared against the **arthropoda_odb10** and **crustacea_odb12** BUSCO datasets, respectively. Additionally, we compared our annotation against the proteome of four other barnacle species (see Methods) using **OrthoFinder** [Emms&Kelly2015; Emms&Kelly2019], resulting in the assignment of 16,978 (83.91%) *B. glandula* genes to orthologous groups across barnacles and the identification of 2,576 single-copy orthologs across all five species.

In addition to protein-coding genes, we also annotated non-coding elements in the *B. glandula* reference assembly. We identified 7,532 lncRNA, 1,203 tRNA, and 24 rRNA genes. In total, across protein-coding and non-coding transcripts, we were able to annotate 28,993 genes in the genome.

Conserved synteny across barnacle genome

We performed a conserved synteny analysis using the software **Genespace** to identify patterns of large-scale structure across barnacle genomes. We compared our *B. glandula* genome against the chromosome-level genome assemblies of *Amphibalanus amphitrite* (Balanidae) [Han2024] and *Pollicipes pollicipes* (Pollicipidae) [Bernot2022].

Discussion

Nulla ac nisl. Nullam urna nulla, ullamcorper in, interdum sit amet, gravida ut, risus. Aenean ac enim. In luctus. Phasellus eu quam vitae turpis viverra pellentesque. Duis feugiat felis ut enim. Phasellus pharetra, sem id porttitor sodales, magna nunc aliquet nibh, nec blandit nisl mauris at pede. Suspendisse risus risus, lobortis eget, semper at, imperdiet sit amet, quam. Quisque scelerisque dapibus nibh. Nam enim. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Nunc ut metus. Ut metus justo, auctor at, ultrices eu, sagittis ut, purus. Aliquam aliquam.




Figure 1: Conserved synteny across barnacle genomes.

Materials and Methods

Nulla ac nisl. Nullam urna nulla, ullamcorper in, interdum sit amet, gravida ut, risus. Aenean ac enim. In luctus. Phasellus eu quam vitae turpis viverra pellentesque. Duis feugiat felis ut enim. Phasellus pharetra, sem id porttitor sodales, magna nunc aliquet nibh, nec blandit nisl mauris at pede. Suspendisse risus risus, lobortis eget, semper at, imperdiet sit amet, quam. Quisque scelerisque dapibus nibh. Nam enim. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Nunc ut metus. Ut metus justo, auctor at, ultrices eu, sagittis ut, purus. Aliquam aliquam.

Data Availability

https://github.com/arcolon14/balanus_genome_ms

Acknowledgments

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Supplements