

Case study: Unveiling the seasonal dynamics of protist communities in tropical coastal ecosystems using long-read metabarcoding

BIO9905MERG1 – Bioinformatics for Environmental Sequencing

Denise Ong Rui Ying
University of Oslo (IBV, AKVA)
Genomics and Ecology of Eukaryotes Lab (GEEK Lab)
8 April 2025

About me


- Postdoctoral research fellow at University of Oslo, Norway
- PhD at Nanyang Technological University, Singapore

My research




- Marine planktonic microbial eukaryotes (protists)
- Biological carbon pump – carbon fixation and carbon export
- DNA metabarcoding


Protist communities are diverse in size




Protist communities are taxonomically diverse




Ecologically significant groups of coastal protists









Haptophytes











Climate seasonality affects the marine plankton community




Time series sampling

- Repeated sampling and measurements
 - Hourly
 - Daily
 - Weekly
 - Monthly
- Different methods to characterize the community
 - Microscopy
 - Flow cytometry cell counting
 - Chlorophyll a/photosynthetic pigment
 - DNA metabarcoding
- Temporal dynamics
 - Yearly (seasons), daily
 - Long-term
- What biotic and abiotic factors that drive the marine plankton community variation?

Temperate regions have strong seasonality





Temperate regions have strong seasonality




What about tropical regions?

Tropical coastal ecosystem – Singapore Strait






Monsoon seasons in the Singapore Strait



Monsoon season	Duration
Northeast monsoon	15 Nov to end Feb
Intermonsoon 1	01 Mar to 15 May
Southwest monsoon (coastal darkening)	15 May to 15 Sept
Intermonsoon 2	15 Sept to 15 Nov


Monsoon seasons in the Singapore Strait



Is the tropical coastal protist community seasonal?


Monthly sampling to collect surface seawater

- Five-year time-series
 - Feb 2018 to Dec 2022



Long-read metabarcoding

eukaryotic rRNA operon




- Increased genetic resolution (400 bp to 5000 bp)
- Differentiate between closely related clades

Covered tomorrow by Embla Stokke




Clarence Sim Ph.D., GEEK Lab


Bioinformatic process



Bioinformatic process – clustering ASVs




Overall protist community




Mostly stable community

Southwest monsoon community is distinct from other seasons




Is the protist community seasonal?



- Comparing the similarity (Bray-Curtis) of the community at different months sampled
- Mantel test; R=0.1556, p< 0.001
- Significant but weak correlation

How can we differentiate between seasonal and non-seasonal ASVs?



Recurrence Index from Giner et al. (2018)

Received: 11 May 2018 | Revised: 8 October 2018 | Accepted: 30 October 2018

DOI: 10.1111/mec.14929


FROM THE COVER | WILEY | MOLECULAR ECOLOGY


Quantifying long-term recurrence in planktonic microbial eukaryotes

Caterina R. Giner¹ | Vanessa Balagué¹ | Anders K. Krabberød² | Isabel Ferrera¹ | Albert Reñé¹ | Esther Garcés¹ | Josep M. Gasol¹ | Ramiro Logares^{1,2} | Ramon Massana¹


Autocorrelation

- Linear relationship between lagged values of a time series
- Correlation between the relative abundance of an ASV against its lagged values (1 month apart, etc.)
- At every time interval...






How can we differentiate between seasonal and non-seasonal ASVs?




Using autocorrelation to calculate the Recurrence Index






RAD-A - 663dd8632e - non-seasonal - RI 1.05




A small proportion of the community is seasonal



Different types of seasonality


Long-term

Mantoniella sp.



Pulses

Micromonas bravo




Different types of seasonality

Long-term

Mantoniella sp.


Mamiellophyceae - b7fb08b504 - seasonal - RI 1.38



Pulses


Micromonas bravo

Mamiellophyceae - 079e786a79 - seasonal - RI 1.32




Seasonal ASVs: Long term vs pulses


• Long term ASVs: present in more than ¾ of the samples



More abundant taxa have more seasonal ASVs




Seasonality does not dominate in any one taxa




Why long-read metabarcoding?

Bathycoccus




Newly proposed clade present in the Singapore Strait

Bathycoccus sp., Clade III



Mamiellophyceae - 5744d6a833 - seasonal - RI 1.47



Key takeaways

- The overall community has very little seasonality
- Identify seasonality for a small proportion of ASVs
- Understanding at a lower taxonomic level might help us to better identify the causes of plankton blooms (HABs, Noctiluca scintillans) in Singapore coastal waters
- Better classification of the plankton community with long-read metabarcoding

Acknowledgements







- Adriana Lopes dos Santos (UiO)
- Clarence Sim (NTU)
- Daniel Vaulot (CNRS/UiO)
- Avneet Kaur
- Rae Chua
- Patrick Martin (NTU)
- Woo Oon Yee (NTU)

Funding:

Singapore Ministry of Education, Academic Research Fund Tier 1