

Today:

11.4

] Review problem]

tangent plane
Linear Approximation
Differentials


$f(x, y)$

11.6

] Chain rule

$$df = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$$

EXAMPLE 4 The base radius and height of a right circular cone are measured as 10 cm and 25 cm, respectively, with a possible error in measurement of as much as 0.1 cm in each. Use differentials to estimate the maximum error in the calculated volume of the cone.



$$V(r, h) = \frac{1}{3} \pi r^2 h$$

$$dV = ??$$

$$dV = \frac{\partial V}{\partial r} dr + \frac{\partial V}{\partial h} dh$$

$$dr = dh = 0.1$$

$$dV = \frac{2}{3} \pi r h dr + \frac{1}{3} \pi r^2 dh$$

$$\begin{aligned} r &= 10, \quad h = 25 \\ dr &= dh = 0.1 \\ &= 20\pi \end{aligned}$$

$$\begin{aligned} \frac{dV}{V} &= ?? \% \\ &= \frac{3}{125} \\ &\approx 0.024 \\ &= 2.4 \% \end{aligned}$$

$$f = f(x, y)$$

$$df = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$$

1-6 ■ Find an equation of the tangent plane to the given surface at the specified point.

$$z = y \cos(x - y), \quad (2, 2, 2)$$


$$z - z_0 = \frac{\partial z}{\partial x} (x - x_0) + \frac{\partial z}{\partial y} (y - y_0)$$

$$\frac{\partial z}{\partial x} = -y \sin(x - y) \Big|_{\substack{x=2 \\ y=2}} = 0$$

$$\frac{\partial z}{\partial y} = [y \sin(x - y) + \cos(x - y)] \Big|_{\substack{x=2 \\ y=2}} = 1$$

$$z - 2 = 0(x - 2) + 1(y - 2)$$

$$\boxed{z = y}$$



30. The pressure, volume, and temperature of a mole of an ideal gas are related by the equation $PV = 8.31T$, where P is measured in kilopascals, V in liters, and T in kelvins. Use differentials to find the approximate change in the pressure if the volume increases from 12 L to 12.3 L and the temperature decreases from 310 K to 305 K.

$$P = 8.31T/V$$

$$dP = ?? \quad \text{given} \quad V = 12, \quad T = 310, \quad dV = 0.3, \quad dT = -5$$


$$dP = \frac{\partial P}{\partial V} dV + \frac{\partial P}{\partial T} dT$$

$$= -8.8$$




11.5

THE CHAIN RULE



$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial t}$$





$$\frac{\partial f}{\partial u} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u}$$

$$\frac{\partial f}{\partial v} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial v}$$

$$\frac{\partial f}{\partial s} = \frac{\partial f}{\partial y} \frac{\partial y}{\partial s}$$

Q:

$T(x, y) = x + y$
Temperature at point (x, y)



$$x = \cos(t)$$

$$y = \sin(t)$$

$$0 < t < \infty$$

what path is this ??

$$\vec{r}(t) = \cos t \hat{i} + \sin t \hat{j}$$

Q: rate of change of temperature T
w.r.t. time t

$$\frac{dT}{dt} = ??$$



$$\text{at } t = \pi/2$$

$$= \frac{\partial T}{\partial x} \frac{dx}{dt} + \frac{\partial T}{\partial y} \frac{dy}{dt}$$

$$= \left[1(-\sin t) + 1 \cdot \cos(t) \right]_{t=\pi/2}$$


$$= -1$$

EXAMPLE 1 If $z = x^2y + 3xy^4$, where $x = \sin 2t$ and $y = \cos t$, find dz/dt when $t = 0$.



$$x = \sin 2t$$

$$y = \cos t$$



$$\frac{dz}{dt} = \frac{\partial z}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt}$$


$$= (2xy + 3y^4)(2\cos 2t) + (x^2 + 12xy^3)(-\sin t)$$

$$t = 0, \quad x = 0, \quad y = 1$$

$$= 6$$

EXAMPLE 3 If $z = e^x \sin y$, where $x = st^2$ and $y = s^2t$, find $\partial z / \partial s$ and $\partial z / \partial t$.


$$z = e^x \sin y$$



$$\frac{\partial z}{\partial s} = e^x \sin y \cdot t^2 + e^x \cos y \cdot 2st$$

$$\frac{\partial z}{\partial t} = e^x \sin y \cdot 2st + e^x \cos y \cdot s^2$$

V EXAMPLE 5 If $u = x^4y + y^2z^3$, where $x = rse^t$, $y = rs^2e^{-t}$, and $z = r^2s \sin t$, find the value of $\partial u / \partial s$ when $r = 2$, $s = 1$, $t = 0$.



$$\frac{\partial u}{\partial s} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial s} + \frac{\partial u}{\partial z} \frac{\partial z}{\partial s}$$

$$= 4x^3y r e^t + (x^4 + 2yz^3) r s e^{-t} + (2y^2z^2) r^2 \sin t$$

$$r=2, s=1, t=0$$

$$x=2, y=2, z=0$$


$$= 192$$

Today:

→ Review : chain rule

→ 11.6 : Directional Derivatives
& gradients

32. The radius of a right circular cone is increasing at a rate of 1.8 in/s while its height is decreasing at a rate of 2.5 in/s. At what rate is the volume of the cone changing when the radius is 120 in. and the height is 140 in.?



$$\frac{dh}{dt} = -2.5$$

$$\frac{dr}{dt} = 1.8$$

$$V = \frac{1}{3} \pi r^2 h$$


$$\left. \frac{dV}{dt} \right|_{\substack{r=120 \\ h=140}} = ??$$


$$= \frac{\partial V}{\partial r} \frac{dr}{dt} + \frac{\partial V}{\partial h} \frac{dh}{dt}$$

$$= \frac{2}{3} \pi r h \frac{dr}{dt} + \frac{\pi r^2}{3} \frac{dh}{dt}$$

$$= \frac{2}{3} \pi (120)(140)(1.8) + \frac{\pi}{3} (120)^3 (-2.5)$$

$$= 8160 \pi \text{ in}^3/\text{s}$$






$$\frac{\partial f}{\partial u} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u}$$


$$\frac{\partial f}{\partial v} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v}$$

29. The temperature at a point (x, y) is $T(x, y)$, measured in degrees Celsius. A bug crawls so that its position after t seconds is given by $x = \sqrt{1+t}$, $y = 2 + \frac{1}{3}t$, where x and y are measured in centimeters. The temperature function satisfies $T_x(2, 3) = 4$ and $T_y(2, 3) = 3$. How fast is the temperature rising on the bug's path after 3 seconds?



$$t = 3 \quad x = 2 \quad y = 3$$

$$\begin{aligned} \left. \frac{dT}{dt} \right|_{t=3} &= \frac{\partial T}{\partial x} \frac{dx}{dt} + \frac{\partial T}{\partial y} \frac{dy}{dt} \\ &= \left(4 \cdot \frac{1}{2\sqrt{1+t}} + 3 \cdot \frac{1}{3} \right) \bigg|_{t=3} \\ &= 1 + 1 = 2 \end{aligned}$$



- 29.** The temperature at a point (x, y) is $T(x, y)$, measured in degrees Celsius. A bug crawls so that its position after t seconds is given by $x = \sqrt{1+t}$, $y = 2 + \frac{1}{3}t$, where x and y are measured in centimeters. The temperature function satisfies $T_x(2, 3) = 4$ and $T_y(2, 3) = 3$. How fast is the temperature rising on the bug's path after 3 seconds?

- 29.** The temperature at a point (x, y) is $T(x, y)$, measured in degrees Celsius. A bug crawls so that its position after t seconds is given by $x = \sqrt{1+t}$, $y = 2 + \frac{1}{3}t$, where x and y are measured in centimeters. The temperature function satisfies $T_x(2, 3) = 4$ and $T_y(2, 3) = 3$. How fast is the temperature rising on the bug's path after 3 seconds?


11.6

DIRECTIONAL DERIVATIVES AND THE GRADIENT VECTOR

2 DEFINITION The directional derivative of f at (x_0, y_0) in the direction of a unit vector $\mathbf{u} = \langle a, b \rangle$ is

$$D_{\mathbf{u}}f(x_0, y_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + ha, y_0 + hb) - f(x_0, y_0)}{h}$$

if this limit exists.



• blue surface is the graph of $f(x, y)$


• base point $(x_0, y_0) = P'$

• \hat{u} : directional derivative at point P' in the direction \hat{u}

= rate of change of f if we start moving from point P' along the direction \hat{u}

fixed direction

d := sketch the line passing through P & parallel to \hat{u}



$$\hat{u} = a\hat{i} + b\hat{j}$$

3 THEOREM If f is a differentiable function of x and y , then f has a directional derivative in the direction of any unit vector $\mathbf{u} = \langle a, b \rangle$ and

$$D_{\mathbf{u}}f(x, y) = f_x(x, y)a + f_y(x, y)b$$

$$D_{\mathbf{u}}f(x, y) = \lim_{h \rightarrow 0} \frac{f[(x, y) + h(x, y)] - f(x, y)}{h} = \frac{\partial f}{\partial x} a + \frac{\partial f}{\partial y} b$$


$$= \underbrace{\left(\frac{\partial f}{\partial x} \hat{i} + \frac{\partial f}{\partial y} \hat{j} \right)}_{\text{gradient } \nabla f} \cdot \hat{\mathbf{u}}$$

ok??

$$D_{\mathbf{u}}f(x, y) = \nabla f \cdot \hat{\mathbf{u}}$$

$$\nabla f = \frac{\partial f}{\partial x} \hat{i} + \frac{\partial f}{\partial y} \hat{j}$$

EXAMPLE 1 Find the directional derivative $D_{\mathbf{u}}f(x, y)$ if $f(x, y) = x^3 - 3xy + 4y^2$ and \mathbf{u} is the unit vector given by angle $\theta = \pi/6$. What is $D_{\mathbf{u}}f(1, 2)$?



$$D_{\mathbf{u}}f(1, 2) = \nabla f \cdot \hat{\mathbf{u}}$$

$$\nabla f = \frac{\partial f}{\partial x} \hat{\mathbf{i}} + \frac{\partial f}{\partial y} \hat{\mathbf{j}}$$

$$= (3x^2 - 3y) \hat{\mathbf{i}} + (-3x + 8y) \hat{\mathbf{j}}$$

$$\nabla f(1, 2) = -3 \hat{\mathbf{i}} + 13 \hat{\mathbf{j}}$$

$$\hat{\mathbf{u}} = ?? = \cos \frac{\pi}{6} \hat{\mathbf{i}} + \sin \frac{\pi}{6} \hat{\mathbf{j}}$$

$$= \frac{\sqrt{3}}{2} \hat{\mathbf{i}} + \frac{1}{2} \hat{\mathbf{j}}$$

$$D_{\mathbf{u}}f(1, 2) = (-3 \hat{\mathbf{i}} + 13 \hat{\mathbf{j}}) \cdot \left(\frac{\sqrt{3}}{2} \hat{\mathbf{i}} + \frac{1}{2} \hat{\mathbf{j}} \right)$$

$$= \frac{(3 - 3\sqrt{3})}{2}$$

8 DEFINITION If f is a function of two variables x and y , then the **gradient** of f is the vector function ∇f defined by

$$\nabla f(x, y) = \langle f_x(x, y), f_y(x, y) \rangle = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j}$$

next time

V EXAMPLE 3 Find the directional derivative of the function $f(x, y) = x^2y^3 - 4y$ at the point $(2, -1)$ in the direction of the vector $\mathbf{v} = 2\mathbf{i} + 5\mathbf{j}$.

V EXAMPLE 4 If $f(x, y, z) = x \sin yz$, (a) find the gradient of f and (b) find the directional derivative of f at $(1, 3, 0)$ in the direction of $\mathbf{v} = \mathbf{i} + 2\mathbf{j} - \mathbf{k}$.

MAXIMIZING THE DIRECTIONAL DERIVATIVE

EXAMPLE 5

- (a) If $f(x, y) = xe^y$, find the rate of change of f at the point $P(2, 0)$ in the direction from P to $Q(\frac{1}{2}, 2)$.
- (b) In what direction does f have the maximum rate of change? What is this maximum rate of change?

EXAMPLE 6 Suppose that the temperature at a point (x, y, z) in space is given by $T(x, y, z) = 80/(1 + x^2 + 2y^2 + 3z^2)$, where T is measured in degrees Celsius and x, y, z in meters. In which direction does the temperature increase fastest at the point $(1, 1, -2)$? What is the maximum rate of increase?

