

Stability of Bianchi attractors in gauged supergravity

Karthik Inbasekar

Tata Institute of Fundamental Research, Mumbai

National String Meeting, Dec 2013

Ref: [1206.3887](#), [1307.1314](#)

Plan of the talk

Introduction

Motivation

Overview

Bianchi Attractors

Symmetries

Example

Stability of Bianchi attractors in gauged sugra

Gauged Supergravity

Linearised equations

Stability conditions

Stable Bianchi Attractors

Summary and Future outlook

Stability of Bianchi
attractors

Introduction

Motivation

Overview

Bianchi Attractors

Stability

Summary and
Future outlook

- In gauge/gravity correspondence, black branes are **holographic duals** to field theories at finite temperature.
- **Extremal** branes exhibit vanishing entropy density at zero temperature and describe ground states of the field theory.
- **Bianchi attractors**: Classification of **homogeneous anisotropic extremal black brane horizons** in $d = 5$.
- Lifshitz, Schrödinger geometries belong to Bianchi Type I in the classification.
- Appear as **generalised attractor** solutions in extensions of attractor mechanism to **gauged supergravity**.

Cacciatori-Klemm , Kachru-Kallosh-Shmakova

Generalised attractors

Introduction

Motivation
Overview

Bianchi Attractors

Stability

Summary and
Future outlook

- **Generalised attractors**: solutions to equations of motion that reduce to algebraic equations when all the fields, curvature tensors are constants in tangent space.
- Gauge field, and Einstein equations reduce to **algebraic equations** at the **attractor point**.
- Scalar field equations reduce to a minimisation condition on an **attractor potential**.
- **Generalised attractor geometries** are characterised by **constant anholonomy** coefficients.

$$[e_a, e_b] = c_{ab}^c e_c ; \quad e_a \equiv e_a^\mu \partial_\mu$$

$$c_{ab}^c = e_a^\mu e_b^\nu (\partial_\mu e_\nu^c - \partial_\nu e_\mu^c)$$

- **Bianchi attractors** have constant anholonomy by construction.

Bianchi Attractors in gauged supergravity

- Bianchi type metrics can be easily realised in simple Einstein-Maxwell systems with **massive gauge fields**.

- Typical scalar kinetic term of gauged supergravities,

$$g_{xy} \mathcal{D}_\mu \phi^x \mathcal{D}^\mu \phi^y; \quad \mathcal{D}_\mu \phi^x \equiv \partial_\mu \phi^x + g A_\mu^I K_I^x(\phi).$$

- At **attractor** points **scalars are constant**, terms like

$$g^2 g_{xy} K_I^x K_J^y A_\mu^I A^{J\mu}$$

act as **effective mass term** for the **gauge field**.

- Several **Bianchi attractors** were embedded in gauged supergravity as **generalised attractors**. Inbasekar-Tripathy

Motivation: Stability of Bianchi metrics

- **Instabilities** due to scalar fluctuations exist in large class of such metrics. Donos-Gauntlett-Pantelidou, Cremonini-Sinkovics, Andrade-Ross, Keeler
- Studying such instabilities help in understanding how the **geometry might get corrected in the IR**.
- **A common recipe** to study the stability of Bianchi type metrics will be useful.
- Embedding Bianchi type metrics as generalised attractors in gauged supergravity provided an ideal platform for this study.

Motivation: Stability of generalised attractors

- Generalised attractor analysis does not involve susy, relies on extremisation of an attractor potential.
- Solutions were found at critical points, not at absolute minima of attractor potential.
- Preliminary susy analysis of existing solutions using KSI indicated broken susy.
- Non-susy attractors can be unstable to scalar fluctuations about critical value.

- Analyse the **stability of Bianchi attractors** in gauged supergravity under scalar **fluctuations about the attractor value**.
- Examine the field equations at the **linearised level** and demand that **fluctuations vanish near the horizon**.
- Determine **conditions of stability**.
- Identify the class of Bianchi attractors which satisfy the condition.

Bianchi Attractors

Stability of Bianchi
attractors

Introduction

Motivation

Overview

Introduction

Bianchi Attractors

Symmetries

Example

Stability

Summary and
Future outlook

Bianchi Attractors

Symmetries

Example

Stability of Bianchi attractors in gauged sugra

Gauged Supergravity

Linearised equations

Stability conditions

Stable Bianchi Attractors

Summary and Future outlook

Bianchi Attractors: Symmetries

- **Bianchi Attractors:** Five dimensional **extremal black brane horizons** with **homogeneous symmetries** in spatial directions. **Iizuka-Kachru-Kundu-Narayan-Sircar-Trivedi**
- **Homogeneous symmetries:** **invariant basis** $\tilde{e}_i, i = 1, 2, 3$ that commutes with Killing vectors.

$$[\xi_j, \tilde{e}_i] = 0, \quad [\tilde{e}_i, \tilde{e}_j] = c_{ij}{}^k \tilde{e}_k$$

- **Invariant vectors** close to form a **Lie algebra** - isomorphic to **Bianchi classification** (I-IX) of 3d real Lie algebras **Bianchi**.
- Metric written in terms of **invariant one forms** ω^i dual to \tilde{e}_i displays **manifest homogeneous symmetries**.

$$d\omega^k = \frac{1}{2} c_{ij}{}^k \omega^i \wedge \omega^j$$

- Additional symmetries: **scale invariance**, **time translation invariance**

$$\hat{r} \rightarrow \lambda \hat{r} , \quad \hat{t} \rightarrow \lambda^{-u_0} \hat{t} , \quad \omega^i \rightarrow \lambda^{-u_i} \omega^i$$

- Fix the form of the metric completely.

$$ds^2 = L^2 \left[-\hat{r}^{2u_0} d\hat{t}^2 + \frac{d\hat{r}^2}{\hat{r}^2} + \hat{r}^{2(u_i+u_j)} \eta_{ij} \omega^i \otimes \omega^j \right]$$

- Have **constant anholonomy coefficients** by construction.

Example: Bianchi II

- One forms, invariant vectors, structure constants,

$$c_{23}^1 = 1 = -c_{32}^1,$$

$$\xi_1 = \partial_{\hat{y}}, \quad \tilde{e}_1 = \partial_{\hat{y}}, \quad \omega^1 = d\hat{y} - \hat{x}d\hat{z}, \quad d\omega^1 = \omega^2 \wedge \omega^3,$$

$$\xi_2 = \partial_{\hat{z}}, \quad \tilde{e}_2 = \hat{x}\partial_{\hat{y}} + \partial_{\hat{z}}, \quad \omega^2 = d\hat{z}, \quad d\omega^2 = 0,$$

$$\xi_3 = \partial_{\hat{x}} + \hat{z}\partial_{\hat{y}}, \quad \tilde{e}_3 = \partial_{\hat{x}}, \quad \omega^3 = d\hat{x}, \quad d\omega^3 = 0$$

- scaling in coordinates,

$$(\hat{x}, \hat{y}, \hat{z}) \rightarrow (\lambda^{-u_1} \hat{x}, \lambda^{-(u_1+u_3)} \hat{y}, \lambda^{-u_3} \hat{z})$$

- scaling in one forms,

$$(\omega^1, \omega^2, \omega^3) \rightarrow (\lambda^{-(u_1+u_3)} \omega^1, \lambda^{-u_3} \omega^2, \lambda^{-u_1} \omega^3)$$

- metric

$$ds^2 = L^2 \left[-\hat{r}^{2u_0} d\hat{t}^2 + \frac{d\hat{r}^2}{\hat{r}^2} + \hat{r}^{2(u_1+u_3)} (\omega^1)^2 + \hat{r}^{2u_3} (\omega^2)^2 + \hat{r}^{2u_1} (\omega^3)^2 \right]$$

Example: Bianchi IX

- Invariant one forms,

$$\begin{aligned}\omega^1 &= -\sin(\hat{z})d\hat{x} + \sin(\hat{x})\cos(\hat{z})d\hat{y}, & d\omega^1 &= \omega^2 \wedge \omega^3, \\ \omega^2 &= \cos(\hat{z})d\hat{x} + \sin(\hat{x})\sin(\hat{z})d\hat{y}, & d\omega^2 &= \omega^3 \wedge \omega^1, \\ \omega^3 &= \cos(\hat{x})d\hat{y} + d\hat{z}, & d\omega^3 &= \omega^1 \wedge \omega^2\end{aligned}$$

- no scaling symmetry in $(\hat{x}, \hat{y}, \hat{z})$ coordinates and one forms

$$ds^2 = L^2 \left[-\hat{r}^{2u_0} d\hat{t}^2 + \frac{d\hat{r}^2}{\hat{r}^2} + (\omega^1)^2 + (\omega^2)^2 + (\omega^3)^2 \right]$$

- Metric can be rewritten in direct product form

$$AdS_2 \times M_{IX}$$

$$ds^2 = L_1^2 \left(-\tilde{r}^2 d\tilde{t}^2 + \frac{d\tilde{r}^2}{\tilde{r}^2} \right) + L_2^2 \left((\omega^1)^2 + (\omega^2)^2 + (\omega^3)^2 \right)$$

Stability of Bianchi attractors in gauged sugra

Introduction

Motivation

Overview

Bianchi Attractors

Symmetries

Example

Stability of Bianchi attractors in gauged sugra

Gauged Supergravity

Linearised equations

Stability conditions

Stable Bianchi Attractors

Summary and Future outlook

- The most general $\mathcal{N} = 2, d = 5$ gauged sugra has gravity coupled to vector, tensor and hypermultiplets.

Ceresole-Dall'Agata

- The scalars in the theory parametrise a manifold that factorises into a direct product of a **very special** and **quaternionic manifold**,

$$\mathcal{M}_{scalar} = \mathcal{S}(n_V + n_T) \otimes \mathcal{Q}(n_H).$$

$$C_{IJK} h^I h^J h^K = 1, \quad h^I \equiv h^I(\phi).$$

- The **R symmetry** group is $SU(2)_R$.
- We consider the truncated theory with vector multiplets, abelian gauging of the isometries of the scalar manifold and gauging of $U(1)_R$ symmetry.

Gauged Sugra: Gauging the Symmetries

- Gauging: Suitable subgroup K of the isometry group G of the full scalar manifold \mathcal{M}_{scalar} , and the $U(1)_R$ symmetry group.
- Ordinary derivatives on scalar and fermions are replaced with K -covariant derivatives.

$$\partial_\mu \phi^x \rightarrow \mathcal{D}_\mu \phi^x \equiv \partial_\mu \phi^x + g A_\mu^I K_I^x(\phi)$$

- Gauging the $U(1)_R$ Symmetry:

$$\nabla_\mu \psi_{\nu i} \rightarrow \nabla_\mu \psi_{\nu i} + ig_R A_\mu^I V_I \psi_{\nu i}.$$

- Gauging leads to scalar potentials in the theory - possibility of AdS vacuum.

- The bosonic part of the Lagrangian:

$$\hat{e}^{-1} \mathcal{L}_{Bosonic}^{\mathcal{N}=2} = -\frac{1}{2}R - \frac{1}{4}a_{IJ}F_{\mu\nu}^I F^{J\mu\nu} - \frac{1}{2}g_{xy}\mathcal{D}_\mu\phi^x\mathcal{D}^\mu\phi^y \\ - \mathcal{V}(\phi) + \frac{\hat{e}^{-1}}{6\sqrt{6}}C_{IJK}\epsilon^{\mu\nu\rho\sigma\tau}F_{\mu\nu}^I F_{\rho\sigma}^J A_\tau^K$$

- The potential in this case comes only from the R symmetry gauging,

$$\mathcal{V}(\phi) = -g_R^2[2P_{ij}P^{ij} - P_{ij}^x P^{xij}], \\ P_i{}^k \equiv h^I V_I \delta_i{}^k, \quad P_i{}^x{}^k \equiv h^{xI} V_I \delta_i{}^k.$$

Fluctuations about attractor value

- Fluctuations

$$\phi_c^x + \epsilon \delta \phi^x,$$

$$A_{\mu}^I + \epsilon \delta A_{\mu}^I,$$

$$g_{\mu\nu} + \epsilon \gamma_{\mu\nu},$$

- Gauge field equations,

$$\begin{aligned} a_{IJ}|_{\phi_c} \nabla_{\mu} F_{\delta}^{I\mu\nu} - g^2 K_{IJ}|_{\phi_c} \delta A^{\nu J} = \\ - \left(\frac{\partial a_{IJ}}{\partial \phi^z} \Big|_{\phi_c} \nabla_{\mu} (F^{I\mu\nu} \delta \phi^z) - g^2 \frac{\partial K_{IJ}}{\partial \phi^z} \Big|_{\phi_c} \delta \phi^z A^{\nu J} \right) \\ + g K_{Iy}|_{\phi_c} \partial^{\nu} \delta \phi^y \end{aligned}$$

- **Regularity** of the gauge fields requires well behaved scalar fluctuations near the horizon.

Linearised Einstein equations

- Linearised Einstein equation,

$$\nabla^\alpha \nabla_\alpha \bar{\gamma}_{\mu\nu} + 2R_{(\mu}^{\alpha}{}_{\nu)}{}^{\beta} \bar{\gamma}_{\beta\alpha} - 2R_{(\mu}^{\beta} \bar{\gamma}_{\nu)\beta} + g_{\mu\nu} (R_{\alpha\beta} \bar{\gamma}^{\alpha\beta} + \frac{2}{2-D} R \bar{\gamma}) + R \bar{\gamma}_{\mu\nu} \\ + 2\dot{T}_{\mu\nu}^{attr} (g_{\alpha\beta} + \epsilon \gamma_{\alpha\beta})|_{\epsilon=0} + 2\dot{T}_{\mu\nu} (\phi_c + \epsilon \delta\phi)|_{\epsilon=0} = 0$$

- Stress energy dependence on $\gamma_{\mu\nu}$ and $\delta\phi^z$

$$\dot{T}_{\mu\nu}^{attr} (g_{\alpha\beta} + \epsilon \gamma_{\alpha\beta})|_{\epsilon=0} = \mathcal{V}_{attr}(\phi_c) (\bar{\gamma}_{\mu\nu} + \frac{2\bar{\gamma}}{2-D} g_{\mu\nu}) \\ - (\bar{\gamma}_{\lambda\sigma} + \frac{\bar{\gamma}}{2-D} g_{\lambda\sigma}) (\frac{1}{2} T_{attr}^{\lambda\sigma} g_{\mu\nu} + a_{IJ}|_{\phi_c} F_{\mu}^I{}^{\lambda} F_{\nu}^J{}^{\sigma})$$

$$\dot{T}_{\mu\nu} (\phi_c + \delta\phi)|_{\epsilon=0} = T_{\mu\nu}^{attr}|_{\phi_c} \\ + g K_{yI}|_{\phi_c} \left(A^{\lambda I} \partial_\lambda (\delta\phi^y) g_{\mu\nu} - A_{\mu}^I \partial_\nu (\delta\phi^y) - A_{\nu}^I \partial_\mu (\delta\phi^y) \right) \\ - \left[\frac{\partial a_{IJ}}{\partial \phi^z} \right]_{\phi_c} F_{\mu\lambda}^I F_{\nu}^J{}^{\lambda} + g^2 \frac{\partial K_{IJ}}{\partial \phi^z} \Big|_{\phi_c} A_{\mu}^I A_{\nu}^J \Big] \delta\phi^z$$

Stress energy tensor: Backreaction at first order

- For Gauged sugra with generic gauging, trace of Einstein equation,

$$R(g_{\mu\nu}, \gamma_{\mu\nu}) \frac{(2-D)}{2} = T_{\mu}^{\text{attr}\mu}|_{\phi_c} + (D-2)g K_{yI}|_{\phi_c} A^{\lambda I} \partial_{\lambda}(\delta\phi^y) \\ + g^2 \frac{\partial K_{IJ}}{\partial \phi^z} \Big|_{\phi_c} A_{\mu}^I A^{J\mu} \delta\phi^z$$

$$T_{\mu}^{\text{attr}\mu}|_{\phi_c} = \mathcal{V}_{\text{attr}}(\phi_c) D - \left[a_{IJ}|_{\phi_c} F_{\mu\nu}^I F^{\mu\nu J} + g^2 K_{IJ}|_{\phi_c} A_{\mu}^I A^{\mu J} \right]$$

$$K_{IJ} = g_{xy} K_I^x K_J^y$$

- Scalar fluctuation terms indicate backreaction even at first order perturbation.
- Relevant boundary conditions for scalars should be such that they are well behaved near the horizon.
- For $U(1)_R$ gauging, $g = 0$ and back reaction is absent.

Scalar fluctuation equations

- Scalar fluctuation equation for arbitrary gauged sugra,

$$\nabla_\mu \nabla^\mu \delta\phi^x - g^{zx} \frac{\partial^2 \mathcal{V}_{attr}}{\partial \phi^z \partial \phi^y} \Big|_{\phi_c} \delta\phi^y + 2g (g^{zx} \tilde{\nabla}_y K_{|z})|_{\phi_c} A^{\mu l} \nabla_\mu \delta\phi^y = 0$$

$\tilde{\nabla}$ - covariant derivative w.r.t g_{xy} .

∇ - covariant derivative w.r.t near horizon metric.

- higher order metric/gauge field fluctuations can be ignored for solving the above equation at lowest order.
- Laplacian for any given 5d Bianchi type metric,

$$\nabla_\mu \nabla^\mu = \frac{1}{L^2} \left[\hat{r}^2 \partial_{\hat{r}}^2 + (m+2) \hat{r} \partial_{\hat{r}} - \frac{1}{\hat{r}^{2u_0}} \partial_{\hat{t}}^2 \right]$$

$$m = -1 + \sum_I c_I u_I, \quad c_I > 0, \quad c_0 = 1.$$

Scalar fluctuation equations

- For the specific gauged supergravity model fluctuation equation reduce to ,

$$\left[\hat{r}^2 \partial_{\hat{r}}^2 + (m+2) \hat{r} \partial_{\hat{r}} - \frac{1}{\hat{r}^{2u_0}} \partial_{\hat{t}}^2 - \lambda \right] \delta\phi^x = 0$$

λ - **Eigenvalue** of double derivative of attractor potential.

Sign of λ - indicates nature of critical point.

- For ansatz $\delta\phi(\hat{r}, \hat{t}) = f(\hat{r})e^{ik\hat{t}}$ (with k real), we get **Bessel equation**

$$\left[\hat{r}^2 \partial_{\hat{r}}^2 + (m+2) \hat{r} \partial_{\hat{r}} + \left(\frac{k^2}{\hat{r}^{2u_0}} - \lambda \right) \right] f(\hat{r}) = 0$$

Scalar fluctuations

- Scalar fluctuations

$$f(X) = \left(\frac{X}{2}\right)^{\nu_0} \left[C_1 H_{\nu_\lambda}^1(X) [\Gamma(1 - \nu_\lambda) e^{i\nu_\lambda \pi} + \Gamma(1 + \nu_\lambda)] \right. \\ \left. + C_2 H_{\nu_\lambda}^2(X) [\Gamma(1 - \nu_\lambda) e^{-i\nu_\lambda \pi} + \Gamma(1 + \nu_\lambda)] \right]$$

$$X = \frac{k}{u_0 \hat{r}^{u_0}}, \quad \nu_\lambda = \frac{\sqrt{(1+m)^2 + 4\lambda}}{2u_0}, \quad \nu_0 = \frac{(1+m)}{2u_0}$$

- Consistency condition for ν_λ real,

$$\nu_\lambda = \frac{\sqrt{(1+m)^2 + 4\lambda}}{2u_0} = \frac{\sqrt{(\sum_I c_I u_I)^2 + 4\lambda}}{2u_0} \leq 1$$

- implies $\lambda < 0$,

$$-\frac{(\sum_I c_I u_I)^2}{4} \leq \lambda < 0$$

- Scalar fluctuations - well defined for critical points which are maxima of attractor potential.

Conditions for stability

- In our coordinate system horizon is located at $\hat{r} = 0$, $X \simeq 1/\hat{r}$, consider **asymptotic expansion** of $f(X)$

$$f(X) \sim \left(\frac{X}{2}\right)^{\nu_0 - \frac{1}{2}} \sqrt{\frac{1}{\pi}} \left[C_1 e^{i(X - \frac{\pi}{2}(\nu_\lambda + \frac{1}{2}))} [\Gamma(1 - \nu_\lambda) e^{i\nu_\lambda \pi} + \Gamma(1 + \nu_\lambda)] \right. \\ \left. + C_2 e^{-i(X - \frac{\pi}{2}(\nu_\lambda + \frac{1}{2}))} [\Gamma(1 - \nu_\lambda) e^{-i\nu_\lambda \pi} + \Gamma(1 + \nu_\lambda)] \right]$$

- **Leading divergent term is absent only when,**

$$\nu_0 = \frac{(1+m)}{2u_0} = \frac{\sum_I c_I u_I}{2u_0} \leq \frac{1}{2}$$

- since $c_0 = 1$,

$$\sum_{I, I \neq 0} c_I u_I \leq 0$$

- But $u_I \geq 0$ for regular horizon, therefore **stability conditions** are:

$$u_0 \neq 0, \quad u_I = 0 \quad \forall I \neq 0$$

Stable Bianchi attractors

- Bianchi attractors with scale invariance in all directions,

$$ds^2 = L^2 \left[-\hat{r}^{2u_0} d\hat{t}^2 + \frac{d\hat{r}^2}{\hat{r}^2} + \hat{r}^{2(u_i+u_j)} \eta_{ij} \omega^i \otimes \omega^j \right]$$

- Stability condition,

$$u_0 \neq 0, \quad u_I = 0 \quad \forall I \neq 0$$

- Stable Bianchi attractors in gauged supergravity are a subclass with scale invariance only in \hat{r} and \hat{t} .

$$ds^2 = L^2 \left(-\hat{r}^{2u_0} d\hat{t}^2 + \frac{d\hat{r}^2}{\hat{r}^2} \right) + L^2 (\eta_{ij} \omega^i \otimes \omega^j)$$

- They are of the direct product form $AdS_2 \times M$.

$$ds^2 = L_1^2 \left(-\tilde{r}^2 d\tilde{t}^2 + \frac{d\tilde{r}^2}{\tilde{r}^2} \right) + L_2^2 (\eta_{ij} \omega^i \otimes \omega^j)$$

Stable Bianchi attractors

- Unstable generalised attractors

Geometry	λ	u_0	$u_l, l \neq 0$
Lifshitz	-34	3	1
Bianchi II	$-\frac{22}{3}$	$\sqrt{2}$	$u_1 = u_3 = \frac{1}{2\sqrt{2}}$
Bianchi VI $h < 0$	$-1 + \frac{14h}{3} - h^2$	$\frac{1}{\sqrt{2}}(1 - h)$	$u_2 = -\frac{1}{\sqrt{2}}h, u_3 = \frac{1}{\sqrt{2}}$

- Stable generalised attractors in direct product form

Geometry	λ	u_0	$u_l, l \neq 0$
$Lif_{u_0}(2) \times M_I$	$-\frac{5u_0^2}{3}$	any $u_0 > 0$	0
$AdS_2 \times M_I$	$-\frac{5}{3}$	1	0
$Lif_{u_0}(2) \times M_{II}$	$-\frac{61}{6}$	$\sqrt{\frac{11}{2}}$	0
$Lif_{u_0}(2) \times M^*$	$\lambda < 0$	any $u_0 > 0$	0

Summary and Future outlook

Stability of Bianchi
attractors

Introduction

Motivation

Overview

Introduction

Bianchi Attractors

Stability

Summary and
Future outlook

Bianchi Attractors

Symmetries

Example

Stability of Bianchi attractors in gauged sugra

Gauged Supergravity

Linearised equations

Stability conditions

Stable Bianchi Attractors

Summary and Future outlook

- Bianchi attractors are generalised attractor solutions in gauged supergravity.
- Generalised attractor procedure relies on extremisation of an attractor potential rather than susy.
- non-supersymmetric fixed points can be unstable attractors.
- We studied scalar fluctuations about the attractor value and derived stability conditions by demanding regularity near the horizon.

Results

- Stress energy tensor in gauged supergravity depends on scalar fluctuations even at first order.
- Instability - Ill behaved fluctuations near the horizon will backreact strongly \implies significant deviation from the attractor geometry.
- Consistency condition on scalar fluctuations: critical point is a maxima of the attractor potential.
- Regularity of the fluctuations near the horizon require the near horizon geometry to factorise as $AdS_2 \times M$,

$$ds^2 = L_1^2 \left(-\tilde{r}^2 d\tilde{t}^2 + \frac{d\tilde{r}^2}{\tilde{r}^2} \right) + L_2^2 (\eta_{ij} \omega^i \otimes \omega^j)$$

$M = M_I, M_{II} \dots M_{IX}$ - 3d homogeneous subspaces invariant under the Bianchi type symmetries.

- Purely **radial fluctuations** $k = 0$,


$$\left[\hat{r}^2 \partial_{\hat{r}}^2 + (m+2) \hat{r} \partial_{\hat{r}} - \lambda \right] \delta \phi^x = 0.$$

$$\delta \phi^x = C_1 r^{\left(\sqrt{4\lambda+(1+m)^2}-(1+m)\right)/2} + C_2 r^{\left(-\sqrt{4\lambda+(1+m)^2}-(1+m)\right)/2}$$

- Fluctuations vanishing as $\hat{r} \rightarrow 0$ exist when $\lambda > 0$, $C_2 = 0$.
- However, no bianchi attractors were found for critical points with $\lambda > 0$ - possible model dependent artifacts.

- Based on [stability](#) analysis we conclude that [Bianchi attractors](#) of the form $AdS_2 \times M$ are stable geometries in the deep IR.
- The [factorisation of the near horizon geometry](#) is reminiscent of the situation for [extremal black holes](#).
- It is known that the [near horizon geometry](#) of extremal black holes [preserve supersymmetry](#).
- Reasonable to expect $AdS_2 \times M$ geometries to preserve some fraction of the [supersymmetry](#) (work in progress).

Thank You!



Minimal Surface