

$2 \rightarrow 2$ scattering in supersymmetric matter Chern-Simons theories at large N

Karthik Inbasekar



Young Researchers' Conference, IOPB

29 Sept 2015

Based on

- K.I, S.Jain, S.Mazumdar, S.Minwalla, V.Umesh,
S.Yokoyama: Arxiv [1505.06571](#), accepted in *JHEP*

Related earlier work

- S.Jain, M.Mandlik, S.Minwalla, T.Takimi, S.Wadia,
S.Yokoyama: Arxiv [1404.6373](#), *JHEP* **1504** (2015) 129

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Plan of the talk

Introduction

Preliminaries

Scattering in CS matter theories

Delta function and modified crossing

Our work

Onshell supersymmetry and the S matrix

Onshell supersymmetry

S matrix in onshell superspace

Exact computation of all orders S matrix

Theory

Exact propagator in large N limit

Exact four point correlator in large N limit

S matrix in non-anyonic channels of scattering

S matrix in (singlet) S channel

Unitarity

Pole structure

Summary

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Preliminaries

Scattering in CS
matter theories

Delta function and
modified crossing

Our work

Onshell supersymmetry and the S matrix

Exact computation of all orders S matrix

Unitarity

Pole structure

Summary

CS matter theories: Preliminaries

2 → 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

- Non-Abelian $U(N)$ gauge theories in 2+1 dimensions are rich.

- Yang-Mills + Chern-Simons action

$$\frac{i\kappa}{4\pi} \int \text{Tr} \left(A dA + \frac{2}{3} A^3 \right) - \frac{1}{4g_{YM}^2} \int d^3x \text{Tr} F_{\mu\nu}^2$$

- Describes massive gluons with mass $\propto \kappa g_{YM}^2$.
- Low energies : pure Chern-Simons theory, topological.
- Chern-Simons gauge theory coupled to matter gives rise to interesting dynamics.

Introduction

Preliminaries

Scattering in CS
matter theories

Delta function and
modified crossing

Our work

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

CS matter theories: Preliminaries

- Equations of motion for **abelian theory with scalar matter** of unit charge

$$\kappa \varepsilon^{\mu\nu\rho} F_{\nu\rho} = 2\pi J^\mu$$

- Chern-Simons interaction ties $\frac{1}{\kappa}$ units of **flux** to the charged scalar.
- Aharonov-Bohm** effect: **Exchange** of two unit charge particles result in a **phase** $\frac{\pi}{\kappa}$.
- Chern-Simons** gauge field interacting **with matter** turns them into **anyons** with anyonic phase $\pi\nu = \frac{\pi}{\kappa}$.
- non-abelian** case: for eg **exchange** of $U(N)$ matter quanta R_1 and R_2 gives a **phase operator** $(R_1 \times R_2 = \sum_m R_m)$

$$\nu_m = \frac{T_{R_1} \cdot T_{R_2}}{\kappa} = \frac{C_2(R_1) + C_2(R_2) - C_2(R_m)}{2\kappa}$$

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Preliminaries

Scattering in CS
matter theories
Delta function and
modified crossing
Our work

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

CS matter theories: Preliminaries

- Lots of activity in $U(N)$ Chern-Simons theories with fundamental matter. AdS/CFT, **Vasiliev duality**, limit of ABJ theory.
- **Solvability**: The theory is **solvable in large N** limit.
- Bose-fermi duality:

$$\begin{aligned} &U(N_B) \text{ CS+fundamental boson at Wilson Fisher limit} \\ &\quad \Leftarrow \text{dual} \Rightarrow \\ &U(N_F) \text{ CS+fundamental fermion} \end{aligned}$$

- **Evidence for duality**: Matching of three point functions, spectrum of single trace operators, thermal partition functions, **$2 \rightarrow 2$ S matrices**...

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Preliminaries

Scattering in CS
matter theories

Delta function and
modified crossing
Our work

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Scattering in CS matter theories

- The statement of **duality** is actually a statement of **bosonization of fermions**.
- Bosonic and fermionic S matrices related by **duality** is equivalent to a **bosonization map**.
- Such a mapping is **possible in 2+1 dimensions**: Dirac equation **uniquely** determines the **polarization spinors** as a function of the **momentum**.
- In **large N** limit, only **planar diagrams** contribute. Possible to get **exact results** as a function of 't Hooft coupling λ .
- It has been shown that the **S matrices** for $2 \rightarrow 2$ processes in the **CS+bosonic** theory map to the **CS+fermionic theory** under **duality**.

[Jain, Mandlik, Minwalla, Takimi, Wadia, Yokoyama]

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Preliminaries

Scattering in CS
matter theories

Delta function and
modified crossing

Our work

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Scattering in CS matter theories: Peculiarities

- Scattering results **consistent with duality**.
- In **singlet channel** (particle-Antiparticle) S matrices obtained from **naive crossing symmetry** rules are **inconsistent with unitarity** and have **incorrect non-relativistic limit**.
- **Consistency with unitarity** requires
 - **Delta function** term at forward scattering.
 - **Modified crossing symmetry** rules.
- **Conjecture**: Singlet channel S matrices have the form

$$S = 8\pi\sqrt{s}\cos(\pi\lambda)\delta(\theta) + i\frac{\sin(\pi\lambda)}{\pi\lambda}\mathcal{T}^{S;\text{naive}}(s, \theta)$$

- $\mathcal{T}^{S;\text{naive}}$ is the matrix obtained from **naive analytic continuation** of **particle-particle** scattering.

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Preliminaries

Scattering in CS
matter theories

Delta function and
modified crossing

Our work

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Nature of the conjecture: Delta function and modified crossing rules

$$\mathcal{S} = 8\pi\sqrt{s}\cos(\pi\lambda)\delta(\theta) + i\frac{\sin(\pi\lambda)}{\pi\lambda}\mathcal{T}^{\mathcal{S};\text{naive}}(s, \theta)$$

- The conjectured \mathcal{S} matrix has a **non-analytic $\delta(\theta)$** piece.
- **delta function** is already known to be **necessary to unitarize** non-relativistic **Aharonov-Bohm scattering** [Ruijsenaars; Bak,Jackiw,Pi].
- $\cos(\pi\lambda)$ is due to the **interference of the Aharonov-Bohm phases** of the wave packets.
- modified crossing factor $\frac{\sin(\pi\lambda)}{\pi\lambda}$ is the **expectation value of circular wilson loop** on S^3 in **pure Chern-Simons theory** [Witten].

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Preliminaries

Scattering in CS
matter theories

**Delta function and
modified crossing**

Our work

Onshell
supersymmetry
and the \mathcal{S} matrix

Exact computation
of all orders \mathcal{S}
matrix

Unitarity

Pole structure

Summary

Unitarity and anyonic behavior in Singlet Channel

- **Unitarity** $i(T^\dagger - T) = TT^\dagger$: **non-trivial** only for **singlet channel** in the large N limit.

$$T_{Asym} \sim T_{Sym} \sim T_{Adj} \sim O\left(\frac{1}{N}\right), \quad T_{sing} \sim O(1)$$

- The anyonic phase operator $\nu_m = \frac{C_2(R_1) + C_2(R_2) - C_2(R_m)}{2\kappa}$

$$\nu_{Asym} \sim \nu_{Sym} \sim \nu_{Adj} \sim O\left(\frac{1}{N}\right), \quad \nu_{Sing} \sim O(\lambda)$$

- **symmetric/antisymmetric** channels and **adjoint** channel are effectively **non-anyonic** in large N.
- Particle-Antiparticle scattering in the **singlet channel** is effectively **anyonic** - **usual crossing rules fail unitarity**.
- **Remedy**: **delta function** and **modified crossing rules**.

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Preliminaries

Scattering in CS
matter theories

Delta function and
modified crossing

Our work

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Universality and tests

- delta function and modified crossing rules - appear to be universal
- Tests:
 - Unitarity of the S matrix
 - Bose-Fermi duality
 - Non-relativistic limit gives Aharanov-Bohm
- All the tests have been explicitly verified for
 - $U(N)$ Chern-Simons coupled to fundamental bosons
 - $U(N)$ Chern-Simons coupled to fundamental fermions
 - $\mathcal{N} = 1, 2$ Supersymmetric $U(N)$ Chern-Simons matter theories
- Further investigations ongoing for $\mathcal{N} = 3, 4, 5, 6$ susy CS matter theories [K.I, S.Jain, S.Minwalla, S.Yokoyama]

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Preliminaries

Scattering in CS
matter theories

Delta function and
modified crossing

Our work

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Our work

- Test the **conjecture** in the most general renormalizable supersymmetric $\mathcal{N} = 1$ Chern-Simons matter theory.
- Superspace - **manifest supersymmetry**
- Work in **large N** - only **planar diagrams** .
- Compute **off-shell four point correlator**, take **on-shell limit** and extract the **S matrix**.
- Provide evidence for **duality** and subject the **conjecture** to stringent **unitarity test**.

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Preliminaries

Scattering in CS
matter theories

Delta function and
modified crossing

Our work

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Main results

- Results in perfect agreement with duality.
- Unitarity requires the delta function at forward scattering and crossing symmetry rules modified by exactly the same way as conjectured in [Jain, Mandlik, Minwalla, Takimi, Wadia, Yokoyama]
- Substantial evidence for universality of the conjecture.

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Preliminaries

Scattering in CS
matter theories

Delta function and
modified crossing

Our work

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Bonus results

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

- Results of $\mathcal{N} = 2$ theory obtained at **special value** of quartic scalar coupling.
- **Non-renormalization of pole mass and vertex for $\mathcal{N} = 2$ theory** - good things happen with more susy .
- $\mathcal{N} = 1$ S matrix has **interesting pole structure**, with **vanishing pole mass** on a self-dual codimension one surface in the space of couplings.

Introduction

Preliminaries

Scattering in CS
matter theories

Delta function and
modified crossing

Our work

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Onshell supersymmetry and the S matrix

Introduction

Preliminaries

Scattering in CS matter theories

Delta function and modified crossing

Our work

Onshell supersymmetry and the S matrix

Onshell supersymmetry

S matrix in onshell superspace

Exact computation of all orders S matrix

Theory

Exact propagator in large N limit

Exact four point correlator in large N limit

S matrix in non-anyonic channels of scattering

S matrix in (singlet) S channel

Unitarity

Pole structure

Summary

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Onshell
supersymmetry
S matrix in onshell
superspace

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Supersymmetric scattering

- $2 \rightarrow 2$ scattering amplitude: transition between free incoming and free outgoing onshell particles.
- Initial and final states of Φ_i are effectively subject to free equations of motion

$$(D^2 + m) \Phi = 0$$

- Solution

$$\Phi(x, \theta) = \int \frac{d^2 p}{\sqrt{2p^0}(2\pi)^2} \left[\left(a(\mathbf{p})(1 + m\theta^2) + \theta^\alpha u_\alpha(\mathbf{p}) \alpha(\mathbf{p}) \right) e^{ip \cdot x} + \left(a^{c\dagger}(\mathbf{p})(1 + m\theta^2) + \theta^\alpha v_\alpha(\mathbf{p}) \alpha^{c\dagger}(\mathbf{p}) \right) e^{-ip \cdot x} \right]$$

- action of off-shell supersymmetry operator on onshell superfields $[Q_\alpha, \Phi] = Q_\alpha \Phi = i \left(\frac{\partial}{\partial \theta^\alpha} - \theta^\beta p_{\beta\alpha} \right) \Phi$

$$\begin{aligned} -iQ_\alpha &= u_\alpha(\mathbf{p}_i) (a \partial_\alpha + a^c \partial_{\alpha^c}) + u_\alpha^*(\mathbf{p}_i) (-\alpha \partial_a + \alpha^c \partial_{a^c}) \\ &\quad + v_\alpha(\mathbf{p}_i) (a^\dagger \partial_{\alpha^\dagger} + (a^c)^\dagger \partial_{(\alpha^c)^\dagger}) + v_\alpha^*(\mathbf{p}_i) (\alpha^\dagger \partial_{a^\dagger} + (\alpha^c)^\dagger \partial_{(a^c)^\dagger}) \end{aligned}$$

Onshell superspace

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

- Introduce creation and annihilation operator superfields

$$A_i(\mathbf{p}) = a_i(\mathbf{p}) + \alpha_i(\mathbf{p})\theta_i ,$$

$$A_i^\dagger(\mathbf{p}) = a_i^\dagger(\mathbf{p}) + \theta_i \alpha_i^\dagger(\mathbf{p}) .$$

- Action of supersymmetry operator

$$[Q_\alpha, A_i(\mathbf{p}_i, \theta_i)] = Q_\alpha^1 A_i(\mathbf{p}_i, \theta_i)$$

$$[Q_\alpha, A_i^\dagger(\mathbf{p}_i, \theta_i)] = Q_\alpha^2 A_i^\dagger(\mathbf{p}_i, \theta_i)$$

$$Q_\beta^1 = i \left(-u_\beta(\mathbf{p}) \frac{\overrightarrow{\partial}}{\partial \theta} - v_\beta(\mathbf{p}) \theta \right)$$

$$Q_\beta^2 = i \left(v_\beta(\mathbf{p}) \frac{\overrightarrow{\partial}}{\partial \theta} - u_\beta(\mathbf{p}) \theta \right) .$$

Introduction

Onshell
supersymmetry
and the S matrix

Onshell
supersymmetry
S matrix in onshell
superspace

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Supersymmetry and dual supersymmetry

- Action of **bose-fermi duality**

$$a^D = \alpha, \quad \alpha^D = a \quad m^D = -m$$

- **dual supersymmetry** operator has the form

$$(Q^D)_\beta^1 = i \left(-u_\beta(\mathbf{p}, -m) \frac{\overrightarrow{\partial}}{\partial \theta} - v_\beta(\mathbf{p}, -m) \theta \right),$$

$$(Q^D)_\beta^2 = i \left(v_\beta(\mathbf{p}, -m) \frac{\overrightarrow{\partial}}{\partial \theta} - u_\beta(\mathbf{p}, -m) \theta \right)$$

- using $u(m, p) = -v(-m, p)$, $v(m, p) = -u(-m, p)$
and $\theta \leftrightarrow \frac{\partial}{\partial \theta}$

$$(Q^D)^1 \propto Q^1, \quad (Q^D)^2 \propto Q^2$$

- Quantities invariant under usual **supersymmetry** also invariant under **dual supersymmetry**.

- **Onshell supersymmetry commutes with duality**

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Onshell
supersymmetry
S matrix in onshell
superspace

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

S matrix in onshell superspace

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

- 2 \rightarrow 2 S matrix: $p_1 + p_2 \rightarrow p_3 + p_4$

$$S(\mathbf{p}_1, \theta_1, \mathbf{p}_2, \theta_2, \mathbf{p}_3, \theta_3, \mathbf{p}_4, \theta_4) \sqrt{(2p_1^0)(2p_2^0)(2p_3^0)(2p_4^0)} = \\ \langle 0 | A_4(\mathbf{p}_4, \theta_4) A_3(\mathbf{p}_3, \theta_3) U A_2^\dagger(\mathbf{p}_2, \theta_2) A_1^\dagger(\mathbf{p}_1, \theta_1) | 0 \rangle$$

- Supersymmetric ward identity for the S matrix

$$\left(\vec{Q}_\alpha^1(\mathbf{p}_1, \theta_1) + \vec{Q}_\alpha^1(\mathbf{p}_2, \theta_2) \right. \\ \left. + \vec{Q}_\alpha^2(\mathbf{p}_3, \theta_3) + \vec{Q}_\alpha^2(\mathbf{p}_4, \theta_4) \right) S(\mathbf{p}_1, \theta_1, \mathbf{p}_2, \theta_2, \mathbf{p}_3, \theta_3, \mathbf{p}_4, \theta_4) = 0$$

Introduction

Onshell
supersymmetry
and the S matrix

Onshell
supersymmetry
S matrix in onshell
superspace

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

S matrix in onshell superspace

- S matrix **solution** (in-states: p_1, p_2 , out-states p_3, p_4) is determined **in terms of two functions** \mathcal{S}_B and \mathcal{S}_F of momenta, couplings and mass.

$$\begin{aligned} S(\mathbf{p}_1, \theta_1, \mathbf{p}_2, \theta_2, \mathbf{p}_3, \theta_3, \mathbf{p}_4, \theta_4) = & \mathcal{S}_B + \mathcal{S}_F \theta_1 \theta_2 \theta_3 \theta_4 + \\ & \left(\frac{1}{2} C_{12} \mathcal{S}_B - \frac{1}{2} C_{34}^* \mathcal{S}_F \right) \theta_1 \theta_2 + \left(\frac{1}{2} C_{13} \mathcal{S}_B - \frac{1}{2} C_{24}^* \mathcal{S}_F \right) \theta_1 \theta_3 \\ & + \left(\frac{1}{2} C_{14} \mathcal{S}_B + \frac{1}{2} C_{23}^* \mathcal{S}_F \right) \theta_1 \theta_4 + \left(\frac{1}{2} C_{23} \mathcal{S}_B + \frac{1}{2} C_{14}^* \mathcal{S}_F \right) \theta_2 \theta_3 \\ & + \left(\frac{1}{2} C_{24} \mathcal{S}_B - \frac{1}{2} C_{13}^* \mathcal{S}_F \right) \theta_2 \theta_4 + \left(\frac{1}{2} C_{34} \mathcal{S}_B - \frac{1}{2} C_{12}^* \mathcal{S}_F \right) \theta_3 \theta_4 \end{aligned}$$

- **No θ term:** **four boson** scattering, **four θ term :** **four fermion** scattering.
- All **other processes** (two boson to two fermion etc) **determined completely** in terms of the two independent functions \mathcal{S}_B and \mathcal{S}_F .

S matrix in onshell superspace

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

$$\begin{aligned}\frac{1}{2}C_{12} &= -\frac{1}{4m}v^*(\mathbf{p}_1)v^*(\mathbf{p}_2) & \frac{1}{2}C_{23} &= -\frac{1}{4m}v^*(\mathbf{p}_2)u^*(\mathbf{p}_3) \\ \frac{1}{2}C_{13} &= -\frac{1}{4m}v^*(\mathbf{p}_1)u^*(\mathbf{p}_3) & \frac{1}{2}C_{24} &= -\frac{1}{4m}v^*(\mathbf{p}_2)u^*(\mathbf{p}_4) \\ \frac{1}{2}C_{14} &= -\frac{1}{4m}v^*(\mathbf{p}_1)u^*(\mathbf{p}_4) & \frac{1}{2}C_{34} &= -\frac{1}{4m}u^*(\mathbf{p}_3)u^*(\mathbf{p}_4)\end{aligned}$$

$$\begin{aligned}\frac{1}{2}C_{12}^* &= \frac{1}{4m}v(\mathbf{p}_1)v(\mathbf{p}_2) & \frac{1}{2}C_{23}^* &= \frac{1}{4m}v(\mathbf{p}_2)u(\mathbf{p}_3) \\ \frac{1}{2}C_{13}^* &= \frac{1}{4m}v(\mathbf{p}_1)u(\mathbf{p}_3) & \frac{1}{2}C_{24}^* &= \frac{1}{4m}v(\mathbf{p}_2)u(\mathbf{p}_4) \\ \frac{1}{2}C_{14}^* &= \frac{1}{4m}v(\mathbf{p}_1)u(\mathbf{p}_4) & \frac{1}{2}C_{34}^* &= \frac{1}{4m}u(\mathbf{p}_3)u(\mathbf{p}_4)\end{aligned}$$

Introduction

Onshell
supersymmetry
and the S matrix

Onshell
supersymmetry
S matrix in onshell
superspace

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Exact computation of all orders S matrix

Introduction

Preliminaries

Scattering in CS matter theories

Delta function and modified crossing

Our work

Onshell supersymmetry and the S matrix

Onshell supersymmetry

S matrix in onshell superspace

Exact computation of all orders S matrix

Theory

Exact propagator in large N limit

Exact four point correlator in large N limit

S matrix in non-anyonic channels of scattering

S matrix in (singlet) S channel

Unitarity

Pole structure

Summary

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation of all orders S matrix

Theory

Exact propagator in
large N limit

Exact four point
correlator in large N
limit

S matrix in
non-anyonic channels
of scattering

S matrix in (singlet) S
channel

Unitarity

Pole structure

Summary

Theory

- General renormalizable $\mathcal{N} = 1$ theory coupled to single fundamental matter multiplet Φ

$$\begin{aligned} \mathcal{S}_{\mathcal{N}=1} = - \int d^3x d^2\theta \left[\frac{\kappa}{2\pi} \text{Tr} \left(-\frac{1}{4} D_\alpha \Gamma^\beta D_\beta \Gamma^\alpha \right. \right. \\ \left. \left. - \frac{1}{6} D^\alpha \Gamma^\beta \{ \Gamma_\alpha, \Gamma_\beta \} - \frac{1}{24} \{ \Gamma^\alpha, \Gamma^\beta \} \{ \Gamma_\alpha, \Gamma_\beta \} \right) \right. \\ \left. - \frac{1}{2} (D^\alpha \bar{\Phi} + i \bar{\Phi} \Gamma^\alpha) (D_\alpha \Phi - i \Gamma_\alpha \Phi) \right. \\ \left. + m_0 \bar{\Phi} \Phi + \frac{\pi w}{\kappa} (\bar{\Phi} \Phi)^2 \right] \end{aligned}$$

- Φ : complex superfield, Γ_α : real superfield

$$\Phi = \phi + \theta\psi - \theta^2 F, \quad \bar{\Phi} = \bar{\phi} + \theta\bar{\psi} - \theta^2 \bar{F},$$

$$\Gamma^\alpha = \chi^\alpha - \theta^\alpha B + i\theta^\beta A_\beta^\alpha - \theta^2 (2\lambda^\alpha - i\partial^{\alpha\beta} \chi_\beta).$$

- Integer parameters N, κ , matter coupling constant w , 't Hooft coupling $\lambda = \frac{N}{\kappa}$.

Conjectured Duality for susy matter CS

- Jain, Minwalla, Yokoyama conjectured that $\mathcal{N} = 1, 2$ supersymmetric matter coupled Chern-Simons theories are self dual

$$Theory(\lambda', w', m') \iff Theory(\lambda, w, m)$$

- under the map

$$\lambda' = \lambda - \text{Sgn}(\lambda), \quad w' = \frac{3 - w}{1 + w} \quad m'_0 = \frac{-2m_0}{1 + w}$$

$$N' = |\kappa| - N + 1, \quad \kappa' = -\kappa$$

- with a pole mass

$$m = \frac{2m_0}{2 + (-1 + w)\lambda \text{Sgn}(m)}$$

- $m' = -m$ under duality and $\lambda m(m_0, w) \geq 0$

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Theory

Exact propagator in
large N limit

Exact four point
correlator in large N
limit

S matrix in
non-anyonic channels
of scattering

S matrix in (singlet) S
channel

Unitarity

Pole structure

Summary

Supersymmetric light cone gauge

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

- Supersymmetric generalisation of light cone gauge

$$\Gamma_- = 0 \Rightarrow A_- = A_1 + iA_2 = 0$$

- Gauge self interactions **vanish**

$$S = - \int d^3x d^2\theta \left[- \frac{\kappa}{8\pi} \text{Tr}(\Gamma^- i \partial_- \Gamma^-) - \frac{1}{2} D^\alpha \bar{\Phi} D_\alpha \Phi \right. \\ \left. - \frac{i}{2} \Gamma^- (\bar{\Phi} D_- \Phi - D_- \bar{\Phi} \Phi) \right. \\ \left. + m_0 \bar{\Phi} \Phi + \frac{\pi W}{\kappa} (\bar{\Phi} \Phi)^2 \right]$$

- Susy light cone gauge maintains **manifest supersymmetry**.

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Theory

Exact propagator in
large N limit

Exact four point
correlator in large N
limit

S matrix in
non-anyonic channels
of scattering

S matrix in (singlet) S
channel


Unitarity

Pole structure

Summary

Exact propagator in large N limit

- Integral equation for self-energy

$$\Sigma(p, \theta_1, \theta_2) =$$


$$\begin{aligned} \Sigma(p, \theta_1, \theta_2) = & 2\pi\lambda w \int \frac{d^3r}{(2\pi)^3} \delta^2(\theta_1 - \theta_2) P(r, \theta_1, \theta_2) \\ & - 2\pi\lambda \int \frac{d^3r}{(2\pi)^3} D_-^{\theta_2, -p} D_-^{\theta_1, p} \left(\frac{\delta^2(\theta_1 - \theta_2)}{(p-r)_{--}} P(r, \theta_1, \theta_2) \right) \\ & + 2\pi\lambda \int \frac{d^3r}{(2\pi)^3} \frac{\delta^2(\theta_1 - \theta_2)}{(p-r)_{--}} D_-^{\theta_1, r} D_-^{\theta_2, -r} P(r, \theta_1, \theta_2) \end{aligned}$$

Exact propagator in large N limit

- Solution to **exact propagator** is extremely simple

$$P(p, \theta_1, \theta_2) = \frac{D^2 - m}{p^2 + m^2} \delta^2(\theta_1 - \theta_2)$$

- Same structure as the **bare propagator** with m_0 replaced by m .
- m is the **pole mass**

$$m = \frac{2m_0}{2 + (-1 + w)\lambda \text{Sgn}(m)}$$

is **duality invariant**, agrees with the pole mass computed by **Jain, Minwalla, Yokoyama**

- **Bonus:** In the $\mathcal{N} = 2$ limit ($w = 1$), **no mass renormalization for $\mathcal{N} = 2$ theory** !

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Theory

Exact propagator in
large N limit

Exact four point
correlator in large N
limit

S matrix in
non-anyonic channels
of scattering

S matrix in (singlet) S
channel

Unitarity

Pole structure

Summary

An integral equation for the four point function




Figure: The diagrams in the first line pictorially represents the Schwinger-Dyson equation for offshell four point function. The second line represents the tree level contributions from the gauge superfield interaction and the quartic interactions.

An integral equation for the four point function

- Schematically the integral equations are of the form

$$V(\theta_i, p_i) = V_0(\theta_i, p_i) + \int \frac{d^3 r}{(2\pi)^3} d^2 \theta'_j V_0(\theta_i, \theta'_j, p_i, r) \\ P(\theta'_j, p_i + r) P(\theta'_j, r) V(\theta'_j, \theta_i, p_i)$$

- solved** integral equations **exactly** in large N limit for **arbitrary** values of the **t'Hooft coupling** λ and determined the **offshell four point function** in the kinematic regime $q_{\pm} = 0$.
- Onshell limit** directly gives the **S matrix** for **T (adjoint)**, **U_d (symm)** and **U_e (Asymm)** channels of scattering (q_{μ} is momentum transfer).
- Impossible to extract **S (singlet)** channel S matrix since q_{μ} is center of mass energy (cannot be spacelike).

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Theory

Exact propagator in
large N limit

Exact four point
correlator in large N
limit

S matrix in
non-anyonic channels
of scattering

S matrix in (singlet) S
channel

Unitarity

Pole structure

Summary

S matrix in T , U_d , U_e channels of scattering

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

- S matrix: onshell limit of off-shell four point correlator

$$\mathcal{T}_B = \frac{4i\pi}{\kappa} \epsilon_{\mu\nu\rho} \frac{q^\mu (p-k)^\nu (p+k)^\rho}{(p-k)^2} + J_B(q, \lambda) ,$$

$$\mathcal{T}_F = \frac{4i\pi}{\kappa} \epsilon_{\mu\nu\rho} \frac{q^\mu (p-k)^\nu (p+k)^\rho}{(p-k)^2} + J_F(q, \lambda) ,$$

$$J_B(q, \lambda) = \frac{4\pi q}{\kappa} \frac{N_1 N_2 + M_1}{D_1 D_2} ,$$

$$J_F(q, \lambda) = \frac{4\pi q}{\kappa} \frac{N_1 N_2 + M_2}{D_1 D_2} ,$$

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Theory

Exact propagator in
large N limit

Exact four point
correlator in large N
limit

S matrix in
non-anyonic channels
of scattering

S matrix in (singlet) S
channel

Unitarity

Pole structure

Summary

S matrix in T , U_d , U_e channels of scattering

$$N_1 = \left(\left(\frac{2|m| + iq}{2|m| - iq} \right)^{-\lambda} (w-1)(2m+iq) + (w-1)(2m-iq) \right) ,$$

$$N_2 = \left(\left(\frac{2|m| + iq}{2|m| - iq} \right)^{-\lambda} (q(w+3) + 2im(w-1)) + (q(w+3) - 2im(w-1)) \right) ,$$

$$M_1 = -8mq((w+3)(w-1) - 4w) \left(\frac{2|m| + iq}{2|m| - iq} \right)^{-\lambda} ,$$

$$M_2 = -8mq(1+w)^2 \left(\frac{2|m| + iq}{2|m| - iq} \right)^{-\lambda} ,$$

$$D_1 = \left(i \left(\frac{2|m| + iq}{2|m| - iq} \right)^{-\lambda} (w-1)(2m+iq) - 2im(w-1) + q(w+3) \right) ,$$

$$D_2 = \left(\left(\frac{2|m| + iq}{2|m| - iq} \right)^{-\lambda} (-q(w+3) - 2im(w-1)) + (w-1)(q+2im) \right) .$$

Bonus: S matrix in T , U_d , U_e channels for $\mathcal{N} = 2$ theory

- Remarkable **simplification** in the $\mathcal{N} = 2$ limit ($w=1$)

$$\mathcal{T}_B^{\mathcal{N}=2} = \frac{4i\pi}{\kappa} \epsilon_{\mu\nu\rho} \frac{q^\mu (p-k)^\nu (p+k)^\rho}{(p-k)^2} - \frac{8\pi m}{\kappa} ,$$

$$\mathcal{T}_F^{\mathcal{N}=2} = \frac{4i\pi}{\kappa} \epsilon_{\mu\nu\rho} \frac{q^\mu (p-k)^\nu (p+k)^\rho}{(p-k)^2} + \frac{8\pi m}{\kappa}$$

- All orders S matrix is just tree level - **no loop corrections** - non renormalization.

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Theory

Exact propagator in
large N limit

Exact four point
correlator in large N
limit

S matrix in
non-anyonic channels
of scattering

S matrix in (singlet) S
channel

Unitarity

Pole structure

Summary

Duality invariance of $\mathcal{N} = 1$ and $\mathcal{N} = 2$ S matrices

- Under the duality transformation

$$w' = \frac{3-w}{w+1}, \lambda' = \lambda - \text{sgn}(\lambda), m' = -m, \kappa' = -\kappa$$

$$J_B(q, \kappa', \lambda', w', m') = -J_F(q, \kappa, \lambda, w, m),$$

$$J_F(q, \kappa', \lambda', w', m') = -J_B(q, \kappa, \lambda, w, m).$$

- Duality maps the purely bosonic and purely fermionic S matrices into one another upto overall phase.
- Concrete evidence for duality.
- Supersymmetric ward identity guarantees the duality invariance of all other processes.

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Theory

Exact propagator in
large N limit

Exact four point
correlator in large N
limit

S matrix in
non-anyonic channels
of scattering

S matrix in (singlet) S
channel

Unitarity

Pole structure

Summary

S matrix in (singlet) S channel

- We cannot extract the S channel S matrix directly because of kinematic restriction $q_{\pm} = 0$.
- Usual rules of crossing symmetry in quantum field theory predict particle - anti particle scattering from particle particle scattering or vice-versa
- Naive analytic continuation gives a non-unitary S matrix in the S channel as observed in earlier work.
- Any analytic continuation cannot produce the non-analytic delta function piece required for unitarization.
- Remedy: Modify crossing symmetry rules as conjectured in [Jain, Mandlik, Minwalla, Takimi, Wadia, Yokoyama]

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Theory

Exact propagator in
large N limit

Exact four point
correlator in large N
limit

S matrix in
non-anyonic channels
of scattering

S matrix in (singlet) S
channel

Unitarity

Pole structure

Summary

Conjectured S matrix in S channel $\mathcal{N} = 1$ theory

- Conjectured S matrix for the $\mathcal{N} = 1$ theory in center of mass frame

$$\mathcal{S}_B^S(s, \theta) = 8\pi\sqrt{s}\cos(\pi\lambda)\delta(\theta) + i\frac{\sin(\pi\lambda)}{\pi\lambda} \left(4\pi i\lambda\sqrt{s}\cot(\theta/2) + J_B(\sqrt{s}, \lambda) \right) ,$$

$$\mathcal{S}_F^S(s, \theta) = 8\pi\sqrt{s}\cos(\pi\lambda)\delta(\theta) + i\frac{\sin(\pi\lambda)}{\pi\lambda} \left(4\pi i\lambda\sqrt{s}\cot(\theta/2) + J_F(\sqrt{s}, \lambda) \right) .$$

$$J_B(\sqrt{s}, \lambda) = -4\pi i\lambda\sqrt{s}\frac{N_1N_2 + M_1}{D_1D_2} ,$$

$$J_F(\sqrt{s}, \lambda) = -4\pi i\lambda\sqrt{s}\frac{N_1N_2 + M_2}{D_1D_2}$$

Conjectured S matrix in S channel $\mathcal{N} = 1$ theory

$$N_1 = \left((w-1)(2m + \sqrt{s}) + (w-1)(2m - \sqrt{s})e^{i\pi\lambda} \left(\frac{\sqrt{s} + 2|m|}{\sqrt{s} - 2|m|} \right)^\lambda \right),$$

$$N_2 = \left((-i\sqrt{s}(w+3) + 2im(w-1)) + (-i\sqrt{s}(w+3) - 2im(w-1))e^{i\pi\lambda} \left(\frac{\sqrt{s} + 2|m|}{\sqrt{s} - 2|m|} \right)^\lambda \right),$$

$$M_1 = 8mi\sqrt{s}((w+3)(w-1) - 4w)e^{i\pi\lambda} \left(\frac{\sqrt{s} + 2|m|}{\sqrt{s} - 2|m|} \right)^\lambda,$$

$$M_2 = 8mi\sqrt{s}(1+w)^2e^{i\pi\lambda} \left(\frac{\sqrt{s} + 2|m|}{\sqrt{s} - 2|m|} \right)^\lambda,$$

$$D_1 = \left(i(w-1)(2m + \sqrt{s}) - (2im(w-1) + i\sqrt{s}(w+3))e^{i\pi\lambda} \left(\frac{\sqrt{s} + 2|m|}{\sqrt{s} - 2|m|} \right)^\lambda \right),$$

$$D_2 = \left((\sqrt{s}(w+3) - 2im(w-1)) + (w-1)(-i\sqrt{s} + 2im)e^{i\pi\lambda} \left(\frac{\sqrt{s} + 2|m|}{\sqrt{s} - 2|m|} \right)^\lambda \right)$$

Straightforward non-relativistic limit of the $\mathcal{N} = 1$ S matrix

- Non-rel limit: $\sqrt{s} \rightarrow 2m$ with all other parameters held fixed.

$$\mathcal{T}_B^S(s, \theta) = -8\pi i \sqrt{s} (\cos(\pi\lambda) - 1) \delta(\theta) + 4\sqrt{s} \sin(\pi\lambda) (i \cot(\theta/2) - 1) ,$$

$$\mathcal{T}_F^S(s, \theta) = -8\pi i \sqrt{s} (\cos(\pi\lambda) - 1) \delta(\theta) + 4\sqrt{s} \sin(\pi\lambda) (i \cot(\theta/2) + 1) .$$

- conjectured S channel S matrix has simple non-relativistic limit leading to known Aharonov-Bohm result.
- Surprisingly this result is also same as the $\mathcal{N} = 2$ S channel S matrix.
- Presumably supersymmetry enhancement in non-relativistic limit.

Unitarity

Introduction

Preliminaries

Scattering in CS matter theories

Delta function and modified crossing

Our work

Onshell supersymmetry and the S matrix

Onshell supersymmetry

S matrix in onshell superspace

Exact computation of all orders S matrix

Theory

Exact propagator in large N limit

Exact four point correlator in large N limit

S matrix in non-anyonic channels of scattering

S matrix in (singlet) S channel

Unitarity

Pole structure

Summary

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Unitarity equation

- Define on-shell superfield S^\dagger as

$$S^\dagger(\mathbf{p}_1, \theta_1, \mathbf{p}_2, \theta_2, \mathbf{p}_3, \theta_3, \mathbf{p}_4, \theta_4) = S^*(\mathbf{p}_3, \theta_3, \mathbf{p}_4, \theta_4, \mathbf{p}_1, \theta_1, \mathbf{p}_2, \theta_2)$$

- Supersymmetric ward identity for S^\dagger implies S^\dagger is supersymmetric if and only if S is supersymmetric.
- The supersymmetric unitarity equation is

$$(S \star S^\dagger - I) = 0$$

- Recall that the superfield expansion for S is completely specified by \mathcal{S}_B and \mathcal{S}_F .
- Sufficient to check the LHS for no θ and four θ terms.
- Supersymmetric ward identity guarantees the rest of the terms will obey the unitarity equations.

Unitarity equations for T , U_d and U_e channels

- Writing $\mathcal{S}_B = I + i\mathcal{T}_B$, $\mathcal{S}_F = I + i\mathcal{T}_f$
- The S matrices in the T , U_d and U_e channels are all $O(1/N)$ - unitarity equation is linear

$$\mathcal{T}_B(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4) = \mathcal{T}_B^*(\mathbf{p}_3, \mathbf{p}_4, \mathbf{p}_1, \mathbf{p}_2) ,$$

$$\mathcal{T}_F(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4) = \mathcal{T}_F^*(\mathbf{p}_3, \mathbf{p}_4, \mathbf{p}_1, \mathbf{p}_2)$$

- Linearity: No branch cuts in the physical domain of scattering in these channels.
- Explicitly checked that unitarity conditions are obeyed using

$$J_B(q, \lambda) = J_B^*(-q, \lambda) , \quad J_F(q, \lambda) = J_F^*(-q, \lambda)$$

- The S matrix in the S channel is $O(1)$ - the unitarity conditions are non-linear

Unitarity equations in the S channel

- Consider the general structure ($T(\theta) = i \cot(\theta/2)$.)

$$\mathcal{T}_B^S = H_B T(\theta) + W_B - iW_2 \delta(\theta) , \quad \mathcal{T}_F^S = H_F T(\theta) + W_F - iW_2 \delta(\theta) ,$$

- first unitarity equation

$$H_B - H_B^* = \frac{1}{8\pi\sqrt{s}}(W_2 H_B^* - H_B W_2^*) ,$$

$$W_2 + W_2^* = -\frac{1}{8\pi\sqrt{s}}(W_2 W_2^* + 4\pi^2 H_B H_B^*) ,$$

$$W_B - W_B^* = \frac{1}{8\pi\sqrt{s}}(W_2 W_B^* - W_2^* W_B) - \frac{i}{4\sqrt{s}}(H_B H_B^* - W_B W_B^*) - \frac{iY}{4\sqrt{s}}(W_B - W_F)(W_B^* - W_F^*)$$

- Second unitarity equation

$$H_F - H_F^* = \frac{1}{8\pi\sqrt{s}}(W_2 H_F^* - H_F W_2^*) ,$$

$$W_2 + W_2^* = -\frac{1}{8\pi\sqrt{s}}(W_2 W_2^* + 4\pi^2 H_F H_F^*) ,$$

$$W_F - W_F^* = \frac{1}{8\pi\sqrt{s}}(W_2 W_F^* - W_2^* W_F) - \frac{i}{4\sqrt{s}}(H_F H_F^* - W_F W_F^*) - \frac{iY}{4\sqrt{s}}(W_B - W_F)(W_B^* - W_F^*)$$

Unitarity equation in the S channel

- Unitarity equations are verified using

$$H_B = H_F = 4\sqrt{s} \sin(\pi\lambda), \quad W_2 = 8\pi\sqrt{s}(\cos(\pi\lambda)-1), \quad T(\theta) = i \cot(\theta/2)$$

$$W_B = J_B(\sqrt{s}, \lambda) \frac{\sin(\pi\lambda)}{\pi\lambda},$$

$$W_F = J_F(\sqrt{s}, \lambda) \frac{\sin(\pi\lambda)}{\pi\lambda}.$$

- **Algebraic-miracle**: Non-linear unitarity equations obeyed by very complicated functions.
- unitarity is an extremely sensitive test ¹.
- Important to note that the **crossing symmetry rules** have to be modified exactly as conjectured in [Jain, Mandlik, Minwalla, Takimi, Wadia, Yokoyama], else the unitarity test fails.

¹Tagline: **one sign is doom**

Unitarity in the S channel - $\mathcal{N} = 2$ case

- The $\mathcal{N} = 2$ T matrix is tree level exact in T, U channels.
- Naive crossing symmetry would imply the same for S channel, unitarity equation $i(T^\dagger - T) = TT^\dagger$ would never be obeyed (LHS would be zero).
- modified crossing rules resolve this puzzle:

$$\mathcal{T}_B^{S;\mathcal{N}=2}(s, \theta) = -8\pi i \sqrt{s} (\cos(\pi\lambda) - 1) \delta(\theta) + \sin(\pi\lambda) (4i\sqrt{s} \cot(\theta/2) - 8m) ,$$

$$\mathcal{T}_F^{S;\mathcal{N}=2}(s, \theta) = -8\pi i \sqrt{s} (\cos(\pi\lambda) - 1) \delta(\theta) + \sin(\pi\lambda) (4i\sqrt{s} \cot(\theta/2) + 8m) .$$

- Non-analytic piece makes $\mathcal{T}_B, \mathcal{T}_F$ not Hermitian, both LHS and RHS are non-zero and non-linear unitarity equation is obeyed.

Pole structure of the S matrix

Introduction

Preliminaries

Scattering in CS matter theories

Delta function and modified crossing

Our work

Onshell supersymmetry and the S matrix

Onshell supersymmetry

S matrix in onshell superspace

Exact computation of all orders S matrix

Theory

Exact propagator in large N limit

Exact four point correlator in large N limit

S matrix in non-anyonic channels of scattering

S matrix in (singlet) S channel

Unitarity

Pole structure

Summary

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Pole structure of the singlet channel S matrix

- Both bosonic and fermionic S matrices have a pole at threshold ($s = 4m^2$) for $w \leq -1$. For $w \leq -1 + \epsilon$ the pole is close to threshold.

- As w is decreased further and as it hits a critical value $w = w_c$ the pole becomes massless!

$$w = w_c(\lambda) = 1 - \frac{2}{|\lambda|}$$

- As w is further decreased and as $w \rightarrow -\infty$ the pole approaches threshold once again.
- To summarize, a one parameter tuning of the superpotential interaction parameter w - sufficient to produce massless bound states in a massive theory.
- w can be scaled to w_c - possible decoupled QFT description of light states.

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Summary

Introduction

Preliminaries

Scattering in CS matter theories

Delta function and modified crossing

Our work

Onshell supersymmetry and the S matrix

Onshell supersymmetry

S matrix in onshell superspace

Exact computation of all orders S matrix

Theory

Exact propagator in large N limit

Exact four point correlator in large N limit

S matrix in non-anyonic channels of scattering

S matrix in (singlet) S channel

Unitarity

Pole structure

Summary

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Summary

- Computations and conjectures for the all orders $w \rightarrow 2$ S matrix in the most general renormalizable $\mathcal{N} = 1$ Chern-Simons matter theory with a single fundamental matter multiplet.
- Used supersymmetric ward identity to derive conditions and constraints on off-shell correlators, on-shell S matrices and derive unitarity conditions.
- Computed exact offshell four point correlators in the large N limit in kinematic regime $q_{\pm} = 0$.
- Obtained S matrices by taking onshell limit of offshell four point correlator.
- Conjectured S matrix in the singlet channel.

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Summary

- Results are consistent with duality.
- Results are consistent with unitarity if and only if we assume that the usual results of crossing symmetry are modified in precisely the manner proposed in [Jain, Mandlik, Minwalla, Takimi, Wadia, Yokoyama].
- Non-relativistic limit of the S matrix reproduces the known Aharonov-Bohm result.
- The S channel S matrix has an interesting analytic structure. In a certain range of superpotential parameters the S matrix has a bound state pole.
- A one parameter tuning of superpotential parameters can be used to set the pole mass to zero.

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Open questions

- $\mathcal{N} = 2$ S matrices are tree level exact in non-anyonic channels and depend on λ very simple way in the anyonic channel - can it be reproduced from general principles and $\mathcal{N} = 2$ supersymmetry?
- Generalisation to higher supersymmetry - mass deformed $\mathcal{N} = 3, 4, 5$, and mass deformed $\mathcal{N} = 6$ ABJ theory. [K.I, S.Jain, S.Minwalla, S. Yokoyama]
- Four point correlator: useful in computation of 2,3,4 point functions of gauge invariant currents - explicit computation in $\mathcal{N} = 2$ theory?, possible $\mathcal{N} = 2$ generalisation of Maldacena-Zhiboedov solutions. [K.I, S.Jain, P.Nayak]

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Open questions

- Rigorous proof of delta function and modified crossing rules, generalisation to finite N and κ .
- From perturbative pov modified crossing rules could be related to IR divergences.
- IR divergences can be summed up and exponentiated [Grammer,Yennie; Bern,Dickson,Smirnov]
- Modified crossing factor $\frac{\sin(\pi\lambda)}{\pi\lambda}$ is identical to the expectation value of circular wilson loop in pure Chern-Simons theory on S^3 .
- To explore: Path integral derivation of Witten's result, crossing and fusion rules in RCFT's.

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Thank You!

Level rank duality in CS matter theory

- $U(N_B)$ Chern-Simons theory coupled to fundamental boson [Giombi, Minwalla, Prakash, Trivedi, Wadia]

$$S = \int d^3x \left(i\epsilon^{\mu\nu\rho} \frac{\kappa_B}{4\pi} \text{Tr}(A_\mu \partial_\nu A_\rho - \frac{2i}{3} A_\mu A_\nu A_\rho) \right. \\ \left. D_\mu \bar{\phi} D^\mu \phi + \sigma \bar{\phi} \phi + N_B \frac{m_B^2}{b_4} \sigma - N_B \frac{\sigma^2}{2b_4} \right)$$

- Wilson-Fisher limit

$$b_4 \rightarrow \infty, \quad m_B \rightarrow \infty, \quad 4\pi \frac{m_B^2}{b_4} = \text{fixed}$$

- $U(N_F)$ Chern-Simons theory coupled to fundamental fermion

$$S = \int d^3x \left(i\epsilon^{\mu\nu\rho} \frac{\kappa_F}{4\pi} \text{Tr}(A_\mu \partial_\nu A_\rho - \frac{2i}{3} A_\mu A_\nu A_\rho) \right. \\ \left. + \bar{\psi} \gamma^\mu D_\mu \psi + m_F \bar{\psi} \psi \right)$$

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Level rank duality in CS matter theory

- Statement of duality

$U(N_B)$ CS+fundamental boson at Wilson Fisher limit

\Leftarrow dual \Rightarrow

$U(N_F)$ CS+fundamental fermion

- under the duality map

$$\kappa_F = -\kappa_B$$

$$N_F = |\kappa_B| - N_B$$

$$\lambda_B = \lambda_F - \text{sgn}(\lambda_F)$$

$$m_F = -m_B^{\text{cri}} \lambda_B$$

- with condition

$$\lambda_F m_F > 0$$

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Evidence for duality

- Spectrum of single trace operators and three point functions on both sides match.
[Giombi, Minwalla, Prakash, Trivedi, Wadia] ,
[Aharony, Gur-Ari, Yacoby], [Maldacena, Zhiboedov]
- Thermal partition functions on both sides match.
[Jain, Trivedi, Wadia, Yokoyama] ,
[Aharony, Giombi, Gur-Ari, Maldacena, Yacoby], [Takimi]
- Duality follows from a deformation of the known Giveon-Kutasov duality in supersymmetric theory.
[Jain, Minwalla, Yokoyama]
- $2 \rightarrow 2$ S matrices for purely bosonic and purely fermionic theories have been computed to all orders in t'Hooft coupling and map to one another under duality
[Jain, Mandlik, Minwalla, Takimi, Wadia, Yokoyama]

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix


Unitarity

Pole structure

Summary

Need for a conjecture

- $P_i(p_1) + A^j(p_2) \rightarrow P_m(p_3) + A^n(p_4)$



- Work in light-cone gauge in the frame $q_{\pm} = 0$.
- Adjoint** channel (from top) - $q_{\pm} = 0$ is a frame choice, full answer can be obtained by covariantizing.

$$p_1 = p + q, \quad p_2 = -k - q, \quad p_3 = p, \quad p_4 = -k$$

- Singlet** channel (from left), **exchange momentum cannot be spacelike!**

$$p_1 = p + q, \quad p_2 = -p, \quad p_3 = k + q, \quad p_4 = -k$$

- Cannot compute singlet channel directly.
- Using **usual crossing symmetry** gives a **non-unitary S matrix** for **singlet** channel.

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Bare Propagators

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

- The bare scalar superfield propagator:

$$\langle \bar{\Phi}(\theta_1, p) \Phi(\theta_2, -p') \rangle = \frac{D_{\theta_1, p}^2 - m_0^2}{p^2 + m_0^2} \delta^2(\theta_1 - \theta_2) (2\pi)^3 \delta^3(p - p')$$

- The gauge superfield propagator:

$$\langle \Gamma^-(\theta_1, p) \Gamma^-(\theta_2, -p') \rangle = -\frac{8\pi}{\kappa} \frac{\delta^2(\theta_1 - \theta_2)}{p_{--}} (2\pi)^3 \delta^3(p - p')$$

where $p_{--} = -(p_1 + ip_2) = -p_-$.

- Gauge field component propagators have same form as non-susy light cone gauge

$$\langle A_+(p) A_3(-p') \rangle = \frac{4\pi i}{\kappa} \frac{1}{p_-} (2\pi)^3 \delta^3(p - p')$$

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Susy constraints on two-point correlator

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

- Supersymmetric ward identity for two point correlator

$$(Q_{\theta_1,p} + Q_{\theta_2,-p})\langle\bar{\Phi}(\theta_1,p)\Phi(\theta_2,-p)\rangle = 0$$

- Exact propagator solves the ward identity

$$\langle\bar{\Phi}(p,\theta_1)\Phi(-p',\theta_2)\rangle = (2\pi)^3\delta^3(p-p')P(\theta_1,\theta_2,p)$$

$$P(\theta_1,\theta_2,p) = (C_1(p^\mu)D_{\theta_1,p}^2 + C_2(p^\mu))\delta^2(\theta_1-\theta_2)$$

- eg for bare propagator

$$C_1 = \frac{1}{p^2 + m_0^2}, \quad C_2 = \frac{m_0}{p^2 + m_0^2}$$

Introduction

Onshell
supersymmetry
and the S matrix


Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Susy constraints on four-point function



- Supersymmetric ward identity for four point function

$$(Q_{\theta_1, p+q} + Q_{\theta_2, -p} + Q_{\theta_3, -k-q} + Q_{\theta_4, k})V(\theta_1, \theta_2, \theta_3, \theta_4, p, k, q) = 0$$

$$\begin{aligned} \langle \bar{\Phi}((p+q + \frac{l}{4}), \theta_1) \Phi(-p + \frac{l}{4}, \theta_2) \Phi(-(k+q) + \frac{l}{4}, \theta_3) \bar{\Phi}(k + \frac{l}{4}, \theta_4) \rangle \\ = (2\pi)^3 \delta(l) V(\theta_1, \theta_2, \theta_3, \theta_4, p, k, q) \end{aligned}$$

Susy constraints on four-point function

- Solution of the ward identity

$$V = \exp \left(\frac{1}{4} X \cdot (p \cdot X_{12} + q \cdot X_{13} + k \cdot X_{43}) \right) F(X_{12}, X_{13}, X_{43}, p, q, k)$$

$$X = \sum_{i=1}^4 \theta_i, \quad X_{ij} = \theta_i - \theta_j,$$

- F is a shift invariant function $\theta_i \rightarrow \theta_i + \gamma$.
- V may be taken to be invariant under the \mathbb{Z}_2 symmetry

$$p \rightarrow k + q, k \rightarrow p + q, q \rightarrow -q, \\ \theta_1 \rightarrow \theta_4, \theta_2 \rightarrow \theta_3, \theta_3 \rightarrow \theta_2, \theta_4 \rightarrow \theta_1$$

An integral equation for the four point function

- Most general form of F can be parameterized in terms of 32 bosonic functions of p, k and q .
- leads to 32 coupled integral equations - tedious.
- In the kinematic regime $q_{\pm} = 0$ the ansatz

$$V = \exp\left(\frac{1}{4}X.(p.X_{12} + q.X_{13} + k.X_{43})\right) F(X_{12}, X_{13}, X_{43}, p, q, k)$$
$$F = X_{12}^+ X_{43}^+ \left(A(p, k, q) X_{12}^- X_{43}^- X_{13}^+ X_{13}^- + B(p, k, q) X_{12}^- X_{43}^- \right. \\ \left. + C(p, k, q) X_{12}^- X_{13}^+ + D(p, k, q) X_{13}^+ X_{43}^- \right)$$

is closed under the multiplication rule induced by the RHS of the integral equation.

Product of S matrices

- General multiplication rule for two S matrices

$$S_1 \star S_2 \equiv \int d\Gamma S_1(\mathbf{p}_1, \theta_1, \mathbf{p}_2, \theta_2, \mathbf{k}_3, \phi_1, \mathbf{k}_4, \phi_2) \\ \exp(\phi_1\phi_3 + \phi_2\phi_4) 2k_1^0 (2\pi)^2 \delta^{(2)}(\mathbf{k}_3 - \mathbf{k}_1) 2k_2^0 (2\pi)^2 \delta^{(2)}(\mathbf{k}_4 - \mathbf{k}_2) \\ S_2(\mathbf{k}_1, \phi_3, \mathbf{k}_2, \phi_4, \mathbf{p}_3, \theta_3, \mathbf{p}_4, \theta_4)$$

$$d\Gamma = \frac{d^2 k_3}{2k_3^0 (2\pi)^2} \frac{d^2 k_4}{2k_4^0 (2\pi)^2} \frac{d^2 k_1}{2k_1^0 (2\pi)^2} \frac{d^2 k_2}{2k_2^0 (2\pi)^2} d\phi_1 d\phi_3 d\phi_2 d\phi_4$$

- supersymmetry invariant Identity operator

$$I(\mathbf{p}_1, \theta_1, \mathbf{p}_2, \theta_2, \mathbf{p}_3, \theta_3, \mathbf{p}_4, \theta_4) = \exp(\theta_1\theta_3 + \theta_2\theta_4) I(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4)$$

$$I(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4) = 2p_3^0 (2\pi)^2 \delta^{(2)}(\mathbf{p}_1 - \mathbf{p}_3) 2p_4^0 (2\pi)^2 \delta^{(2)}(\mathbf{p}_2 - \mathbf{p}_4)$$

- Multiplication rule with Identity operator

$$S \star I = I \star S = S$$

- More generally product of two supersymmetric S matrices is supersymmetric.

Unitarity equation in center of mass frame

- Writing $\mathcal{S}_B = I + i\mathcal{T}_B$, $\mathcal{S}_F = I + i\mathcal{T}_f$

- No theta term:

$$\frac{1}{8\pi\sqrt{s}} \int d\theta \left(-Y(s)(\mathcal{T}_B(s, \theta) + 4Y(s)\mathcal{T}_f(s, \theta))(\mathcal{T}_B^*(s, -(\alpha - \theta)) + 4Y(s)\mathcal{T}_f^*(s, -(\alpha - \theta))) \right. \\ \left. + \mathcal{T}_B(s, \theta)\mathcal{T}_B^*(s, -(\alpha - \theta)) \right) = i(\mathcal{T}_B^*(s, -\alpha) - \mathcal{T}_B(s, \alpha))$$

- Four theta term:

$$\frac{1}{8\pi\sqrt{s}} \int d\theta \left(Y(s)(\mathcal{T}_B(s, \theta) + 4Y(s)\mathcal{T}_f(s, \theta))(\mathcal{T}_B^*(s, -(\alpha - \theta)) + 4Y(s)\mathcal{T}_f^*(s, -(\alpha - \theta))) \right. \\ \left. - 16Y(s)^2\mathcal{T}_f(s, \theta)\mathcal{T}_f^*(s, -(\alpha - \theta)) \right) = i4Y(s)(-\mathcal{T}_f(s, \alpha) + \mathcal{T}_f^*(s, -\alpha))$$

$$Y(s) = \frac{-s + 4m^2}{16m^2}$$

Unitarity equations in the S channel

- The S matrix in the S channel is $O(1)$ - the unitarity conditions are non-linear

$$\frac{1}{8\pi\sqrt{s}} \int d\theta \left(-Y(s)(\mathcal{T}_B^S(s, \theta) - \mathcal{T}_F^S(s, \theta))(\mathcal{T}_B^{S*}(s, -(\alpha - \theta)) - \mathcal{T}_F^{S*}(s, -(\alpha - \theta))) \right. \\ \left. + \mathcal{T}_B^S(s, \theta)\mathcal{T}_B^{S*}(s, -(\alpha - \theta)) \right) = i(\mathcal{T}_B^{S*}(s, -\alpha) - \mathcal{T}_B^S(s, \alpha))$$

$$\frac{1}{8\pi\sqrt{s}} \int d\theta \left(Y(s)(\mathcal{T}_B^S(s, \theta) - \mathcal{T}_F^S(s, \theta))(\mathcal{T}_B^{S*}(s, -(\alpha - \theta)) - \mathcal{T}_F^{S*}(s, -(\alpha - \theta))) \right. \\ \left. - \mathcal{T}_F^S(s, \theta)\mathcal{T}_F^{S*}(s, -(\alpha - \theta)) \right) = i(\mathcal{T}_F^S(s, \alpha) - \mathcal{T}_F^{S*}(s, -\alpha))$$

- Under duality $\mathcal{T}_B \rightarrow \mathcal{T}_F$ and vice versa; both the equations map to one another.
- Unitarity conditions are compatible with duality.

Unitarity equations in the S channel

- In the **center of mass frame**, the supersymmetric unitarity equations are

$$\frac{1}{8\pi\sqrt{s}} \int d\theta \left(-Y(s)(\mathcal{T}_B^S(s, \theta) - \mathcal{T}_F^S(s, \theta))(\mathcal{T}_B^{S*}(s, -(\alpha - \theta)) - \mathcal{T}_F^{S*}(s, -(\alpha - \theta))) \right. \\ \left. + \mathcal{T}_B^S(s, \theta)\mathcal{T}_B^{S*}(s, -(\alpha - \theta)) \right) = i(\mathcal{T}_B^{S*}(s, -\alpha) - \mathcal{T}_B^S(s, \alpha))$$

$$\frac{1}{8\pi\sqrt{s}} \int d\theta \left(Y(s)(\mathcal{T}_B^S(s, \theta) - \mathcal{T}_F^S(s, \theta))(\mathcal{T}_B^{S*}(s, -(\alpha - \theta)) - \mathcal{T}_F^{S*}(s, -(\alpha - \theta))) \right. \\ \left. - \mathcal{T}_F^S(s, \theta)\mathcal{T}_F^{S*}(s, -(\alpha - \theta)) \right) = i(\mathcal{T}_F^S(s, \alpha) - \mathcal{T}_F^{S*}(s, -\alpha))$$

- Under **duality** $\mathcal{T}_B \rightarrow \mathcal{T}_F$ and vice versa; both the equations map to one another.
- **Unitarity conditions are compatible with duality.**

Poles of the S matrix

2 \rightarrow 2 scattering
in supersymmetric
matter
Chern-Simons
theories at large N

- Both bosonic and fermionic S matrix have a **pole at threshold for $w \leq -1$** . Near $w = -1 - \delta w, y = 1 - \delta y$ the S matrix has the pole structure ($y = \sqrt{s}/2m$)

$$\mathcal{T}_B \sim \frac{\left(\frac{\delta y}{2}\right)^{|\lambda|}}{\delta w - 2\left(\frac{\delta y}{2}\right)^{|\lambda|}}, \quad \mathcal{T}_F \sim \frac{\left(\frac{\delta y}{2}\right)^{1+|\lambda|}}{\delta w - 2\left(\frac{\delta y}{2}\right)^{|\lambda|}}$$

- As w is decreased further and as it hits a **critical value $w = w_c$**

$$w = w_c(\lambda) = 1 - \frac{2}{|\lambda|}$$

- the **pole becomes massless!**. Near $w = w_c - \delta w$ and $y = \delta y$ the poles **approach zero mass quadratically**

$$\mathcal{T}_B \sim \mathcal{T}_F - \frac{64|m|\sin(\pi\lambda)(-1 + |\lambda|)}{|\lambda|(\delta w^2\lambda^2 - 4\delta y^2(1 - |\lambda|)^2)}$$

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary

Poles of the S matrix

- As w is further decreased and as $w \rightarrow -\infty$ the pole approaches threshold once again. Near $w = -\frac{1}{\delta w}$, $y = 1 - \delta y$ the S matrix has the pole structure

$$\mathcal{T}_B \sim \frac{\left(\frac{\delta y}{2}\right)^{2-|\lambda|}}{\delta w - \frac{1}{2} \left(\frac{\delta y}{2}\right)^{1-|\lambda|}}, \quad \mathcal{T}_F \sim \frac{\left(\frac{\delta y}{2}\right)^{1-|\lambda|}}{\delta w - \frac{1}{2} \left(\frac{\delta y}{2}\right)^{1-|\lambda|}}$$

- To summarize, a one parameter tuning of the superpotential interaction parameter w - sufficient to produce massless bound states in our massive theory.
- w can be scaled to w_c - possible decoupled QFT description of light states.
- Is this a $\mathcal{N} = 1$ Wilson-Fischer theory made of single real superfield?

Analytic structure of S channel S matrix

- The S matrix in the singlet channel has an **interesting analytic structure**.
- As a function of s (at fixed t), there is the expected two particle branch cut starting at $s = 4m^2$.
- For smaller but **positive values of s** there exist **poles in the S matrix** for a range of coupling parameters.
- These **poles represent bound states** that exist at large but finite N .
- At some **critical value** of the scalar coupling $w = w_c(\lambda)$ the **pole becomes massless!**
- To summarize, **a one parameter tuning** of the superpotential interaction parameter w - **sufficient to produce massless bound states in a massive theory.**

$2 \rightarrow 2$ scattering
in supersymmetric
matter
Chern-Simons
theories at large N

Introduction

Onshell
supersymmetry
and the S matrix

Exact computation
of all orders S
matrix

Unitarity

Pole structure

Summary