

EE5609 Matrix Theory

Kranthi Kumar P

Download the latex-file codes from

https://github.com/kranthiakssy/AI20RESCH14002_PhD_IITH/tree/master/EE5609_Matrix_Theory/Assignment-14

1 PROBLEM

(ugcdec2014, 29) :

The determinant of $n \times n$ permutation matrix

$$\begin{pmatrix} & & & & 1 \\ & & & 1 & \\ & & \cdot & & \\ & & \cdot & & \\ & & \cdot & & \\ 1 & & & & \end{pmatrix}$$

- 1) $(-1)^n$
- 2) $(-1)^{\lfloor \frac{n}{2} \rfloor}$
- 3) -1
- 4) 1

2 SOLUTION

Given	$n \times n$ permutation matrix $\begin{pmatrix} & & & & 1 \\ & & & 1 & \\ & & \cdot & & \\ & & \cdot & & \\ & & \cdot & & \\ 1 & & & & \end{pmatrix}$
Solution	<p>The given $n \times n$ permutation matrix can be converted into identity matrix of $n \times n$ dimension by doing row exchange operations.</p> <p>from the row exchange property of determinants the determinant will be multiplied by -1 for every row exchange.</p> <p>the given $n \times n$ matrix requires $\lfloor \frac{n}{2} \rfloor$</p>

	<p>row exchanges to become identity matrix.</p> <p>Hence the determinant of given permutation matrix is</p> $(-1)^{\lfloor \frac{n}{2} \rfloor} \begin{vmatrix} 1 & & & & \\ & 1 & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & 1 \\ & & & & & 1 \end{vmatrix}$ <p>we know that the determinant of identity matrix, $\det(\mathbf{I}) = 1$ \therefore the determinant of given $n \times n$ permutation matrix $= (-1)^{\lfloor \frac{n}{2} \rfloor}$</p>
Conclusion	Option-2 is the right solution

TABLE 1: Solution

3 EXAMPLE

Example-1	<p>Let \mathbf{A} is 5×5 permutation matrix, then</p> $\det(\mathbf{A}) = \begin{vmatrix} & & & & 1 \\ & & & 1 & \\ & & 1 & & \\ & 1 & & & \\ 1 & & & & \end{vmatrix}$ $\xrightarrow{R_1 \leftrightarrow R_5} (-1) \begin{vmatrix} 1 & & & & \\ & & & 1 & \\ & & 1 & & \\ & 1 & & & \\ & & & & 1 \end{vmatrix}$ $\xrightarrow{R_2 \leftrightarrow R_4} (-1)(-1) \begin{vmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{vmatrix}$ $= 1$ <p>substituting $n = 5$ in the solution $(-1)^{\lfloor \frac{5}{2} \rfloor} = 1$</p>
Example-2	<p>Let \mathbf{A} is 6×6 permutation matrix, then</p> $\det(\mathbf{A}) = \begin{vmatrix} & & & & & 1 \\ & & & & 1 & \\ & & & 1 & & \\ & & 1 & & & \\ & 1 & & & & \\ 1 & & & & & \end{vmatrix}$

$ \begin{aligned} &= \xleftrightarrow{R_1 \leftrightarrow R_6} (-1) \begin{vmatrix} 1 & & & & & \\ & 1 & & & & \\ & & 1 & & & \\ & & & 1 & & \\ & & & & 1 & \\ & & & & & 1 \end{vmatrix} \\ &= \xleftrightarrow{R_2 \leftrightarrow R_5} (-1)(-1) \begin{vmatrix} 1 & & & & & \\ & 1 & & & & \\ & & 1 & & & \\ & & & 1 & & \\ & & & & 1 & \\ & & & & & 1 \end{vmatrix} \\ &= \xleftrightarrow{R_3 \leftrightarrow R_4} (-1)(-1)(-1) \begin{vmatrix} 1 & & & & & \\ & 1 & & & & \\ & & 1 & & & \\ & & & 1 & & \\ & & & & 1 & \\ & & & & & 1 \end{vmatrix} \\ &= -1 \end{aligned} $ <p>substituting $n = 6$ in the solution</p> $(-1)^{\lfloor \frac{6}{2} \rfloor} = -1$ <p>Hence the proved that the solution is correct.</p>

TABLE 2: Example