

# EE5609 Matrix Theory

Kranthi Kumar P

Download the python code from

[https://github.com/kranthiakssy/AI20RESCH14002\\_PhD\\_IITH/tree/master/EE5609\\_Matrix\\_Theory/Assignment-2](https://github.com/kranthiakssy/AI20RESCH14002_PhD_IITH/tree/master/EE5609_Matrix_Theory/Assignment-2)

and latex-file codes from

[https://github.com/kranthiakssy/AI20RESCH14002\\_PhD\\_IITH/tree/master/EE5609\\_Matrix\\_Theory/Assignment-2](https://github.com/kranthiakssy/AI20RESCH14002_PhD_IITH/tree/master/EE5609_Matrix_Theory/Assignment-2)

## ASSIGNMENT-2

### Problem

Lines and Planes (81) : In each of the following cases, determine the normal to the plane and the distance from the origin.

- (a)  $\begin{pmatrix} 0 & 0 & 1 \end{pmatrix} \mathbf{x} = 2$
- (b)  $\begin{pmatrix} 1 & 1 & 1 \end{pmatrix} \mathbf{x} = 1$
- (c)  $\begin{pmatrix} 0 & 5 & 0 \end{pmatrix} \mathbf{x} = -8$
- (d)  $\begin{pmatrix} 2 & 3 & -1 \end{pmatrix} \mathbf{x} = 5$

### Solution

If  $ax+by+cz = k$  is a linear equation representing a plane, then the normal  $\mathbf{n}$  to that plane is the coefficients of the linear equation

$$\mathbf{n} = \begin{pmatrix} a \\ b \\ c \end{pmatrix} \quad (0.0.1)$$

The equation of plane can be written as

$$\mathbf{n}^T \mathbf{x} = c \quad (0.0.2)$$

The shortest distance between the plane (0.0.2) and origin is

$$\frac{|c|}{\|\mathbf{n}\|} \quad (0.0.3)$$

$$a) \begin{pmatrix} 0 & 0 & 1 \end{pmatrix} \mathbf{x} = 2$$

$$\text{normal vector, } \mathbf{n} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad (0.0.4)$$

$$c = 2 \quad (0.0.5)$$

As per (0.0.3), shortest distance from origin =

$$\frac{|2|}{\sqrt{0^2 + 0^2 + 1^2}} = 2 \quad (0.0.6)$$

$$b) \begin{pmatrix} 1 & 1 & 1 \end{pmatrix} \mathbf{x} = 1$$

$$\text{normal vector, } \mathbf{n} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad (0.0.7)$$

$$c = 1 \quad (0.0.8)$$

As per (0.0.3), shortest distance from origin =

$$\frac{|1|}{\sqrt{1^2 + 1^2 + 1^2}} = \frac{1}{\sqrt{3}} \quad (0.0.9)$$

$$c) \begin{pmatrix} 0 & 5 & 0 \end{pmatrix} \mathbf{x} = -8$$

$$\text{normal vector, } \mathbf{n} = \begin{pmatrix} 0 \\ 5 \\ 0 \end{pmatrix} \quad (0.0.10)$$

$$c = -8 \quad (0.0.11)$$

As per (0.0.3), shortest distance from origin =

$$\frac{|-8|}{\sqrt{0^2 + 5^2 + 0^2}} = \frac{8}{5} \quad (0.0.12)$$

$$d) \begin{pmatrix} 2 & 3 & -1 \end{pmatrix} \mathbf{x} = 5$$

$$\text{normal vector, } \mathbf{n} = \begin{pmatrix} 2 \\ 3 \\ -1 \end{pmatrix} \quad (0.0.13)$$

$$c = 5 \quad (0.0.14)$$

As per (0.0.3), shortest distance from origin =

$$\frac{|5|}{\sqrt{2^2 + 3^2 + (-1)^2}} = \frac{5}{\sqrt{14}} \quad (0.0.15)$$