

K 223 - Angular $\mu\mu$ - correlations

Preparation tasks

1 - Which distances?

▷ How does the count rate of one detector change with the distance?

$$N \propto \frac{1}{r^2}$$

▷ How does the coincidence rate change with the distance?

$$N_c \propto \frac{1}{r^2}$$

▷ How does the asymmetry seen in the experiment differ from the theoretical prediction?

We can only move one detector? $\rightarrow \frac{1}{r}$

▷ How do the factors influence the error?

$$R_{kk} = \frac{R_{kk}^{exp}}{Q_{kk}} \quad \Delta R_{kk}^2 = \left(\frac{1}{Q_{kk}} \Delta R_{kk}^{exp} \right)^2 + \left(\frac{R_{kk}^{exp}}{Q_{kk}^2} \Delta Q_{kk} \right)^2$$

$\rightarrow 0 \quad Q_{kk} \rightarrow 1$

2 - Which angles?


▷ most important maximum and falling flanks

3 - How to correct for de-adjustment?

▷ if the source is not aligned we do not measure isotropically distributed angular photons

▷ renormalize $n_c \rightarrow \frac{n_c(\theta_0)}{n_c(\theta)} \cdot n_c$

Knowledge



▷ $W(\theta) \propto P_e(\cos \theta) (Chk) \quad P(\mu) = \sum_{\mu_i} G(\mu_i \rightarrow \mu) F_{e \mu \mu}(\theta_1)$

▷ $W(\theta) = \sum_{\mu_i \mu_f} P(\mu) G(\mu \rightarrow \mu_f) F_{e \mu \mu}(\theta)$

$|\theta_1 - \theta_2|$

▷ distributed by HFI interactions

▷ not all unpaired coincidences are real → random coincidences

