


Determination of the beam asymmetry Σ in η - and η' -photoproduction off the proton using Bayesian statistics

Master thesis for the CBELSA/TAPS collaboration

JAKOB KRAUSE

✉ krause@hiskp.uni-bonn.de |  [krausejm](#)

Supervisor: JUN. PROF. DR. ANNIKA THIEL

✉ thiel@hiskp.uni-bonn.de


30th March 2022

Setting the scene

The Standard Model of Particle Physics


- ▶ matter consists of 12 (anti-) *fermions*
- ▶ quarks interact via *strong interaction*
- ▶ form bound states: mesons ($q\bar{q}$) and baryons (qqq)

baryon spectroscopy (photoproduction) gives insight in strong interaction



Setting the scene

Observe resonances N^*/Δ^* in the cross sections $\sigma(\gamma p \rightarrow pM)$



Total cross section $\sigma(\gamma p \rightarrow p\pi^0)$ [Wunderlich et al. 2017]

→goal: (help to) identify contributing resonances as strong bound states!

1. Experimental Setup

2. Preliminary results

3. Conclusion

Theoretical basics

- ▶ resonances are broad, overlapping, require complicated partial-wave-analysis (PWA)
- ▶ constraints for the analysis can be derived from polarization observables
- ▶ ultimate goal: "complete experiment"; unambiguous, model-independent PWA solution → several single and double polarization observables needed

Beam-target polarization observables

photon		target polarization		
		x	y	z
unpolarized	σ_0	-	T	-
linearly polarized	$-\Sigma$	H	$-P$	$-G$
circularly polarized	-	F	-	$-E$

[san]


Theoretical basics

Beam asymmetry Σ

$$\frac{d\sigma}{d\Omega}(E_\gamma, \cos\theta, \varphi) = \frac{d\sigma}{d\Omega_0}(E_\gamma, \cos\theta) \cdot \left[1 - p_\gamma^{\text{lin}} \Sigma \cos(2\varphi) \right]$$

polarization angle φ , polarization degree p_γ^{lin}


[san]



Definition of the polarization angle

CBELSA/TAPS experiment

- ▶ generate photon beam from accelerated electrons via bremsstrahlung, with $E_\gamma \leq 3.2 \text{ GeV}$
- ▶ photon beam impinges on liquid hydrogen target:
 $\gamma p \rightarrow pM \rightarrow pX$
- ▶ measure decay products X of different final states:
 $M = \pi^0/\eta/\eta'/\dots$
- ▶ data set:
July-October 2013,
1065 h beam time




Overview of the experimental area, adapted from [cb]

Event selection of the $\eta' \rightarrow \gamma\gamma$ final state

Analysis performed in 2x6 bins of $(E_\gamma, \cos \theta_{\eta'}^{\text{CMS}})$, $E_\gamma \in [1400, 1800]$ MeV

- ▶ 3 detector hits, 2 uncharged, 1 charged
- ▶ coincident detector hits
- ▶ kinematic cuts derived from energy-momentum conservation
$$p_\gamma + p_p = p'_p + p_{\eta'}$$
- ▶ additional cuts to reduce background contributions

total: ~ 11000 $\eta' \rightarrow \gamma\gamma$ events




Extraction method for $\Sigma_{\eta'}$

- ▶ measure in 2 distinct orthogonal polarization settings \perp, \parallel


- ▶ χ^2 -fit to event yield asymmetries

$$A(E_\gamma, \theta, \phi) = \frac{N^\perp(E_\gamma, \theta, \phi) - N^\parallel(E_\gamma, \theta, \phi)}{p_\gamma^\parallel N^\perp(E_\gamma, \theta, \phi) + p_\gamma^\perp N^\parallel(E_\gamma, \theta, \phi)} = \Sigma(E_\gamma, \theta) \cos(2(\alpha^\parallel - \phi))$$

- ▶ fit from ~ 800 $\eta' \rightarrow \gamma\gamma$ events



Preliminary results for $\Sigma_{\eta'}$



Beam asymmetry $\Sigma_{\eta'}$ for all energy and angle bins, compared with results of [clas]

Confirming pre-published results of Σ_η

- ▶ Polarization observables are needed for different final states ($\pi^0, \eta, \eta', \dots$)
- ▶ results for η have already been published [eta]
- ▶ goal: confirm results using different fitting method

Confirming pre-published results for Σ_η

Event selection (η)

analysis performed in 11x12 bins of $(E_\gamma, \cos \theta)$ by [eta]

Method

- ▶ fit to event yield asymmetries using BAYESIAN inference
- ▶ BAYES' theorem: $p(\theta|y) \propto \mathcal{L}(y|\theta) \cdot \pi(\theta)$
 - ▶ *marginal posteriors*: $p(\theta_j|y) = \prod_{i \neq j} \int d\theta_i p(\theta|y)$
 - ▶ obtained using MARKOV-chain-Monte-Carlo (MCMC)

sampling algorithms: STAN



Confirming pre-published results for Σ_η

Event selection (n)

analy

Before


Point estimate: $\Sigma_{\chi^2} \pm \Delta(\Sigma_{\chi^2})$

Now

marginal posteriors: $p(\Sigma|y)$

Meth


- f
- l



Confirming pre-published results for Σ_η

`../bayes/realdeal/sigma_eta_alt.pdf`

Confirming pre-published results for Σ_η



Beam asymmetry Σ for one energy and all angle bins

Additional advantage: sample only in physically allowed parameter space

Conclusion

Summary

- ▶ Σ extracted for η and η' final state
- ▶ η results obtained with BAYESIAN fit agree with previous results
- ▶ η' results agree with previous results

Outlook

- ▶ extract Σ using unbinned maximum likelihood fit for η/η'
- ▶ apply BAYESIAN approach to above method
- ▶ consider bkg contaminations in results of $\Sigma_{\eta'}$

BACKUP & REFERENCES

Additional theoretical basics

Unpolarized differential cross section

$$\frac{d\sigma}{d\Omega} = \frac{1}{4}\rho \sum_{\text{spins}} |\langle f | \mathcal{F} | i \rangle|^2,$$

where


$$\mathcal{F} = i(\vec{\sigma} \cdot \vec{\epsilon})F_1 + (\vec{\sigma} \cdot \hat{q})(\vec{\sigma} \cdot (\hat{k} \times \vec{\epsilon}))F_2 + i(\vec{\sigma} \cdot \hat{k})(\hat{q} \cdot \vec{\epsilon})F_3 + i(\vec{\sigma} \cdot \hat{q})(\hat{q} \cdot \vec{\epsilon})F_4$$

F_i : complex CGLN Amplitudes

[cgln]

$\frac{d\sigma}{d\Omega} \in \mathbb{R}$, not sufficient to determine \mathcal{F} unambiguously
→ Polarization Observables can be related to F_i

Background estimation using Monte-Carlo simulations




Background estimation using Monte-Carlo simulations

$2\pi^0/\pi^0\eta$ events pass event selection, because $E_{\gamma_i} \lesssim 20$ MeV, or $\theta_{\gamma_i} \approx \theta_{\gamma_j}$

Background reducing cuts

- ▶ p in MT for $E_\gamma < 1500$ MeV
- ▶ $E_{\gamma_i} < 1500$ MeV
- ▶ 1 PED/Cluster for γ_i
- ▶ $\text{Clustersize}(p) < 6$
- ▶ $\text{Clustersize}(\gamma_i)$ in FW




bkg contamination $\sim 13\%$

Diagnostics of a BAYESIAN fit

- ▶ \hat{R} : measure of convergence for chains
- ▶ Monte-Carlo-Standard-Error: measure for adequate sample size
- ▶ *posterior predictive checks*: "goodness of fit"

../bayes/realdeal/ppd_checks_alt/ebin6costbin1.pdf

References I

-  Wunderlich, Y. et al. (May 2017). ‘Determining the dominant partial wave contributions from angular distributions of single- and double-polarization observables in pseudoscalar meson photoproduction’. In: *The European Physical Journal A* 53.5. ISSN: 1434-601X. DOI: 10.1140/epja/i2017-12255-0. URL: <http://dx.doi.org/10.1140/epja/i2017-12255-0>.