

# Determination of the target asymmetry $T$ in $\eta'$ photoproduction

JAKOB MICHAEL KRAUSE

Masterarbeit in Physik  
angefertigt im Helmholtz-Institut für Strahlen- und  
Kernphysik

vorgelegt der  
Mathematisch-Naturwissenschaftlichen Fakultät  
der  
Rheinischen Friedrich-Wilhelms-Universität  
Bonn

Sep 2022

DRAFT

I hereby declare that this thesis was formulated by myself and that no sources or tools other than those cited were used.

Bonn, .....  
Date

.....  
Signature

1. Gutachter: JUN. PROF. DR. ANNIKA THIEL
2. Gutachterin: Prof. Dr. TBD

# Contents

---

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Photoproduction of Pseudoscalar Mesons . . . . .	4
1.2	Motivation and Structure of this Thesis . . . . .	4
<b>2</b>	<b>Experimental Setup</b>	<b>5</b>
2.1	Production of (polarized) high energy photon beam . . . . .	5
2.2	Beam Target . . . . .	6
2.3	Calorimeters . . . . .	6
2.4	Trigger . . . . .	6
<b>A</b>	<b>Useful information</b>	<b>9</b>
	<b>Bibliography</b>	<b>11</b>
	<b>List of Figures</b>	<b>13</b>
	<b>List of Tables</b>	<b>15</b>

DRAFT

## Introduction

The *Standard Model of Particle Physics* (SM) is the most successful model aiming to describe the particles and forces of the universe. It distinguishes between *fermions* and *bosons*. While all matter consists of fermions, bosons are particles that mediate the fundamental interactions.

Matter consists of (anti-)quarks and (anti-)leptons with three generations of each. Table 1.1 shows all elementary fermions including some of their most important properties. Only the first and lightest generation consists of stable particles, i.e. the up and down quark as well as the electron and its neutrino. All other particles are heavier and not stable, they will thus decay fast via the strong, electromagnetic or weak interaction.


There are in fact four interactions described by the SM: strong, electromagnetic, weak and gravitational interaction<sup>1</sup>, where gravitation is mentioned here for the sake of completeness; on the mass scale of elementary particles gravitation is negligible. Strong and weak interaction are restricted to a finite range of the order of the nucleon radius, whereas electromagnetic interaction and gravitation have infinite range. Each interaction has its own coupling (charge). The strong interaction is mediated by gluons and couples to the color charge.

	Generation			el. charge	color charge
	1	2	3		
Quarks	$u$	$c$	$t$	2/3	r,g,b
	$d$	$s$	$b$	1/3	r,g,b
Leptons	$e$	$\mu$	$\tau$	-1	-
	$\nu_e$	$\nu_\mu$	$\nu_\tau$	0	-

**Table 1.1:** Summary of the particles of the SM

Gluons and quarks carry color charge and thus interact strongly. However, an isolated quark or gluon has not been observed. Only color neutral bound systems of quarks are seen, which are called hadrons. Hadrons with integer spin are called mesons and those with half-integer spin are called baryons. Color neutrality demands mesons consist of at least one quark and one anti-quark and baryons consist of at least three quarks.

<sup>1</sup> they are ordered here according to their relative strength



**Figure 1.1:** Running coupling of QCD. The colored data points represent different methods to obtain a value for  $\alpha_s$ . For more details it may be referred to [Zyl+20].


As already mentioned, isolated quarks are not seen. This can be understood in terms of the strong coupling constant  $\alpha_s$ . The coupling constant is a measure of the strength of the strong interaction. Because it is highly dependent on the momentum transfer in the observed strong reaction it is also called running coupling constant, which is depicted in figure 1.1.

For low ( $< 1$  GeV) momentum transfers or large distances the coupling constant approaches infinity whereas it decreases for high ( $\gg 1$  GeV) momentum transfers or short distances. These momentum ranges are referred to as *confinement* and *asymptotic freedom*, respectively; quarks are confined to remain in a bound state since if one tried to pull them apart the color field becomes so strong it will create a new quark anti-quark pair resulting in two new bound states. On the other hand, bound quarks behave quasi-free and can be described using perturbative quantum chromodynamics (pQCD) if probed at sufficiently large momentum transfers.

It is more difficult however to describe QCD at momentum scales of  $\approx 1$  GeV since the coupling is too strong to justify a perturbative approach. Thus explicit modeling of QCD bound states is inevitable. One possibility is to describe baryons consisting of constituent quarks which are bound in a potential. Constituent quark models assume baryons are made up of three constituent quarks with effective masses differing from the bare quark mass. The effective mass is made up mostly from a sea of quark anti-quark pairs and gluons which surround the bare (valence) quarks. The explicit form of the binding potential is determined for each model.

The Bonn model [LMP01], for example, is formulated as a relativistically covariant constituent quark model. A potential increasing linearly with the distance is employed to adequately describe confinement. The binding potential between the constituent quarks is described by an instanton-induced interaction. Baryon resonances are then states with an orbital or angular excitation of one of the quarks. Figure 1.2 shows computed nucleon, that is Isospin  $I = 1/2$  resonances, of the Bonn model [LMP01] on the left side of each column. These are compared to measured resonances and their PDG rating [Zyl+20] in the middle. Uncertainties are indicated by the colored areas. The resonances are

identified by their total angular momentum and their parity  $J\pi$ . In addition also the total internal angular momentum along with isospin and again the total angular momentum  $L_{2T2J}$  is given. While




**Figure 1.2:** Calculated nucleon (isospin  $I = 1/2$ ) resonances compared to measurements. Left in each column are the calculations [LMP01], the middle shows the measurements and PDG rating [Zyl+20]

generally good agreement exists for low lying resonances, especially for high masses there are much more resonances predicted than actually found. This is also known as the problem of the “missing resonances” indicating the poor understanding of QCD in the non-perturbative region. This can be due to several reasons: most of the knowledge about nucleon resonances and their properties was obtained investigating the  $\pi N$  channel, biasing the data for resonances coupling weakly to this channel. Furthermore, the number of excited states with definite quantum numbers is related directly to the effective number of degrees-of-freedom accessible to the underlying theory. As a consequence, the number of degrees-of-freedom should be obtainable by comparing the measured states to the predicted states. Since nucleon resonances decay dominantly hadronic, their resonances are broad and overlapping. Thus on one hand the determination of excitation spectra proves to be a challenge on its own, demanding sophisticated methods, such as partial wave analysis (PWA). On the other hand it is not yet clear how many effective degrees-of-freedom exist for the nucleon in a constituent quark model. They could for example be decreased if the nucleon were made up of a quark and a di-quark structure. In either case it should prove fruitful to investigate the photoproduction of mesons off the nucleon for different final states to access the resonances and transitions between them which are of interest. This should ultimately add to the understanding of QCD in the non-perturbative regime. [KS03]

## 1.1 Photoproduction of Pseudoscalar Mesons

From the scattering theory point of view, photoproduction of mesons is well understood [KS03].



**Figure 1.3:** FEYNMAN diagram for the s-channel photoproduction of pseudoscalar mesons, adapted from [Afz19]

The analysis requires partial wave decomposition in both initial and final states [DT92]

## 1.2 Motivation and Structure of this Thesis

bla



## Experimental Setup

As motivated in the previous chapter, it is promising to study the photoproduction of pseudoscalar mesons in order to determine a complete set of polarization observables. This requires a polarized photon beam and/or a polarized target. It is convenient to study photoproduction off a fixed target and investigate the resonances that occur in the process. Incidentally, these resonances can only be accessed via their decay products, such that suitable calorimeters are needed. The CBELSA/TAPS experiment is located in Bonn at the ELectron Stretcher Accelerator (ELSA), which can be used to generate a high energy photon beam using the *bremsstrahlung* process, meets all above mentioned requirements. This chapter will elaborate on the already mentioned parts of the experiment that was used to collect the data needed for the determination of the target asymmetry in  $\eta'$  photoproduction.

### 2.1 Production of (polarized) high energy photon beam

First of all, a high energy photon beam has to be produced.




Figure 2.1: [Wal]




Figure 2.2: [Wal]

## 2.2 Beam Target



Figure 2.3: [Wal]

## 2.3 Calorimeters

## 2.4 Trigger




Figure 2.4: D. WALTHER in [Urb17]





Figure 2.5: [Wal]



**Figure 2.6:** [Wal]

## Useful information

---

In the appendix you usually include extra information that should be documented in your thesis, but not interrupt the flow.

The L<sup>A</sup>T<sub>E</sub>X WikiBook [[latexwiki](#)] is a useful source of information on L<sup>A</sup>T<sub>E</sub>X.



# Bibliography

---

- [Zyl+20] P. Zyla et al., *Review of Particle Physics*, PTEP **2020** (2020) 083C01 (cit. on pp. 2, 3).
- [LMP01] U. Löring, B. Metsch and H. Petry, *The light-baryon spectrum in a relativistic quark model with instanton-induced quark forces*,  
The European Physical Journal A **10** (2001) 395, ISSN: 1434-601X,  
URL: <http://dx.doi.org/10.1007/s100500170105> (cit. on pp. 2, 3).
- [KS03] B. Krusche and S. Schadmand,  
*Study of nonstrange baryon resonances with meson photoproduction*,  
Prog. Part. Nucl. Phys. **51** (2003) 399, arXiv: [nuc1-ex/0306023](https://arxiv.org/abs/nuc1-ex/0306023) (cit. on pp. 3, 4).
- [Afz19] F. N. Afzal, *Measurement of the beam and helicity asymmetries in the reactions  $\gamma p \rightarrow p\pi^0$  and  $\gamma p \rightarrow p\eta$* ,  
PhD thesis: Rheinische Friedrich-Wilhelms-Universität Bonn, 2019,  
URL: <https://hdl.handle.net/20.500.11811/8064> (cit. on p. 4).
- [DT92] D. Drechsel and L. Tiator, *Threshold pion photoproduction on nucleons*,  
J. Phys. G **18** (1992) 449 (cit. on p. 4).
- [Wal] D. Walther, *Crystal Barrel, A 4 $\pi$  photon spectrometer*,  
URL: <https://www.cb.uni-bonn.de> (visited on 27/09/2021) (cit. on pp. 5–8).
- [Urb17] M. Urban, *Design eines neuen Lichtpulsersystems sowie Aufbau und Inbetriebnahme der neuen APD Auslese für das Crystal-Barrel-Kalorimeter*,  
PhD thesis: Rheinische Friedrich-Wilhelms-Universität Bonn, 2017 (cit. on p. 7).





# List of Figures

---

1.1	Running coupling of QCD. The colored data points represent different methods to obtain a value for $\alpha_s$ . For more details it may be referred to [Zyl+20]. . . . .	2
1.2	Calculated nucleon (isospin $I = 1/2$ ) resonances compared to measurements. Left in each column are the calculations [LMP01], the middle shows the measurements and PDG rating [Zyl+20] . . . . .	3
1.3	FEYNMAN diagram for the s-channel photoproduction of pseudoscalar mesons, adapted from [Afz19] . . . . .	4
2.1	[Wal] . . . . .	5
2.2	[Wal] . . . . .	6
2.3	[Wal] . . . . .	6
2.4	D. WALTHER in [Urb17] . . . . .	7
2.5	[Wal] . . . . .	7
2.6	[Wal] . . . . .	8



# List of Tables

---

1.1 Summary of the particles of the SM . . . . . 1