

Recent Polarization Observable Results in η - and η' -photoproduction off the proton

Master thesis for the CBELSA/TAPS collaboration

JAKOB KRAUSE

✉ krause@hiskp.uni-bonn.de | 🌐 krausejm

Supervisor: JUN. PROF. DR. ANNIKA THIEL

✉ thiel@hiskp.uni-bonn.de


30th March 2022

Setting the scene

The Standard Model of Particle Physics


- ▶ matter consists of 12 (anti-) *fermions*
- ▶ quarks interact via *strong interaction*
- ▶ form bound states: mesons ($q\bar{q}$) and baryons (qqq)

baryon spectroscopy (photoproduction) gives insight in strong interaction



Setting the scene

Observe resonances N^*/Δ^* in the cross sections $\sigma(\gamma p \rightarrow pM)$



Total cross section $\sigma(\gamma p \rightarrow p\pi^0)$ [Wunderlich et al. 2017]

→goal: (help to) identify contributing resonances as strong bound states!

1. Theoretical basics
2. Experimental Setup
3. Preliminary results
4. Conclusion

Theoretical basics

- ▶ resonances are broad, overlapping, require complicated partial-wave-analysis (PWA)
- ▶ constraints for the analysis can be derived from polarization observables
- ▶ ultimate goal: "complete experiment"; unambiguous, model-independent PWA solution → several single and double polarization observables needed

Beam-target polarization observables

photon		target polarization		
		x	y	z
unpolarized	σ_0	-	T	-
linearly polarized	$-\Sigma$	H	$-P$	$-G$
circularly polarized	-	F	-	$-E$

[Sandorfi et al. 2011]


Theoretical basics

Beam asymmetry Σ

$$\frac{d\sigma}{d\Omega}(E_\gamma, \cos\theta, \varphi) = \frac{d\sigma}{d\Omega_0}(E_\gamma, \cos\theta) \cdot \left[1 - p_\gamma^{\text{lin}} \Sigma \cos(2\varphi)\right]$$

polarization angle φ , polarization degree p_γ^{lin}


[Sandorfi et al. 2011]



Definition of the polarization angle

CBELSA/TAPS experiment

- ▶ generate photon beam from accelerated electrons via bremsstrahlung, with $E_\gamma \leq 3.2 \text{ GeV}$
- ▶ photon beam impinges on liquid hydrogen target:
 $\gamma p \rightarrow pM \rightarrow pX$
- ▶ measure decay products X of different final states:
 $M = \pi^0/\eta/\eta'/\dots$
- ▶ data set:
July-October 2013,
1065 h beam time




Overview of the experimental area, adapted from [Walther 2022]

Event selection of the $\eta' \rightarrow \gamma\gamma$ final state

Analysis performed in 2x6 bins of $(E_\gamma, \cos \theta_{\eta'}^{\text{CMS}})$, $E_\gamma \in [1400, 1800]$ MeV

- ▶ 3 detector hits, 2 uncharged, 1 charged
- ▶ coincident detector hits
- ▶ kinematic cuts derived from energy-momentum conservation
$$p_\gamma + p_p = p'_p + p_{\eta'}$$
- ▶ additional cuts to reduce background contributions

total: ~ 11000 $\eta' \rightarrow \gamma\gamma$ events




Extraction method for $\Sigma_{\eta'}$

- ▶ measure in 2 distinct orthogonal polarization settings \perp, \parallel


- ▶ χ^2 -fit to event yield asymmetries

$$A(E_\gamma, \theta, \phi) = \frac{N^\perp(E_\gamma, \theta, \phi) - N^\parallel(E_\gamma, \theta, \phi)}{p_\gamma^\parallel N^\perp(E_\gamma, \theta, \phi) + p_\gamma^\perp N^\parallel(E_\gamma, \theta, \phi)} = \Sigma(E_\gamma, \theta) \cos(2(\alpha^\parallel - \phi))$$

- ▶ fit from ~ 800 $\eta' \rightarrow \gamma\gamma$ events



Preliminary results for $\Sigma_{\eta'}$



Beam asymmetry $\Sigma_{\eta'}$ for all energy and angle bins, compared with results of [Collins et al. 2017]

Confirming pre-published results of Σ_η

Preliminary results (η)

Event selection (η)

analysis performed in 11x12 bins of $(E_\gamma, \cos \theta)$ by [Afzal et al. 2020]

Method


- ▶ fit to event yield asymmetries using BAYESIAN inference
- ▶ BAYES' theorem: $p(\theta|y) \propto \mathcal{L}(y|\theta) \cdot \pi(\theta)$
 - ▶ *marginal posteriors*: $p(\theta_j|y) = \prod_{i \neq j} \int d\theta_i p(\theta|y)$
 - ▶ obtained using MARKOV-chain-Monte-Carlo (MCMC)

sampling algorithms: STAN
[Stan development team 2022]




(marginal) posteriors $p(\Sigma|y) \leftrightarrow \Sigma_{\chi^2} \pm \Delta(\Sigma_{\chi^2})$

Preliminary results (η)



Beam asymmetry Σ for all energy and angle bins

Preliminary results (η)



Beam asymmetry Σ for one energy and all angle bins

Additional advantage: sample only in physically allowed parameter space

Conclusion

Summary

- ▶ Σ extracted for η and η' final state
- ▶ η results obtained with BAYESIAN fit agree with previous results
- ▶ η' results agree with previous results

Outlook

- ▶ extract Σ using unbinned maximum likelihood fit for η/η'
- ▶ apply BAYESIAN approach to above method
- ▶ consider bkg contaminations in results of $\Sigma_{\eta'}$

BACKUP & REFERENCES

Additional theoretical basics

Unpolarized differential cross section

$$\frac{d\sigma}{d\Omega} = \frac{1}{4}\rho \sum_{\text{spins}} |\langle f | \mathcal{F} | i \rangle|^2,$$

where


$$\mathcal{F} = i(\vec{\sigma} \cdot \vec{\epsilon})F_1 + (\vec{\sigma} \cdot \hat{q})(\vec{\sigma} \cdot (\hat{k} \times \vec{\epsilon}))F_2 + i(\vec{\sigma} \cdot \hat{k})(\hat{q} \cdot \vec{\epsilon})F_3 + i(\vec{\sigma} \cdot \hat{q})(\hat{q} \cdot \vec{\epsilon})F_4$$

F_i : complex CGLN Amplitudes

[Chew et al. 1957]

$\frac{d\sigma}{d\Omega} \in \mathbb{R}$, not sufficient to determine \mathcal{F} unambiguously
→ Polarization Observables can be related to F_i

Background estimation using Monte-Carlo simulations




Background estimation using Monte-Carlo simulations

$2\pi^0/\pi^0\eta$ events pass event selection, because $E_{\gamma_i} \lesssim 20$ MeV, or $\theta_{\gamma_i} \approx \theta_{\gamma_j}$

Background reducing cuts

- ▶ p in MT for $E_\gamma < 1500$ MeV
- ▶ $E_{\gamma_i} < 1500$ MeV
- ▶ 1 PED/Cluster for γ_i
- ▶ $\text{Clustersize}(p) < 6$
- ▶ $\text{Clustersize}(\gamma_i)$ in FW




bkg contamination $\sim 13\%$




Diagnostics of a BAYESIAN fit

- ▶ \hat{R} : measure of convergence for chains
- ▶ Monte-Carlo-Standard-Error: measure for adequate sample size
- ▶ *posterior predictive checks*: "goodness of fit"





$$1490 \leq E_\gamma < 1550, \quad -0.83 \leq \cos \theta_\eta^{\text{CMS}} < -0.67$$



References I

-  Afzal, F. et al. (Oct. 2020). ‘Observation of the $p\eta'$ Cusp in the New Precise Beam Asymmetry Σ Data for $\gamma p \rightarrow p\eta'$ '. In: *Phys. Rev. Lett.* 125 (15), p. 152002. DOI: 10.1103/PhysRevLett.125.152002. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.125.152002>.
-  Chew, G. F. et al. (June 1957). ‘Relativistic Dispersion Relation Approach to Photomeson Production’. In: *Phys. Rev.* 106 (6), pp. 1345–1355. DOI: 10.1103/PhysRev.106.1345. URL: <https://link.aps.org/doi/10.1103/PhysRev.106.1345>.
-  Collins, P. et al. (2017). ‘Photon beam asymmetry Σ for η and η' photoproduction from the proton’. In: *Phys. Lett. B* 771, pp. 213–221. DOI: 10.1016/j.physletb.2017.05.045. arXiv: 1703.00433 [nucl-ex].

References II

-  Sandorfi, A. M. et al. (Apr. 2011). ‘Determining pseudoscalar meson photoproduction amplitudes from complete experiments’. In: *Journal of Physics G: Nuclear and Particle Physics* 38.5, p. 053001. ISSN: 1361-6471. DOI: 10.1088/0954-3899/38/5/053001. URL: <http://dx.doi.org/10.1088/0954-3899/38/5/053001>.
-  Stan development team (2022). *Stan Modeling Language Users Guide and Reference Manual*. Vol. 2.29. URL: <https://mc-stan.org>.
-  Walther, Dieter (2022). *Crystal Barrel. A 4π photon spectrometer*. URL: <https://www.cb.uni-bonn.de> (visited on 09/03/2022).
-  Wunderlich, Y. et al. (May 2017). ‘Determining the dominant partial wave contributions from angular distributions of single- and double-polarization observables in pseudoscalar meson photoproduction’. In: *The European Physical Journal A* 53.5. ISSN: 1434-601X. DOI: 10.1140/epja/i2017-12255-0. URL: <http://dx.doi.org/10.1140/epja/i2017-12255-0>.