

Determination of the beam asymmetry Σ in η - and η' -photoproduction

JAKOB MICHAEL KRAUSE

Masterarbeit in Physik
angefertigt im Helmholtz-Institut für Strahlen- und
Kernphysik

vorgelegt der
Mathematisch-Naturwissenschaftlichen Fakultät
der
Rheinischen Friedrich-Wilhelms-Universität
Bonn

Sep 2022

DRAFT

I hereby declare that this thesis was formulated by myself and that no sources or tools other than those cited were used.

Bonn,
Date

.....
Signature

1. Gutachter: JUN. PROF. DR. ANNIKA THIEL
2. Gutachterin: Prof. Dr. TBD

DRAFT

Contents

1	Introduction	1
1.1	Photoproduction of Pseudoscalar Mesons	4
1.2	Measurement of Polarization Observables	5
1.3	Introduction to BAYESIAN statistics	5
1.4	Motivation and Structure of this Thesis	5
2	Experimental Setup	7
2.1	Production of (polarized) high energy photon beam	7
2.1.1	Tagger	8
2.2	Beam Target	8
2.3	Calorimeters	8
2.4	Trigger	8
3	Event selection	11
3.1	Preselection and charge cut	11
3.2	Time of particles	12
3.3	Kinematic constraints	14
3.3.1	Coplanarity	14
3.3.2	Polar angle difference	14
3.3.3	Missing mass	15
3.3.4	Invariant mass	15
3.4	Determination of cut ranges	15
3.5	Investigation of of background and additional cuts	15
4	Determination of the beam asymmetry Σ_η	15
4.1	BAYESIAN fit to event yield asymmetries	15
4.1.1	Application of method to toy Monte Carlo data	15
4.1.2	Application of method to real data	15
4.2	Event based fit	15
4.2.1	Application of method to toy Monte Carlo data	15
4.2.2	Application of method to real data	15
4.3	Discussion	15
5	Determination of the beam asymmetry $\Sigma_{\eta'}$	17
5.1	Fit to event yield asymmetries	17
5.1.1	Application of method to toy Monte Carlo data	17

5.1.2	Application of method to real data	17
5.2	Event based fit	17
5.2.1	Application of method to toy Monte Carlo data	17
5.2.2	Application of method to real data	17
5.3	Discussion	17
6	Summary and outlook	19
A	Useful information	21
	Bibliography	23
	List of Figures	25
	List of Tables	27

DRAFT

Event selection

The determination of polarization observables needs to be completed for particular reactions (cf. chapter 1), such as the photoproduction of e.g. a single η' meson. However, the recorded events contain data from the decay products of all possible final states in addition to combinatorical background. Thus, event candidates for the desired reaction have to be extracted before they are considered for further analysis. Table 3.1 shows the five most probable decay modes of the η' meson. Three of these result in final states which only contain photons and are thus reliably measurable with the CBELSA/TAPS experiment. Only the $\eta' \rightarrow \gamma\gamma$ decay channel was considered for further analysis; the $\omega\gamma$ channel provides negligible statistics and considering the acceptance of detecting six photons in the final state, the expected yield of the $\eta' \rightarrow \gamma\gamma$ decays should be roughly equal to the $\eta' \rightarrow \pi^0\pi^0\eta \rightarrow 6\gamma$ final state [Afz22]. Offering a cleaner, three-particle final state, the $\eta' \rightarrow \gamma\gamma$ was then favored in the course of this thesis.

Decay mode		Branching ratio
$\pi^+\pi^-\eta$		42.6%
$\rho^0\gamma$	$\rightarrow \pi^+\pi^-\gamma$	28.9% (28.9%)
$\pi^0\pi^0\eta$	$\rightarrow 6\gamma$	22.8% (8.8%)
$\omega\gamma$	$\rightarrow \pi^+\pi^-\pi^0\gamma/\pi^0\gamma\gamma$	2.52% (2.2%/0.21%)
$\gamma\gamma$		2.3%

Table 3.1: The five most probable decay modes of the η' meson. The most probable further decay with according branching ratio is shown in brackets.[Zyl+20]

The process of *event selection* for the reaction $\gamma p \rightarrow p\eta' \rightarrow p\gamma\gamma$ is outlined in the following chapter. Note that in this thesis the analysis of data from η -photoproduction starts with this process already completed, which is described in detail in reference [Afz19].

3.1 Preselection and charge cut

Events are generally classified depending on the number of particle energy deposits (PED). If the complete four-momenta of three final state particles are measured, they are referred to as 3PED events. Low energy protons however may either be only detected in the scintillators of the inner, forward or

MiniTAPS detector – giving only directional information (2.5 PED) – or lost entirely (2 PED). Only 2.5PED and 3PED events were analyzed since the additional background contributions from 2PED events exceeded the additional signal contributions. It is worth noting that 3PED events are significantly dominant for $\eta' \rightarrow \gamma\gamma$ reactions; the production threshold for η' mesons is $E_\gamma = 1\,447$ MeV, such that the recoil proton will likely be detected. Figure 3.1 shows the distribution of the different event classes for $\eta' \rightarrow \gamma\gamma$ production in MONTE CARLO data, with a clear preference towards 3PED events.




Figure 3.1: Distribution of event classes in $\eta' \rightarrow \gamma\gamma$ production

To further improve the signal to background ratio, the charge information of the final state particles was utilized in the next step. In particular, to select $\eta' \rightarrow \gamma\gamma$ reactions, one charged and two uncharged particles in the final state were demanded.

3.2 Time of particles

Due to its high count rate the tagging system (see section 2.1.1) will not only record beam photons which produce the detectable final state particles, but also several uncorrelated ones. To select only beam photons which will induce a photoproduction process the time information of the detected particles is used. It is shown in figure 3.2 for all particles involved in 2.5PED and 3PED events of η' photoproduction. In all cases prompt peaks centered around 0 ns (the trigger time) are visible. Since the final state photons move with velocity c their timing information does not underlie fluctuations, as is the case for the final state proton on the contrary. The tagged, uncorrelated beam photons are visible as flat background underneath the prompt peak in the time of the beam photon. Naturally, only coincident events may be referred to as η' candidates for the further analysis and thus only events with time information of at least one final state particle are kept. Photons need to be detected in the MiniTAPS or forward detector to acquire time information. To determine coincidence it is convenient




Figure 3.2: Time information of all final state particles and the beam photon for 3PED η' production

to define the *reaction time*

$$t_{\text{reaction}} = \begin{cases} t_{\text{beam}} - t_{\text{meson}} & \text{meson time exists} \\ t_{\text{beam}} - t_{\text{recoil}} & \text{meson time does not exist,} \end{cases} \quad (3.1)$$

where the meson time t_{meson} is appointed either the averaged time of both decay photons or the time of a single photon if only one photon has time information. t_{beam} and t_{recoil} are the time of the beam photon and recoil proton, respectively. Figure 3.3 shows the reaction time for 2.5PED and 3PED events; a clear prompt peak centred at 0 is visible, the colored area indicates the chosen range of $t_{\text{reaction}} \in [-8, 5]\text{ns}$. However, this cut still contains random time background underneath the prompt peak. This may be accounted for by *sideband subtraction*, assuming the background is flat. All events residing in the prompt peak with $t_r \in [-8, 5]\text{ns}$ will be assigned a weight of $w_p = +1$ while sideband events with $t_r \in [-200, -100]\text{ns} \vee t_r \in [100, 200]\text{ns}$ will be assigned a weight of $w_s = -\frac{13}{200}$. Any histogram N that is filled in the following will then consist of prompt peak events N_{prompt} and sideband events N_{sideband}

$$N = N_{\text{prompt}} + w_s \cdot N_{\text{sideband}},$$

such that the random time background underneath the prompt peak is subtracted. In addition, the time difference between meson and proton and between the two photons is demanded to be within $[-10, 10]\text{ns}$. All described cuts to the data, including the sideband subtraction are referred to as the *time cut* in the following.




Figure 3.3: Reaction time t_r for 3PED η' production

3.3 Kinematic constraints

After charge and time cut, additional cuts can be derived from energy and momentum conservation. Let p_{beam} and p_p be the four momenta of the initial state beam photon and proton, respectively. Then

$$p_{\text{beam}} + p_p = p_{\text{recoil}} + p_{\text{meson}} \quad (3.2)$$

holds, with p_{recoil} being the momentum of the recoiling proton and p_{meson} the meson momentum.

3.3.1 Coplanarity

In the initial state there is vanishing transversal momentum p_{xy} since the target protons are at rest and the beam photon impinges in z -direction. Naturally this transversal momentum has to vanish in the final state as well, such that

$$\mathcal{P}_{xy} [p_{\text{recoil}} + p_{\text{meson}}] = 0, \quad (3.3)$$

where \mathcal{P}_{xy} is the projection operator to the transversal plane. Equation (3.3) is valid if and only if meson and proton lie back to back (coplanar) in the x - y plane, which is quantified by the difference of their azimuthal angles ϕ_{meson} and ϕ_{recoil} being 180° in the LAB-frame

$$\Delta\phi := \phi_{\text{meson}}^{\text{LAB}} - \phi_{\text{recoil}}^{\text{LAB}} \stackrel{!}{=} 180^\circ. \quad (3.4)$$

3.3.2 Polar angle difference

If all initial and final state momenta are measured, the reaction described by equation (3.2) is *overdetermined*, such that one final state particle can be treated as a "missing particle" X with

momentum p_X :

$$p_X = p_{\text{beam}} + p_p - p_{\text{meson}}. \quad (3.5)$$

One can then use

$$\Delta\theta := \theta_{p_X}^{\text{LAB}} - \theta_{p_{\text{recoil}}}^{\text{LAB}} \stackrel{!}{=} 0 \quad (3.6)$$

as a further constraint to the data.

3.3.3 Missing mass

The previously described angular cuts are only applicable if all final state particles have been detected. Independently of the detection of the recoil proton the mass of the missing particle $m_X^2 = p_X^2$ can be determined and compared with the proton mass of $m_p = 938.27$ MeV [Zyl+20]. From equation (3.5) it follows that

$$m_X = \sqrt{(E_\gamma + m_p - E_{\text{meson}})^2 - p_{x,\text{meson}}^2 - p_{y,\text{meson}}^2 - (E_\gamma - p_{z,\text{meson}})^2}. \quad (3.7)$$

3.3.4 Invariant mass

The measurement of the invariant mass of the two final state photons does also not require the measurement of the recoil proton. The knowledge of both four-momenta suffices, since

$$m_{\text{meson}} = \sqrt{p_{\text{meson}}^2} = \sqrt{(p_{\gamma_1} + p_{\gamma_2})^2} = \sqrt{2E_{\gamma_1}E_{\gamma_2}(1 - \cos \alpha_{\gamma_1\gamma_2})}, \quad (3.8)$$

where E_{γ_i} are the measured photon energies and $\alpha_{\gamma_1\gamma_2}$ is the angle spanned by the two photon momenta. To select only η' candidates $m_{\text{meson}} = m_{\eta'} = 957.78$ MeV is demanded. Remarkably, the cut on the invariant mass of the final state photons is the only one to uniquely select η' production candidates so far. All other cuts apply similarly to arbitrary meson photoproduction.

3.4 Determination of cut ranges

The constraints described in the previous section must not be understood as strict equalities, cf. equations (3.3),(3.6),(3.7) and (3.8). The quantities of interest will rather describe distributions around the desired value, such that confidence intervals may be extracted by fitting to said distributions. This is done iteratively:

Let $C_I^\theta, C_J^\phi, C_K^{m_X}, C_L^{m_{\text{meson}}}$ be the cut operators that restrain the data such that $\Delta\theta, \Delta\phi, m_X$ and m_{meson} are in the interval I, J, K, L respectively.

3.5 Investigation of background and additional cuts

Bibliography

- [Zyl+20] P. Zyla et al., *Review of Particle Physics*, PTEP **2020** (2020) 083C01 (cit. on pp. 11, 15).
- [LMP01] U. Löring, B. Metsch and H. Petry, *The light-baryon spectrum in a relativistic quark model with instanton-induced quark forces*,
The European Physical Journal A **10** (2001) 395, ISSN: 1434-601X,
URL: <http://dx.doi.org/10.1007/s100500170105>.
- [KS03] B. Krusche and S. Schadmand,
Study of nonstrange baryon resonances with meson photoproduction,
Prog. Part. Nucl. Phys. **51** (2003) 399, arXiv: [nuc1-ex/0306023](https://arxiv.org/abs/nuc1-ex/0306023).
- [Afz19] F. N. Afzal, *Measurement of the beam and helicity asymmetries in the reactions $\gamma p \rightarrow p\pi^0$ and $\gamma p \rightarrow p\eta$* ,
PhD thesis: Rheinische Friedrich-Wilhelms-Universität Bonn, 2019,
URL: <https://hdl.handle.net/20.500.11811/8064> (cit. on p. 11).
- [DT92] D. Drechsel and L. Tiator, *Threshold pion photoproduction on nucleons*,
J. Phys. G **18** (1992) 449.
- [Bar+18] M. Bartelmann et al., *Theoretische Physik 3 — Quantenmechanik*, 2018,
ISBN: 978-3-662-56071-6.
- [Wal] D. Walther, *Crystal Barrel, A 4π photon spectrometer*,
URL: <https://www.cb.uni-bonn.de> (visited on 27/09/2021).
- [Urb17] M. Urban, *Design eines neuen Lichtpulsersystems sowie Aufbau und Inbetriebnahme der neuen APD Auslese für das Crystal-Barrel-Kalorimeter*,
PhD thesis: Rheinische Friedrich-Wilhelms-Universität Bonn, 2017.
- [Afz22] F. Afzal, *Private communication*, 2022 (cit. on p. 11).

List of Figures

1.1	Running coupling of QCD. The colored data points represent different methods to obtain a value for α_s . For more details it may be referred to [Zyl+20].	2
1.2	Calculated nucleon (isospin $I = 1/2$) resonances compared to measurements. Left in each column are the calculations [LMP01], the middle shows the measurements and PDG rating [Zyl+20]	3
1.3	FEYNMAN diagram for the s-channel photoproduction of pseudoscalar mesons, adapted from [Afz19]	4
2.1	[Wal]	7
2.2	[Wal]	8
2.3	[Wal]	8
2.4	D. WALTHER in [Urb17]	9
2.5	[Wal]	9
2.6	[Wal]	10
3.1	Distribution of event classes in $\eta' \rightarrow \gamma\gamma$ production	12
3.2	Time information of all final state particles and the beam photon for 3PED η' production	13
3.3	Reaction time t_r for 3PED η' production	14

List of Tables

1.1	Summary of the particles of the SM	1
1.2	Allowed quantum numbers for the intermediate resonance state N^*/Δ^*	4
3.1	The five most probable decay modes of the η' meson. The most probable further decay with according branching ratio is shown in brackets.[Zyl+20]	11