

Recent Polarization Observable Results in η - and η' -photoproduction off the proton

Master thesis for the CBELSA/TAPS collaboration

JAKOB KRAUSE

✉ krause@hiskp.uni-bonn.de | 🌐 krausejm

Supervisor: JUN. PROF. DR. ANNIKA THIEL

✉ thiel@hiskp.uni-bonn.de


30th March 2022

Setting the scene

The Standard Model of Particle Physics


- ▶ matter consists of 12 (anti-) *fermions*
- ▶ quarks interact via *strong interaction*
- ▶ form bound states: mesons ($q\bar{q}$) and baryons (qqq)

baryon spectroscopy (photoproduction) gives insight in strong interaction



Setting the scene

Observe resonances R^* in the cross sections $\sigma(\gamma p \rightarrow R^* \rightarrow pM)$



Total cross section $\sigma(\gamma p \rightarrow R^* \rightarrow p\pi^0)$ [Wunderlich et al. 2017]

→goal: (help to) identify contributing resonances as strong bound states!

1. Theoretical Basics
2. Experimental Setup
3. Preliminary results
4. Conclusion

1. Theoretical Basics

2. Experimental Setup

3. Preliminary results

4. Conclusion

Theoretical Basics

Unpolarized differential cross section

$$\frac{d\sigma}{d\Omega} = \frac{1}{4}\rho \sum_{\text{spins}} |\langle f | \mathcal{F} | i \rangle|^2,$$

where

$$\mathcal{F} = i(\vec{\sigma} \cdot \vec{\epsilon})F_1 + (\vec{\sigma} \cdot \hat{q})(\vec{\sigma} \cdot (\hat{k} \times \vec{\epsilon}))F_2 + i(\vec{\sigma} \cdot \hat{k})(\hat{q} \cdot \vec{\epsilon})F_3 + i(\vec{\sigma} \cdot \hat{q})(\hat{q} \cdot \vec{\epsilon})F_4$$

F_i : complex CGLN Amplitudes

[Chew et al. 1957]

$\frac{d\sigma}{d\Omega} \in \mathbb{R}$, not sufficient to determine \mathcal{F} unambiguously
→ Polarization Observables

Theoretical Basics


- ▶ resonances are broad, overlapping, require sophisticated analysis (PWA)
- ▶ constraints feeding the analysis can be derived from Polarization observables

Beam asymmetry Σ

$$\frac{d\sigma}{d\Omega}(\varphi) = \frac{d\sigma}{d\Omega_0} \left[1 - p_{\gamma}^{\text{lin}} \Sigma \cos(2\varphi) \right]$$

polarization angle φ , polarization degree p_{γ}^{lin}

[Sandorfi et al. 2011]




Definition of the polarization angle

1. Theoretical Basics
2. Experimental Setup
3. Preliminary results
4. Conclusion

Experimental setup

- ▶ generate photon beam from accelerated electrons via bremsstrahlung, with $E_\gamma \leq 3.2$ GeV
- ▶ photon beam impinges on liquid butanol target: $\gamma p \rightarrow pM \rightarrow pX$
- ▶ measure decay products X of different final states: $M = \pi^0/\eta/\eta'/\dots$



Overview of the experimental area, adapted from [Walther 2022]


1. Theoretical Basics
2. Experimental Setup
3. Preliminary results
4. Conclusion

Preliminary results

Event selection (η')

analysis performed in 2×6 bins of $(E_\gamma, \cos \theta_{\eta'}^{\text{CMS}})$, $E_\gamma \in [1400, 1800]$ MeV

- ▶ 3 detector hits, 2 uncharged, 1 charged
- ▶ coincident detector hits
- ▶ kinematic cuts derived from energy-momentum conservation
 $p_\gamma + p_p = p'_p + p_{\eta'}$
- ▶ additional cuts to reduce background contributions



Preliminary results (η')

Method

- ▶ measure in 2 distinct orthogonal polarization settings \perp, \parallel
- ▶ χ^2 -fit to event yield asymmetries

$$A(E_\gamma, \theta, \phi) = \frac{N^\perp(E_\gamma, \theta, \phi) - N^\parallel(E_\gamma, \theta, \phi)}{p_\gamma^\parallel N^\perp(E_\gamma, \theta, \phi) + p_\gamma^\perp N^\parallel(E_\gamma, \theta, \phi)} = \Sigma(E_\gamma, \theta) \cos\left(2\left(\alpha^\parallel - \phi\right)\right)$$

plot of event yield asymmetries and sigma

Preliminary results (η)

Event selection (η)

analysis performed in 11x12 bins of $(E_\gamma, \cos \theta)$ by FARAH AFZAL *cite Farahs Paper*

Method

- ▶ fit to event yield asymmetries using BAYESIAN inference
- ▶ $p(\theta|y) \propto \mathcal{L}(\theta, y) \cdot \pi(\theta)$, obtained using MARKOV-Chain-MONTE-CARLO (MCMC)

plot of eta sigma

1. Theoretical Basics
2. Experimental Setup
3. Preliminary results
4. Conclusion