

Run 4

Contents

Set up simulation	1
Copy fault model	1
Set up injection history and resulting stressing file	2
Run simulation	4
Results	4
Magnitude-time plot	4
Hypocenter-well distance	5
Pore-fluid pressure at failure	7
Estimation of Omori p -value and aftershock fraction n	9
Aftershock fraction ala <i>Zaliapin and Ben-Zion</i> [2016]	12
Moment due to ΔP	13
Summary	15
To do	16
References	16

```
require(MASS)
require(rstan)
options(mc.cores = parallel::detectCores())
rstan_options(auto_write = TRUE)
```

Set up simulation

Difference from Run3 is slightly lower initial stress on the main fault. Same fault system and injection history going back to Run2, so can reuse fault system and external stressing file.

Copy fault model

Make a simple fault model consisting of a single, vertical, planar strikeslip fault, and then add random small faults around it.

```
file.copy(file.path("../Run2", "test.addRandom.10.flt"), ".")
flt2 = readFault("test.addRandom.10.flt")
```

Plot of resulting fault system

```
# injection point that we will use below
xi = matrix(c(500, 0, -3.5e3), 1, 3) # well location: 500 m to east of center pt. of main fault
```

```
p = plotFault3d(flt2, theta=45, phi=40, xlim=c(-4,4), ylim=c(-6,6), zlim=c(-7,-1), lwd=0.5)
lines(trans3d(c(xi[1,1], xi[1,1])/1e3, c(xi[1,2], xi[1,2])/1e3, c(0, xi[1,3])/1e3, p), lwd=3)
```



Set up injection history and resulting stressing file

Make a simple injection history and RSQSim external stressing file from it. Can just copy this over from Run2. Or, actually, to save disk space (current test.externalStress.txt is 450 MB), should just point to Run2's copy.

```
yr = 365.25*86400

injHist = makeInjectionHistory(xi=xi) # use defaults for all hydrologic params

#t = findSampleTimes(flt2, injHist, tmax=100*yr, eps=0.05, minh=1000, maxErr=100)
#writePoreFluidStressingHistoryFile(flt2, injHist, t, "test.externalStress.txt")
#file.copy(file.path("../", "Run2", "test.externalStress.txt"), ".")

s = readExternalStressingHistoryFile(file.path("../", "Run002", "test.externalStress.txt"), flt2)
```

Plots of pressure histories


The closest element, the farthest element, and the median distance element.

```
dwell = elementWellDist(flt2, injHist)
ord = order(dwell)
plot(s$t/yr, -s$sigma[ord[1],], typ='l', xlab="Time (yrs)",
     ylab="Pore-fluid pressure perturbation (MPa)")
```

```

lines(s$t/yr, -s$sigma[ord[flt2$np],], col="red2")
lines(s$t/yr, -s$sigma[ord[round(flt2$np/2)],], col="orange")

```



Initial stresses

```

set.seed(234)
tau00.main = 62
tau00.random = 63
sdtau.main = 0
sdtau.random = 1
main = which(flt2$segName == "NA")
random = which(flt2$segName != "NA")
tau0 = rep(NA, flt2$np)
tau0[main] = rnorm(length(main), tau00.main, sdtau.main)
tau0[random] = rnorm(length(random), tau00.random, sdtau.random)
write(tau0, "initTau.txt", ncol=1)

```

Run simulation

Copy Run3's parameter file and make adjustments.


```
eqstmp = readEqs("../Run003/eqs.Run3.out")
params = eqstmp$params
rm(eqstmp)
params$outFnameInfix = "Run4"
params$extStressRateFname = "../Run002/test.externalStress.txt"
params$initTauFname = "initTau.txt"
writeRSQSimInputFile(params, paste("test.", params$outFnameInfix, ".in", sep=""))
unlink(paste("eqs.", params$outFnameInfix, ".out", sep=""))
system2("mpirun", args=c("-np 4 ~/RSQSim.github.master/runRSQSim",
                          paste("test.", params$outFnameInfix, ".in", sep="")),
        stdout=paste("test.", params$outFnameInfix, ".std", sep=""),
        stderr=paste("test.", params$outFnameInfix, ".std", sep=""))
```

Results

```
eqs = readEqs(paste("eqs.", params$outFnameInfix, ".out", sep=""))
if (max(eqs$t0) > max(s$t)) {
  eqMax = min(which(eqs$t0 > max(s$t))) - 1
  eqs = readEqs(paste("eqs.", params$outFnameInfix, ".out", sep=""), eqMax = eqMax)
}
```


Magnitude-time plot

```
plot(eqs$t0yr, eqs$M, typ='h', xlab="Origin time (yrs)", ylab="Magnitude")
```



Hypocenter-well distance


```
d = hypoWellDist(eqs, injHist)
plot(eqs$t0yr, d/1e3, ylim=c(0,max(d/1e3)),
      xlab="Origin time (yr)", ylab="Distance to injection point (km)")
abline(v=eqs$t0yr[eqs$M>5], col="red", lty="dashed")
```



The evolution of the distance of the induced events from the injection point with time looks roughly like the expected \sqrt{t} (until the M5+). It is not clear what the exact relationship between the coefficient of \sqrt{t} and the hydraulic parameters should be. *Shapiro et al.* [1997] advocate for $x = \sqrt{4\pi\kappa t}$, where κ is the diffusivity. The argument of the erfc in our pore-fluid pressure expression is equal to one at $x = \sqrt{4\kappa t}$, but there is also a $|x|^{-1}$ factor in front of the erfc. Adding in the least-squares fit to the pre-M5 events:

```
plot(eqs$t0yr, d/1e3, ylim=c(0,max(d/1e3)),
      xlab="Origin time (yr)", ylab="Distance to injection point (km)")
abline(v=eqs$t0yr[eqs$M>5], col="red", lty="dashed")

if (max(eqs$M) < 5) use=1:length(eqs$t0) else use = 1:(which(eqs$M>5)-1)
k = sum(d[use]*sqrt(eqs$t0[use])) / sum(eqs$t0[use]) # in m/s^{1/2}
kplot = k * (1/1e3) * sqrt(86400*365.25) # in km/yr^{1/2}
tplot = seq(0, max(eqs$t0yr), length=1000)
lines(tplot, kplot*sqrt(tplot))
```



The \sqrt{t} is not a great fit: event times are systematically earlier at near distances and later at farther distances than the best fit \sqrt{t} curve. I would attribute this to the $|x|^{-1}$ factor in the pore-fluid pressure expression. It could also be due to RSF evolution effects. If we ignore that and ask what value of κ is implied by the *Shapiro et al.* [1997] expression or just setting the argument of the erfc to 1, we get 2.7×10^{-5} and 8.4×10^{-5} m^2/s , respectively. These are both about two orders of magnitude lower than the actual diffusivity used in generating the pore-fluid pressures ($0.008 \text{ m}^2/\text{s}$). Note that this difference here between the actual diffusivity and that estimated by assuming $x = \sqrt{4\pi\kappa t}$ is not necessarily relevant to the data in *Shapiro et al.* [1997]. In that paper the diffusivity is much higher and the time scales are much shorter (a few days), so that RSF evolution effects may be less important.

Elizabeth mentions that some papers of Talwani estimates the diffusivity by fitting curves of the form $x = \sqrt{\kappa t}$. Also said that might have been for fluid pressures diffusing in 2-d (*e.g.* if a fault is a high permeability conduit) instead of in the 3-d bulk. If we ignore the 2-d/3-d issue and just estimate κ from curves of that form, we get $3.3 \times 10^{-4} \text{ m}^2/\text{s}$, still more than an order of magnitude less than the actual value.

Why are these estimates so much lower here than in the uniform pre-stress cases?

Pore-fluid pressure at failure

As an indication of how important RSF evolution effects may be, let's look at the pore-fluid pressure one each hypocentral element just before it failed. For friction with a fixed failure stress (*e.g.* slip-weakening) and uniform initial stresses as we have here, this would be expected to be the same for all events as long as the stress transfer between the elements is negligible (which it will be in this fault system everywhere except

Note that the instantaneous pore-fluid pressure increment that would cause immediate failure is given by that P which solves

For this simulation, this comes out to about 26.9 MPa.

```
itfail = findInterval(eqs$t0, s$t)
itfail[itfail >= length(s$t)] = length(s$t) - 1
pFail = s$sigma[cbind(eqs$patch, itfail)] +
  (s$sigma[cbind(eqs$patch, itfail+1)] - s$sigma[cbind(eqs$patch, itfail)]) *
  (eqs$t0 - s$t[itfail])/(s$t[itfail+1] - s$t[itfail])
pFail = -pFail
plot(eqs$t0yr[use], pFail[use], xlab="Origin time (yr)",
      ylab="Pore-fluid pressure at failure (MPa)")
```



here because of the variable pre-stress.

So, for now, it is an open question how much of the departure from \sqrt{t} behavior is due to the $|x|^{-1}$ and how much due to RSF evolution effects.

Estimation of Omori p -value and aftershock fraction n

See `templateFamilies.Rmd|pdf`


```
use = 1:length(eqs$t0)
alpha = 0.32 # so 68% CIs
c = 90 # seconds, needs to match dSS2007.krd.fixedloglambdac.stan
st = do.stan(eqs, use=use, alpha=alpha, c=c)
```

Plotting function definition

The functions to make plots like in `templateFamilies.pdf`; modified from there. Not printing them out because they take a lot of page space.

Actually make the plots

```
mkFourPanelPlot(eqs, use, st$bfmi, st$n_div, st$rhat, st$n_eff,
                 st$f.stan,
                 st$p.stan.mode, st$pCI.stan, st$loglambdac.stan.mode,
                 st$n.stan.mode, st$nCI.stan)
```




This estimates the aftershock fraction to be 0.05. Just given the data asked for my opinion, I probably would have counted the 0 events that occurred in the couple of years after the M5+ as aftershocks and said the aftershock fraction was $0/22 = 0$.

Leaving out very late events

Perhaps those last 0 events way off by themselves are biasing things? What happens if we leave them out?

```
use2 = 1:(length(eqs$t0)-nout)
st2 = do.stan(eqs, use=use2, alpha=alpha, c=c)
```

```
mkFourPanelPlot(eqs, use2, st2$bfmi, st2$n_div, st2$rhat, st2$n_eff,
  st2$f.stan,
  st2$p.stan.mode, st2$pCI.stan, st2$loglambdac.stan.mode,
  st2$n.stan.mode, st2$nCI.stan)
```



Aftershock fraction didn't change much: 0.04.

Note that the estimate of *Hainzl et al.* [2006] would give 0.79 and 0.79, when using all events and all the but the last 0 events, respectively.


Both the estimated aftershock fraction from fitting the *Saichev and Sornette* [2007] functional form and the C_v are very affected by these outliers. If we used the 1-norm version of the C_v , call it C_{v1} , $C_{v1} = \text{smad}(dt) / \text{median}(dt)$, it would be essentially completely unaffected: including the last 0 events gives 0.9775074 and excluding them gives 0.9775074.

Aftershock fraction ala *Zaliapin and Ben-Zion* [2016]

```
sderrh = 10
z = zaliapin(eqs, use=use, sderrh=sderrh, est.se=TRUE)
z2 = zaliapin(eqs, use=use2, sderrh=sderrh)
```


Using all events

```
source("../mkZaliapinPlots.R")
mkZaliapinPlots(z)
```





Excluding late events


```
mkZaliapinPlots(z2)
```



Moment due to ΔP

```
source("../plotMomentDueToDeltaP.R")
plotMomentDueToDeltaP(eqs, dwell)
```





Summary

```
source("../writeResults.R")
writeResults(paste(params$outFnameInfix, ".results.txt", sep=""),
            eqs, nout, nas, st, st2, z, z2)
results = readResults(paste(params$outFnameInfix, ".results.txt", sep=""))
```

- Full-main-fault event at NA yrs
- Number of events
 - total: 22
 - before full-main-fault event: NA
 - after full-main-fault event: NA
- Estimated aftershock fractions
 - naive count: 0
 - SS2007
 - * using all events: 0.05
 - * excluding last 0 events: 0.04
 - H2006
 - * using all events: 0.79
 - * excluding last 0 events: 0.79
 - ZB2016
 - * using all events: 0.24

* excluding last 0 events: 0.24

To do

- how to get more realistic time-scales and pore-pressures?
- different uniform stresses on main fault
- variable stresses on main fault
- different forms / amplitudes of variable stresses on random faults
- denser random faults

References

- Hainzl, S., F. Scherbaum, and C. Beauval (2006), Estimating background activity based on interevent-time distribution, *Bulletin of the Seismological Society of America*, *96*(1), 313–320.
- Saichev, A., and D. Sornette (2007), Theory of earthquake recurrence times, *Journal of Geophysical Research: Solid Earth*, *112*(B4).
- Shapiro, S. A., E. Huenges, and G. Borm (1997), Estimating the crust permeability from fluid-injection-induced seismic emission at the ktb site, *Geophysical Journal International*, *131*(2), F15–F18.
- Zaliapin, I., and Y. Ben-Zion (2016), Discriminating characteristics of tectonic and human-induced seismicity, *Bulletin of the Seismological Society of America*, *106*(3), 846–859.