

Notation \leadsto See course book p. 9-12

p : # of predictors in our model:

Matrix


$$X_{n \times p} =$$

$$\begin{bmatrix} x_{11} & \dots & x_{1p} \\ x_{21} & & x_{2p} \\ \vdots & & \vdots \\ x_{n1} & \dots & x_{np} \end{bmatrix}$$

x_1^T all meas. for unit 1
 n units
 x_p

measurements of variable 1
for all units

see p. 10



Reducible vs. irreducible error

Assume f and x are fixed

$$\begin{aligned} E[(Y - \hat{Y})^2] &= E[(f(x) + \varepsilon - \hat{f}(x))^2] \\ &= E[(f(x) - \hat{f}(x))^2] + E[\varepsilon^2] + 2 E[(f(x) - \hat{f}(x)) \cdot \varepsilon] \\ &\quad \underbrace{\hspace{10em}}_{\text{fixed}} \quad \xrightarrow{\text{var}(\varepsilon)} \\ &\quad \quad \quad = 0 \quad (E(\varepsilon) = 0) \end{aligned}$$

$$= (f(x) - \hat{f}(x))^2 + \text{var}(\varepsilon)$$

Bias-variance trade-off

Expected test MSE at x_0

$$\begin{aligned} E[(y_0 - \hat{f}(x_0))^2] &= E[(f(x_0) + \varepsilon - \hat{f}(x_0))^2] \\ &= \underbrace{E[f(x_0)^2]}_{\text{var}(\hat{f}(x_0)) + E[f(x_0)]^2} + E[\varepsilon^2] + \underbrace{E[(\hat{f}(x_0))^2]}_{\text{var}(\hat{f}(x_0)) + E[f(x_0)]^2} \end{aligned}$$

$$\begin{aligned}
& -2E[f(x_0)\hat{f}(x_0)] + \underbrace{2E[\varepsilon \cdot f(x_0)]}_{=0} - \underbrace{2E[\varepsilon \cdot \hat{f}(x_0)]}_{=0} \\
& = \underbrace{f(x_0)^2}_{\text{irreducible error}} + \text{Var}(\varepsilon) + \text{Var}(\hat{f}(x_0)) + \underbrace{E[\hat{f}(x_0)]^2}_{\text{Squared bias}} \\
& \quad - \underbrace{2E[f(x_0)\hat{f}(x_0)]}_{\text{Variance}}
\end{aligned}$$

~~~~~