

# Rounds Codex – Cardiac Arrest (ACLS) Quiz

**1. A patient is unresponsive, not breathing normally, and has no definite carotid pulse within 10 seconds. What is the next step?**

- A. Start high-quality CPR immediately
- B. Obtain a 12-lead ECG
- C. Check pupils
- D. Wait for monitor
- E. Give epinephrine

**Answer:** A — Immediate CPR is the priority.

**2. Recommended adult chest compression rate is:**

- A. 60–80/min
- B. 80–100/min
- C. 100–120/min
- D. 120–140/min
- E. As fast as possible

**Answer:** C — Maintain 100–120/min.

**3. Which rhythm is shockable?**

- A. Asystole
- B. PEA
- C. VF
- D. Sinus bradycardia
- E. AV block

**Answer:** C — VF and pulseless VT are shockable.

**4. Primary purpose of high-quality CPR is:**

- A. Restore normal BP
- B. Maintain coronary and cerebral perfusion
- C. Terminate VF
- D. Treat acidosis
- E. Reverse MI

**Answer:** B — CPR buys time by maintaining perfusion.

**5. Which is NOT one of the Hs and Ts?**

- A. Hypovolemia
- B. Hypoxia
- C. Hyperglycemia
- D. Tamponade
- E. Toxins

**Answer:** C — Hyperglycemia is not an H or T.

**6. During adult cardiac arrest, epinephrine dose is:**

- A. 0.5 mg q10 min
- B. 1 mg IV/IO every 3–5 min
- C. 2 mg once
- D. 5 mg q5 min
- E. 300 mg once

**Answer:** B — Standard ACLS dose.

**7. With an advanced airway in place:**

- A. Pause compressions for breaths
- B. 30:2 continues
- C. Continuous compressions with 1 breath every 6 sec
- D. 1 breath every 3 sec
- E. Stop ventilation

**Answer:** C — Continuous compressions plus 10 breaths/min.

**8. An abrupt sustained rise in ETCO<sub>2</sub> during CPR most likely indicates:**

- A. Hyperventilation
- B. ROSC
- C. Pulmonary embolism
- D. Asystole
- E. Poor compressions

**Answer:** B — Sudden ETCO<sub>2</sub> rise suggests ROSC.

**9. After ROSC, an appropriate MAP target is:**

- A. 40 mmHg
- B. 50 mmHg
- C. ≥65 mmHg
- D. ≥90 mmHg
- E. No target

**Answer:** C — Support organ perfusion.

**10. Which statement best summarizes ACLS priorities?**

- A. Medications save most lives
- B. CPR and early defibrillation are the highest priorities
- C. Troponin guides CPR
- D. CT first
- E. Shock all rhythms

**Answer:** B — Early CPR and defibrillation have greatest impact.