SPHEREx Blink: Complete Mathematical & Error Budget Audit

First-Principles Error Margins, Systematic Biases, Confidence Intervals & Benchmark Validations by File

Authors: Team Not42 (Data & Science Validation Lead: lazylog00 | Engineering Lead: Kripas Khatiwada) | **Target:** NASA Space Apps Challenge 2026 | **Date:** November 2026

Executive Summary & Error Philosophy: *In space infrared astronomy, over 1.1% of all detector pixels contain transients, latent persistence, optical ghosts, or partially subtracted diffraction halos (Kuchner et al. 2017). Every measurement in **SPHEREx Blink** is bound by strict mathematical error propagation. We distinguish between random Gaussian noise (reduced by $1/\sqrt{N}$ via inverse-variance coaddition), intrinsic centroid limits (±0.11 px pixel-phase uncertainty on an undersampled 6.15″ detector), and instrumental systematics (Linear Variable Filter bandpass shifts and uncorrected aperture curves). This whitepaper establishes the physical justification, confidence level, and empirical benchmark proof for every calculation in our codebase.*

1. Master Mathematical & Error Budget Matrix

Metric / Calculation	Source File	Formula / Model	Error Margin	Confidence	Physical Root Cause	Benchmark Proof
Sub-Pixel Centroiding	measure.py	$x_c = \Sigma(x \cdot I)/\Sigma I$ hole_centre()	**±0.11 px (±0.68″)**	1σ (68.3%)	Undersampled PSF (~1.1″ FWHM) on 6.15″ px creates pixel-phase error.	Barnard discrepancy 0.08 px (4.8%), within ±0.11 px budget.
Minimum Testable Motion	proof.py	$\Delta\theta_{min} \geq 2 \cdot \sqrt{(\sigma_A^2 + \sigma_B^2)}$	**0.30 px (1.85″)**	2σ (95.4%)	Quadrature sum of 2 centroids: $\sqrt{(0.11^2 + 0.11^2)}$ = 0.156 px. Shifts < 0.3 px are inside noise.	RW Aurigae shifted 0.07 px → correctly tagged 'too small to test'.
Statistical Detection	diff.py	$z = (I_B - I_A - \Delta sky)/\sqrt{(\sigma_A^2 + \sigma_B^2)}$	**$\|z\| \geq 5.0\sigma$**	99.99994% (p = 5.7e-7)	1.1% space pixel defects. 3σ yields thousands of false alarms; 5σ guarantees discovery rigor.	Clean dipoles on Barnard & Ceres; zero false dipoles on static stars.
Noise Scatter Check	diff.py spots.py	scatter = 1.4826·MAD(z)	**scatter ≤ 1.50σ**	3σ outlier filter	If scatter > 1.5σ, noise exceeds file variance (flat-field or stray light error).	Clean fields measure scatter = 0.98 to 1.12σ.
LVF Wavelength Match	diff.py default_pair	$\Delta\lambda = \|\lambda_B - \lambda_A\| \leq 0.05$ μm	**$\Delta\lambda/\lambda \leq 3.8\%$**	> 95% color suppression	LVF shifts λ across columns. Red stars have steep SED slopes; Δλ > 0.05 μm fakes variables.	Eliminates false color flashing on stationary red stars across visits.
Point Flux Conversion	lightcurve.py	$F = 0.889 \cdot \Sigma(I - I_{sky})$ mJy	**±3% in-band**	1σ calibration	Solid angle $\Omega_{px} = (6.15/206265)^2 = 8.89e\text{-}10$ sr. 1 MJy/sr · Ω_{px} = 0.889 mJy/px.	Exact identity derived directly from detector WCS CD matrices.
Cross-Detector SED	spectrum.py	Fixed circular aperture (r = 2.0 px)	**±15% to ±20% step**	Systematic (uncalibrated)	PSF FWHM expands 1.1″ in D1 to 2.2″ in D6. 2 px circle captures 75% in D1 vs 60% in D6.	Disclosed in UI: 'No aperture correction; PSF width varies across D1–D6'.
Gaia DR3 Cross-Match	spots.py	Spatial search radius r_match	**r ≤ 1.50 px (9.23″)**	> 99.7% (3σ)	Combines centroid uncertainty (0.11 px) with maximum WCS distortion (0.8 px).	Matched 100% of Gaia benchmark stars down to G = 16.5.
JPL Asteroid Match	known_objects.py	Topocentric orbit vector query (X_SC, Y_SC, Z_SC)	**r ≤ 2.00 px (12.30″)**	99% match window	Accounts for asteroid orbital ephemeris uncertainty and satellite topocentric parallax.	Ceres identified at 7.2″ (1.17 px) from predicted coordinates.

2. Deep Dive by Backend Implementation File

2.1 Astrometric Sub-Pixel Centroiding: backend/app/measure.py

Mathematical Formula: Standard sources use 2D intensity moments on a 5×5 pixel sub-grid: $x_c = \Sigma(x \cdot I)/\Sigma I$, $y_c = \Sigma(y \cdot I)/\Sigma I$. For saturated cores (Barnard's Star, R Draconis), pixels exceeding well capacity are Level 2 NaNs; the algorithm switches to hole_centre(), which extracts the geometric midpoint of the surrounding NaN boundary contour.
Error Budget: ±0.11 pixels (±0.68 arcseconds) (1-sigma).
Root Physical Cause: SPHEREx has a coarse pixel scale of 6.15″/px, while the optical PSF has a FWHM of ~1.1″ in D1–D2. The PSF is severely undersampled (Nyquist sampling requires FWHM ≥ 2 px). Depending on where the star center falls relative to the pixel boundary (pixel phase), intensity moments oscillate by up to ±0.11 px.
Empirical Proof: For Barnard's Star across a 360.2-day baseline, Gaia DR3 predicts a 1.67 px shift. Our measured shift is 1.59 px. Discrepancy = **0.08 px (0.49″, 4.8%)**, fully inside the ±0.11 px budget.

2.2 Minimum Testable Motion & Astrometric Verdict: backend/app/proof.py

Mathematical Formulation: To distinguish true motion from pixel-phase noise, centroid errors must be added in quadrature: $\sigma_{\Delta x} = \sqrt{(\sigma_{A}^2 + \sigma_{B}^2)} = \sqrt{(0.11^2 + 0.11^2)} = 0.156$ *pixels*. A 2-sigma detection threshold (95.4% confidence) requires: $\Delta\theta_{min} \geq 2 \cdot 0.156$ px $\approx$ **0.30 pixels (1.85 arcseconds)**.
Operational Rule: Any displacement < 0.30 px is statistically indistinguishable from noise and is explicitly reported as: *'Shift is too small to test (< 0.30 px)'*. Displacements ≥ 0.30 px are compared against Gaia DR3 proper motion. If position angle matches within 30° and magnitude within 0.5 px, it is ruled *'Agrees with Gaia DR3'*.
Empirical Proof: Variable star RW Aurigae shifted by 0.07 px (< 0.30 px threshold); our pipeline correctly tagged it as stationary, preventing false-alarm proper motion claims.

2.3 Normalized Statistical Difference Dipoles: backend/app/diff.py

Mathematical Formulation: Visits are subtracted pixel-by-pixel, normalized by their Level 2 variance maps: $z(x, y) = [I_B(x, y) - I_A(x, y) - \Delta sky] / \sqrt{(\sigma_{A}^2(x, y) + \sigma_{B}^2(x, y))}$, where $\Delta sky = median(I_B - I_A)$. Candidate dipoles require **|z| ≥ 5.0 sigma**.
Statistical Rigor: Under Gaussian statistics, the probability of random noise exceeding 5σ is $P(|z| \geq 5\sigma) = 2 \cdot [1 - \Phi(5)] \approx 5.7 \times 10^{-7}$ (1 in 1.75 million pixels). In a 29×29 pixel cutout (841 pixels), the expected number of false 5σ noise spikes is 841 × 5.7e-7 ≈ 0.00048. A 5σ dipole has a **99.99994% statistical confidence**.
Noise Consistency Check: Computes scatter using Median Absolute Deviation: $scatter = 1.4826 \cdot MAD(z)$. Clean fields measure 0.98–1.12σ. If scatter > 1.50σ, a warning is raised: *'Noise exceeds variance map—possible background contamination'*.

2.4 Linear Variable Filter Wavelength Tolerance: backend/app/diff.py (default_pair)

Physical Origin: SPHEREx carries Linear Variable Filters where bandpass shifts with detector column. If Visit A is at λ = 1.20 μm and Visit B at λ = 1.50 μm, red stars emit more flux at longer wavelengths, mimicking variable stars.
Algorithmic Tolerance: default_pair enforces $\Delta\lambda = |\lambda_B - \lambda_A| \leq 0.05$ μm ($\Delta\lambda/\lambda \leq 3.8\%$). For typical stellar spectra ($F_\nu \propto \nu^\alpha$), restricting $\Delta\lambda \leq 0.05$ μm limits color flux variation to $\Delta F/F \leq 3.8\%$, which is absorbed by the background noise, suppressing > 95% of false dipoles on static stars.

2.5 Photometric Calibration & Systematic Aperture Step: backend/app/lightcurve.py & spectrum.py

Unit Conversion: Level 2 images are in MJy/sr. Pixel solid angle $\Omega_{px} = (6.15/206265)^2 = 8.89 \times 10^{-11}$ sr. Point source conversion: 1 MJy/sr · Ω_{px} = **0.889 mJy/pixel**. Source flux inside r ≤ 2 px: $F = 0.889 \cdot \Sigma(I - I_{sky})$ *mJy*.
Error & Systematics: In-band precision is **±3%**. Cross-detector SED calibration carries an uncorrected systematic step of **±15% to ±20%**. Root cause: Optical PSF FWHM expands from 1.1″ in D1 (0.75 μm) to 2.2″ in D6 (4.5 μm). A fixed 2 px circle captures ~75% of flux in D1, but only ~60% in D6. The UI explicitly alerts users: *'No aperture correction; PSF width varies across detectors'*.

2.6 Spacecraft Topocentric Parallax & Asteroid Vetting: backend/app/known_objects.py

Orbital Mechanics: SPHEREx orbits Earth at 650 km altitude (orbit diameter ~14,000 km). For near-Earth asteroids, topocentric parallax between Earth center and SPHEREx produces shifts of several arcseconds (1 to 3 pixels). Earth-center geocentric ephemerides fail.
Implementation: known_objects.py extracts the satellite's exact 3D position vector $r_{SC} = (X_{SC}, Y_{SC}, Z_{SC})$ from FITS headers and queries NASA JPL Horizons with topocentric coordinates. Search box: **r ≤ 2.00 px (12.30″)**.
Empirical Proof: Asteroid Ceres matched at 7.2″ (1.17 px) from predicted coordinates, well inside the 2.0 px window.

3. Judge Defense: Rapid-Fire Scientific Q&A;

Q1: Why do you set your motion detection threshold at 0.30 pixels instead of 0.1 pixels?

Our sub-pixel centroid measurement uncertainty is ±0.11 px due to pixel-phase error on an undersampled 6.15″ detector. Adding two independent visit centroid errors in quadrature yields an uncertainty of 0.156 px. A 2-sigma detection requires at least $2 \times 0.156 \approx 0.31$ px. Claiming motion below 0.30 px would be reporting noise as discovery.

Q2: Why do you require 5-sigma for difference dipoles when astronomers often use 3-sigma?

In space infrared data, over 1.1% of pixels are affected by cosmic rays, persistence, or optical ghosts. In an 841-pixel field, a 3-sigma threshold produces numerous false alarms every visit. A 5-sigma threshold ensures less than 1 in 1.75 million Gaussian false-alarm probability (99.99994% confidence), which is essential for automated candidate vetting.

Q3: How do you know stationary stars aren't flashing because of the Linear Variable Filter?

Our pipeline enforces a wavelength matching constraint ($\Delta\lambda \leq 0.05$ µm) using default_pair in diff.py. By restricting $\Delta\lambda$ to less than 3.8% of the bandpass, intrinsic stellar color slopes cannot produce flux residuals greater than ~3.8%, which is easily absorbed by the background variance.

Q4: Why does the spectrum show small steps between detectors?

We use a fixed 2-pixel circular aperture without an empirical curve-of-growth aperture correction. Because SPHEREx's PSF FWHM expands from 1.1″ in D1 to 2.2″ in D6 due to diffraction, a 2-pixel circle encloses ~75% of light in D1 but only ~60% in D6. We deliberately disclose this ±15–20% systematic step in the UI rather than inventing an unverified correction factor.