SPHEREx Blink: First-Principles Astrometric Vetting & Discovery Architecture

Executive Scientific Audit, Error Budgets, Physical Traps, and the Candidate Dossier Protocol

Target: NASA International Space Apps Challenge 2026 (SPHEREx: Planet X and Friends)
Authors: Team Not42 (Data & Science Validation Lead: lazylog00 | Engineering Lead: Kripas Khatiwada)
Verified Sources: SPHEREx Explanatory Supplement v2.0, IPAC IRSA TAP, and 10 Literature Papers (researchpdf/)

Executive Summary

This document establishes the scientific ground truth, mathematical foundations, error margins, and candidate vetting protocols for **SPHEREx Blink**. It synthesizes literature findings across 10 peer-reviewed papers with real observational analyses of the 5 official NASA Space Apps Challenge benchmark targets: Barnard's Star, Epsilon Indi B, WISE 0855, RW Aurigae, and R Draconis.

Our primary objective is to bridge the documented gap in the SPHEREx mission: **while the satellite maps the entire sky every 6 months to identify time-domain transients, no official NASA/IRSA tool compares visits across epochs.** We present an open-source, mathematically audited pipeline that transforms raw space telescope cutouts into calibrated, wavelength-matched blinking flipbooks, enforces rigorous artifact vetoes, cross-matches Gaia DR3 proper motions and NASA JPL Horizons orbital telemetry, and produces standardized candidate discovery dossiers with zero hype.

1. The Core Scientific Problem & Mission Gap

1.1 The Official Mission Reality

The primary SPHEREx mission paper (*Bock et al. 2026*, Sections 2.4 and 3.4.6) defines time-domain science as a core deliverable: *'In the all-sky survey, SPHEREx measurements can serve as a reference every 6 months to identify time-variable sources.'* However, Table 5 of *Bock et al. 2026* catalogues all official NASA/IPAC Infrared Science Archive (IRSA) tools: Spectrophotometry, Cutout Service, Source Discovery, Custom Mosaic, and Spectral Cube Cutout. **None of these official tools compare multi-epoch visits.** Furthermore, Table 4 confirms that official NASA legacy catalogs of brown dwarfs and solar system objects will not be released until **May 2028**.

1.2 Why Blind Machine Learning Fails Alone

In wide-field space surveys, automated computer differencing produces massive false-alarm catalogs. *Kuchner et al. 2017* (Backyard Worlds: Planet 9, Section 2) documented that in space infrared data, **over 1.1% of all pixels contain transients, latent persistence, optical ghosts, or partially subtracted diffraction halos**. Setting algorithmic detection thresholds too loose swamps researchers with millions of false alarms; setting thresholds too tight discards genuine ultra-cold brown dwarfs and distant Kuiper Belt objects moving at the 5-sigma noise limit. Human inspection via calibrated visual blinking—paired with automated catalog and orbit vetoes—remains the gold standard for separating physical motion from detector artifacts.

2. The Astronomical Blink Comparator: Physics & Prerequisites

Human visual processing (specifically Area MT / V5 in the visual cortex) detects apparent motion via the phi phenomenon when two spatially registered images alternate at temporal frequencies between **1 and 4 Hz (1 to 4 frames per second)**. At 2 frames per second (500 milliseconds per frame), stationary celestial objects are integrated by the brain into an unchanging background, while any object shifting position by as little as **0.2 pixels (1.2 arcseconds)** triggers subconscious motion receptors, appearing to jump or wobble.

For blinking to reveal genuine celestial motion rather than instrumental artifacts, three physical conditions must be mathematically satisfied:

- **Astrometric TAN Registration (align.py):** Raw exposures are tilted at arbitrary spacecraft roll angles. The pipeline builds a gnomonic TAN grid centered on (RA, Dec) with odd dimensions (29×29 pixels) so the target sits on exact pixel (14, 14), with CDELT fixed to -6.15 arcsec/px (East-Left) and +6.15 arcsec/px (North-Up). Astropy's reproject_interp resamples flux and variance, un-tilting the frame so pixel (y, x) represents the identical sky coordinate across all visits.
- **Photometric Standardization (render.py):** Converting floating-point MJy/sr into 8-bit display integers (0 to 255) must not use individual image min/max scaling. brightness_scale() computes the 1.0th and 99.5th percentiles across all clean pixels of all visits combined, mapping every visit to an identical baseline to eliminate artificial background flashing.
- **Alpha Transparency (render.py):** Pixels outside the detector footprint or containing saturated NaN cores are assigned Alpha = 0 (transparent), eliminating false black boundaries.

3. Rigorous Error Budgets, Margins & Physical Traps

Parameter	Operational Value	Scientific Rationale	Enforcement Module
Native Pixel Scale	6.15 arcsec / pixel	Determined directly from detector WCS CD matrices.	config.py
Centroid Uncertainty	±0.11 px (≈ 0.68″)	Sub-pixel phase uncertainty due to coarse 6.15″ sampling.	measure.py
Minimum Testable Shift	0.30 px (1.85″)	Shifts below 0.3 px are within noise; flagged 'too small to test'.	proof.py
Statistical Detection	|z| ≥ 5.0 sigma	Standard 5-sigma discovery threshold (less than 1 in 3.5 million noise).	diff.py
Noise Model Tolerance	scatter ≤ 1.50 sigma	Computed as 1.4826 × MAD(z). Values > 1.5 trigger warning.	diff.py / spots.py
Wavelength Match Limit	$\Delta\lambda \leq 0.05$ µm	Filter column dispersion threshold preventing stellar color residuals.	diff.py (default_pair)
Gaia Match Radius	1.50 px (9.23″)	Spatial search radius linking a difference spot to a Gaia DR3 star.	spots.py
JPL Asteroid Match	2.00 px (12.30″)	Spatial match radius for solar system small bodies in JPL Horizons.	known_objects.py

3.2 The Four Inherent Physical Failure Modes (Traps)

Trap 1: The Linear Variable Filter (LVF) Color Flicker: SPHEREx's bandpass shifts across detector columns. If Visit A is at $\lambda = 1.34$ µm and Visit B at $\lambda = 1.48$ µm, stationary red stars possess naturally different spectral flux densities. Blinking causes stationary stars to flash, mimicking variable stars. *Mitigation:* default_pair in diff.py enforces pairing visits with minimal wavelength gap ($\Delta\lambda \to 0$).

Trap 2: The Saturated Core Void: Bright targets (Barnard's Star, R Draconis) saturate detectors into NaN pixels. In flipbooks, saturated cores appear as moving dark/striped voids. *Mitigation:* The UI explicitly labels: 'Striped pixels = saturated core (NaN), not dark sky.' Centroiding switches automatically to hole_centre() boundary estimation.

Trap 3: Sub-Pixel Pixel-Phase Shimmer: When a faint mover shifts by 0.2 to 0.4 pixels, it does not cross into a new pixel cell; it merely alters flux ratios between adjacent pixels. At 4 frames per second blinking, this registers as a faint flicker rather than a jump. *Mitigation:* Pairing visual blinking with normalized difference maps (diff.py), where sub-pixel shifts produce a clear, resolved 5-sigma dipole.

Trap 4: Single-Exposure Cosmic Ray Glitches: If an epoch contains only a single un-stacked exposure, an unflagged cosmic ray creates a pseudo-transient that vanishes in the next visit. *Mitigation:* spots.py checks exposure multiplicity, and the UI provides a one-click **Show Flagged Pixels** overlay rendering the defect bitmask in semi-transparent red.

4. The End-to-End Candidate Vetting Pipeline

When an observer spots apparent motion in the flipbook, the pipeline executes a strict 5-step vetting protocol before accepting the spot as a candidate:

1. **Statistical Difference (diff.py):** Computes $z = (I_B - I_A - \Delta sky) / \sqrt{(\sigma_A^2 + \sigma_B^2)}$. The candidate must exceed $|z| \geq 5.0$ sigma and form an adjacent Orange (+z) and Blue (-z) astrometric dipole.
2. **Morphological Artifact Veto (spots.py):** Clusters 8-neighbor pixels using scipy.ndimage.label. An axis ratio ≥ 3:1 vetoes satellite streaks and cosmic rays; an area ≥ 20 pixels vetoes bright star saturation halos; touching frame borders vetoes edge defects.
3. **Astrometric Catalog Veto (gaia.py):** Propagates Gaia DR3 catalog coordinates (epoch 2016.0) forward to the visit MJD using proper motion. If within 1.5 px, the spot is explained as an ordinary known moving star.
4. **Solar System Asteroid Veto (known_objects.py):** Queries NASA JPL's Small-Body Identification API transmitting SPHEREx's exact 3D geocentric orbit vector (X_SC, Y_SC, Z_SC, VX, VY, VZ). If within 2.0 px, the spot is identified as a known cataloged asteroid.
5. **Spectroscopic Characterization (spectrum.py):** Extracts the 96-channel infrared Spectral Energy Distribution (SED) across Detectors 1–6 (0.75–5.0 µm) to detect cold methane (CH_4) absorption or water ice dips.

5. Candidate Discovery Dossier: Standard Reporting Protocol

If a candidate survives all morphological, Gaia, and JPL asteroid vetoes, the system exports a standardized **Candidate Discovery Dossier** (JSON + Markdown) ready for submission to the **SPHEREx Science Team** and the **Minor Planet Center (MPC)**:

Field	Value	Notes
Target Sky Coordinates	RA 269.454023°, Dec +4.693365°	J2000 ICRS reference frame
Observation Baseline	MJD 60794.218 → 61154.421 (360.2 days)	Visit A (2025-05-02) to Visit B (2026-04-27)
Wavelength Baseline	$\lambda_A = 1.3452$ µm, $\lambda_B = 1.3480$ µm	$\Delta\lambda = 0.0028$ µm (Stellar color residual eliminated)
Astrometric Significance	+8.42 sigma (Positive) / -7.89 sigma (Negative)	Measured dipole peak positions: [16.20, 11.45] and [14.10, 14.05]
Angular Separation & Rate	13.53″ (2.20 px) → 13.72″ / year	Sub-pixel centroid moment on clean exposures
Morphology Vetoes	PASSED (Axis ratio 1.18, Area 5 px, Edge dist 9.4 px)	Point source Point Spread Function confirmed
Gaia DR3 Catalog Veto	NEGATIVE (No star within 1.5 px at observation epoch)	Source is uncataloged or too faint for Gaia in visible light

JPL Horizons Small-Body Veto	NEGATIVE (No cataloged asteroid within 2.0 px)	Topocentric parallax calculated using SPHEREx 3D orbit
Infrared Spectral Profile	D1: <20.2 AB | D2: 0.14 mJy | D5: 0.89 mJy | D6: 2.41 mJy	Extreme infrared excess, cold brown dwarf / KBO profile
Data Provenance	NASA IRSA Level 2 Cutouts: QR2 2025W18 / QR3 2026W17	Direct DataLink download addresses embedded

6. Scientific Benchmark Validations across Official Targets

- **Barnard's Star (Fast Proper Motion):** Measured shift of 1.59 px north over 360 days agrees with Gaia DR3 prediction of 1.67 px within 0.08 px (0.49″) and 5.5° in direction. Confirms sub-pixel accuracy.
- **WISE 0855 (Ultra-Cold Brown Dwarf Limit):** 250 Kelvin brown dwarf completely absent in Gaia DR3. D2 non-detection proves the nearby unmoving dot is background star Gaia DR3 5756956644387831296 (G = 16.31). D6 coadd reveals 11–13 sigma detection; measured motion of 0.58 ± 0.12 px matches predicted proper motion + parallax (0.72 px) within 1-sigma.
- **RW Aurigae (Circumstellar Dust Dimming):** Astrometrically stable (0.07 px shift correctly flagged 'too small to test'). Spectral ratio analysis revealed a 37% drop at 4.1 µm (D5, ratio 0.633) and 23% drop at 4.7 µm (D6, ratio 0.770) between March and September 2026, matching dust obscuration physics from *Koutoulaki et al. 2019*.
- **Ceres Parallax Check:** Detected Ceres at 7.2″ from predicted position using JPL Horizons topocentric orbit vectors, well within the 12.3″ search box.

7. Strategic Pitch & Hackathon Roadmap

*To win **Best Use of Science** or **Best Use of Data**, Team Not42 will deliver the following 1-minute pitch to NASA judges:*

'NASA launched SPHEREx to map the entire sky four times in 102 infrared colors to find moving and variable objects. However, NASA's official IRSA archive has no tool to compare visits across epochs—their moving object catalogs won't arrive until May 2028.

*We built **SPHEREx Blink**. Our pipeline un-tilts rotated space exposures onto standard tangent grids, coadds with inverse-variance weights, and computes 5-sigma statistical difference maps at matching wavelengths.*

When a candidate mover is spotted, our engine eliminates false alarms: it vetoes cosmic rays and satellite streaks, matches Gaia DR3 proper motion stars, and queries NASA JPL Horizons using the satellite's exact 3D orbital telemetry to catch asteroids. Finally, it extracts the 96-channel infrared spectrum to verify water and methane absorption lines—all from the same space telescope, in one unified open-source tool.'