

# Investigating the Adult Ixodid Tick Populations and Their Associated *Anaplasma*, *Ehrlichia*, and *Rickettsia* Bacteria at a Rocky Mountain Spotted Fever Hotspot in Western Tennessee

Rebecca T. Trout Fryxell,<sup>1</sup> Brain M. Hendricks,<sup>1</sup> Kimberly Pompo,<sup>1</sup> Sarah E. Mays,<sup>1</sup>  
Dave J. Paulsen,<sup>1</sup> Darwin J. Operario,<sup>2</sup> and Allan E. Houston<sup>3</sup>

## Abstract

Ehrlichiosis and rickettsiosis are two common bacterial tick-borne diseases in the southeastern United States. Ehrlichiosis is caused by ehrlichiae transmitted by *Amblyomma americanum* and rickettsiosis is caused by rickettsiae transmitted by *Amblyomma maculatum* and *Dermacentor variabilis*. These ticks are common and have overlapping distributions in the region. The objective of this study was to identify *Anaplasma*, *Ehrlichia*, and *Rickettsia* species associated with questing ticks in a Rocky Mountain spotted fever (RMSF) hotspot, and identify habitats, time periods, and collection methods for collecting questing-infected ticks. Using vegetation drags and CO<sub>2</sub>-baited traps, ticks were collected six times (May–September 2012) from 100 sites (upland deciduous, bottomland deciduous, grassland, and coniferous habitats) in western Tennessee. Adult collections were screened for *Anaplasma* and *Ehrlichia* (simultaneous polymerase chain reaction [PCR]) and *Rickettsia* using genus-specific PCRs, and resulting positive amplicons were sequenced. *Anaplasma* and *Ehrlichia* were only identified within *A. americanum* (*Ehrlichia ewingii*, *Ehrlichia chaffeensis*, Panola Mountain *Ehrlichia*, and *Anaplasma odocoilei* sp. nov.); more *Ehrlichia*-infected *A. americanum* were collected at the end of June regardless of habitat and collection method. *Rickettsia* was identified in three tick species; “*Candidatus Rickettsia amblyommii*” from *A. americanum*, *R. parkeri* and *R. andeanae* from *A. maculatum*, and *R. montanensis* (= *montana*) from *D. variabilis*. Overall, significantly more *Rickettsia*-infected ticks were identified as *A. americanum* and *A. maculatum* compared to *D. variabilis*; more infected-ticks were collected from sites May–July and with dragging. In this study, we report in the Tennessee RMSF hotspot the following: (1) *Anaplasma* and *Ehrlichia* are only found in *A. americanum*, (2) each tick species has its own *Rickettsia* species, (3) a majority of questing-infected ticks are collected May–July, (4) *A. americanum* and *A. maculatum* harbor pathogenic bacteria in western Tennessee, and (5) *R. rickettsii* remains unidentified.

**Keywords:** *Ehrlichia*, *Rickettsia*, ticks

## Introduction

**R**ICKETTSIOSIS AND EHRLICHIOSIS are two tick-borne diseases (TBDs) with expanding case and vector distributions and are common throughout the southeastern United States (CDC 2013a, b). Although many tick species harbor *Rickettsia* and/or *Ehrlichia*, studies in the southeast rarely investigate the role of bacterial coinfections (more than one bacteria per tick host), spillover effects (specific bacterial

species identified in more than one tick species), or the epidemiological behavior of questing-infected ticks (Mixon et al. 2006, Carmichael and Fuerst 2010, Heise et al. 2010, Berrada et al. 2011, Stromdahl et al. 2011). Five states (North Carolina, Arkansas, Tennessee, Oklahoma, and Missouri) report more than 60% of the cases of Rocky Mountain spotted fever (RMSF) and Ehrlichiosis (CDC 2013a, b). These clusters combined with the absence of *R. rickettsii* in ticks warrant additional investigations into which ticks and

<sup>1</sup>Department of Entomology and Plant Pathology, University of Tennessee, Knoxville, Tennessee.

<sup>2</sup>Division of Infectious Diseases and International Health, University of Virginia Health System, Charlottesville, Virginia.

<sup>3</sup>Department of Forestry, Wildlife and Fisheries, University of Tennessee, Knoxville, Tennessee.

pathogens are involved in transmission of TBDs in the southeastern United States.

The most common southeastern TBD is RMSF, caused by *R. rickettsii*, which is often fatal if left untreated (McDade and Newhouse 1986, CDC 2013b). The primary vector is *Dermacentor variabilis* (Ricketts 1906), yet *R. rickettsii* has not been recovered from southern collections of *D. variabilis* (Stromdahl et al. 2011). While *Amblyomma americanum* harbors the currently nonpathogenic “*Candidatus Rickettsia amblyommii*” (Goddard and Norment 1986), it was recently suggested as an additional vector of *R. rickettsii* in North Carolina (Breitschwerdt et al. 2011) and in Kansas (Berrada et al. 2011). In addition *Amblyomma maculatum* has a role in many pathogen transmission cycles, including rickettsiosis (*R. parkeri*) in humans (Sumner et al. 2007, Goddard and Varela-Stokes 2009, Trout et al. 2010). Diagnostic tests for RMSF are often cross-reactive with other spotted fever group rickettsiae (SFGR) (Parola et al. 2005, Ereemeeva et al. 2006a, Raoult and Parola 2008) and the presence of *R. parkeri* and “*Ca. Rickettsia amblyommii*” has raised further concerns about the true prevalence of RMSF infections in the United States (Paddock et al. 2004, Paddock 2005, Whitman et al. 2007, Apperson et al. 2008, Trout Fryxell et al. 2015b). Consequently, test results may incorrectly implicate RMSF instead of another SFGR (McCall et al. 2001). It is possible that cross-reactivity of *R. parkeri*, *R. montanensis*, and “*Ca. R. amblyommii*” is responsible for the increased incidence, declining case fatality, occurrence of multiple symptoms of spotted fevers, and increase in winter cases. Together, this has sparked several questions regarding whether a diagnostic error is occurring (e.g., false or cross-reactive positives), or circulating strains of *R. rickettsii* are becoming less virulent (Dantas-Torres 2007, Raoult and Parola 2008).

Ehrlichiosis is the second most common TBD in the southeast and is attributed to several different bacteria, including *Ehrlichia chaffeensis*, *Ehrlichia ewingii*, and *Panola Mountain Ehrlichia* (Holland et al. 1985, Anderson et al. 1991, Ewing et al. 1995, Buller et al. 1999, Loftis et al. 2008). The primary vector and amplifying reservoir of *Ehrlichia* species is *A. americanum*, a dominant tick species likely responsible for a majority of tick bites in the southeast (Anderson et al. 1993, Childs and Paddock 2003, Stromdahl and Hickling 2012). Within the United States, the number of ehrlichiosis cases has risen in the last several years (CDC 2013a). Complicating matters further, the clinical presentations of ehrlichiosis and rickettsiosis are similar.

Entomological investigations of tick vectors in Kentucky and Tennessee included screening field-collected specimens for *Rickettsia* or *Ehrlichia* (Cohen et al. 2010, Moncayo et al. 2010, Fritzen et al. 2011, Pagac et al. 2014, Harmon et al. 2015); rarely is the same tick screened for both organisms. Cohen et al. (2010) screened field-collected *A. americanum*, *D. variabilis*, *Ixodes texanus* (Banks), *I. cookie* Packard, *I. scapularis* (Say), and *A. maculatum* from 29 Tennessee counties for *Ehrlichia* and found only *A. americanum* to be infected (2.6% *E. chaffeensis* and 0.8% *E. ewingii*). Harmon et al. (2015) screened 303 *A. americanum* collected from a community experiencing an ehrlichiosis outbreak and found *E. ewingii* (5.3%), *E. chaffeensis* (2.0%), and *Panola Mountain Ehrlichia* (2.0%) in the questing population. Moncayo et al. (2010) screened host-collected and field-collected ticks from 49 counties for *Rickettsia* and found both “*Ca. R. amblyommii*” and *R. montana* in both *D. variabilis* and


*A. americanum*, and *R. cooleyi* in *I. scapularis*. In Kentucky, Fritzen et al. (2011) screened *A. americanum* and *D. variabilis* and identified *Borrelia lonestari*, *E. chaffeensis*, and “*Ca. R. amblyommii*” in *A. americanum*, and *R. montana* and *R. parkeri* in *D. variabilis*. In that study they also identified four “*Ca. R. amblyommii*”-positive *A. americanum* coinfecting with *E. ewingii* (2 specimens), *E. chaffeensis* (1 specimen), and *B. lonestari* (1 specimen). In addition, Pagac et al. (2014) screened 105 *A. maculatum* and 299 *D. variabilis* collected from military training sites in Kentucky and Tennessee and identified *R. parkeri* in *A. maculatum* (14.3%) and *R. montanensis* in *D. variabilis* (3.3%). None of the recovered *Rickettsia* sequences was identified as *R. rickettsii* (Moncayo et al. 2010, Fritzen et al. 2011, Pragac et al. 2014). While these studies provide a foundation for which pathogen and tick are in the area, they do not explain the increased incidence of TBDs in the region; epidemiological studies are needed.

As mentioned, in the United States, RMSF fatalities are decreasing, while diagnoses are increasing and severity or health outcomes are variable (CDC 2013b). Of interest are the RMSF cases in Tennessee. Tennessee accounts for only 2.4% of hospitalized RMSF cases in the United States; however, hospitalizations in the Tennessee Valley region result in 26% of the nation’s RMSF fatalities with a majority of these cases occurring in central and western Tennessee, indicating that this area of the country is an RMSF hotspot (Chapman 2006, Adjemian et al. 2009). The objective of this study was (1) to identify *Anaplasma*, *Ehrlichia*, and *Rickettsia* species associated with questing ticks in a rickettsiosis hotspot in a complex environment representative of the southeastern landscape and (2) to identify habitats, time periods, and collection methods for collecting questing-infected ticks. In this study, we collected more than 9000 ticks representing three species in a single year, screened the adults for *Anaplasma*, *Ehrlichia*, and *Rickettsia*, and analyzed the collections by collection variables (month, habitat, and trapping method) to better understand the complex TBD associations in an RMSF hotspot.

## Materials and Methods

### Site selection

Ames Plantation (35.1153N, -89.2167W) (AMES), a University of Tennessee-managed research and education facility in western Tennessee, is located within the rickettsiosis hotspot identified by Adjemian et al. (2009) and has had a number of employees who live at the center who have been previously diagnosed with SFGR or similar tick-borne ailments. The facility is used for research (forestry, wildlife, ecological, and historical), education (many courses are taught at the facility), crop production (timber, livestock, field and row crops), and private recreation (sport hunting and host to the annual National Championship field trials) making the site frequented by numerous people. The tick sampling effort was spread across AMES using a superimposed 100-acre grid (0.40 km<sup>2</sup>). One hundred grids were randomly selected and a 100 m transect placed to represent a single habitat type: 25 native warm season grassland, 37 upland deciduous, 19 bottomland deciduous, and 19 coniferous sites (Fig. 1a). These habitats were designated based on the vegetation and soil type within each site, and additional details are reported in Trout Fryxell et al. (2015a).



**FIG. 1.** Ixodid ticks were collected from Ames Plantation (AMES) during 2012 from 100 sites representing native warm season grassland, upland deciduous, bottomland deciduous, and coniferous sites (A). The collections consisted of three tick species, *Amblyomma americanum*, *Amblyomma maculatum*, and *Dermacentor variabilis*; total numbers collected at each site are presented next to each pie chart representing the proportion of each species collected (adults and nymphs) (B). Adult ticks were positive for *Ehrlichia*/*Anaplasma* (C) and *Rickettsia* (D) and were collected throughout the study area. Figure 1 can be viewed in greater detail online at [www.liebertpub.com/vbz](http://www.liebertpub.com/vbz)

#### Tick collections

Ticks were collected six times from May through September 2012, when a majority of Tennessee TBDs are diagnosed from each transect using vegetation drags and carbon dioxide (CO<sub>2</sub>)-baited traps (Falco and Fish 1992, Mays et al. 2016). Drags were constructed from a 1 m<sup>2</sup> piece of corduroy cloth attached to a wooden dowel rod. The drag was inspected and ticks collected every 20 m along a 100 m transect. CO<sub>2</sub>-baited traps were made from a 1.9-L cylindrical blue cooler (Igloo Corporation, Katy, TX) with holes drilled around the circumference (Wilson et al. 1972). They were filled with ~2.3 kg of dry ice and placed on a 1.37 m<sup>2</sup> piece of light-colored duck cloth at the beginning of the 100 m transect and remained on site 12–16 h. Ticks attracted to the cloth and the cooler were removed in the field. CO<sub>2</sub>-baited traps were used at 10 sites in May, and then all 100 sites June through September. All ticks were collected and stored in trap/site-specific vials containing 80% ethanol, and specimens were

identified to species, life stage, and sex (Cooley and Kohls 1944, Keirans and Litwak 1989, Keirans and Durden 1998).

#### Bacterial screening of adult ixodid ticks

Adult specimens were used in this study because they had the greatest chance of acquiring pathogens. Molecular methods were conducted in different facilities with equipment specific to each procedure to prevent contamination. Tick disruption procedures were identical to those reported in Hendricks (2013). Briefly, ticks were longitudinally bisected with a sterile razor blade and DNA was extracted using the Fermentas Gene Jet Genomic DNA Purification Kit and protocol (ThermoFisher Scientific, Pittsburgh, PA), which yielded 200 µL of genomic DNA; the total genomic DNA (tick and potential pathogen/host) was stored at –20°C until bacterial screening.

After DNA extraction, ticks were screened for *Rickettsia* species by PCR amplification of the 190 kDa protein antigen

gene also referred to as the *ompA* gene hereafter (Eremeeva et al. 1994) and for *Ehrlichia/Anaplasma* species by a simultaneous nested PCR amplification of the *groEL* gene (Tabara et al. 2007; Takano et al. 2009), in separate master mixes using Maxima Hot Start Green PCR Master Mix (Thermo Scientific, Pittsburgh, PA). Extracts were initially tested in pools: 5  $\mu$ L of 10 individual extracted specimens were pooled into a single tube ( $\sim 50$   $\mu$ L pooled DNA), and 5  $\mu$ L of that pooled DNA was used for primary screening for each pathogen in a 50  $\mu$ L reaction (5  $\mu$ L pooled DNA, 25  $\mu$ L of Maxima Hot Start Green PCR Master Mix, 1  $\mu$ L each of forward and reverse primers, and 18  $\mu$ L nuclease-free water). If a pool tested positive, individual extracts from the positive pool were then screened for each pathogen in a 25  $\mu$ L reaction (2  $\mu$ L of individual tick DNA, 12  $\mu$ L of Hot Start Master Mix, 0.5  $\mu$ L each of the forward and reverse primers, and 10  $\mu$ L of nuclease-free water). Primers and thermal cycler conditions are provided in Hendricks (2013). Positive controls consisted of tick DNA extractions previously testing positive for *E. chaffeensis* (99% identical to GenBank KJ907753L10917) or “*Ca. R. amblyommii*” (99% identical to GenBank HM446483). Positive PCR products were identified via gel electrophoresis (1.5% agarose gel: 1 $\times$ TAE buffer with ethidium bromide for 2 h at 120 V).

Selected positive amplicons were bidirectionally sequenced at the University of Tennessee Division of Biology Sequencing Facility to confirm positivity, to identify the bacterial species, and to classify the bacterial genotype (Trout et al. 2010). Due to the few *groEL* PCR positives, all *groEL* amplicons were sequenced. A random sample of *Rickettsia* PCR products were selected for sequencing because many of the ticks were positive. Resulting sequence reads were aligned in Sequencher 5.1 (Gene Codes Corporation Ann Arbor, MI) and compared to GenBank submissions via NCBI BLAST using the default conditions (other database, nonmodel organism optimized for highly similar sequences). All unique sequences were submitted to GenBank (KX158256-158278).

#### Identifying bacterial coinfections within adult ixodid ticks

In addition, to check for coinfections (more than one *Ehrlichia*, *Anaplasma*, or *Rickettsia* within a tick sample), all specimens PCR positive for either *Ehrlichia* or *Rickettsia* PCR were subjected to species-specific PCR testing as described in Gaines et al. (2014). Briefly, the six-plex test described in Gaines et al. was split into two three-plex reactions based on genus. Testing was performed in the ABI 7900 HT Fast Real-time PCR system (#4351405 Life Technologies; Thermo Fisher Scientific, Inc. Pittsburgh, PA). Reactions were performed with the TaqMan Real-time PCR master mix (Life Technologies) in a 25  $\mu$ L reaction volume with 5  $\mu$ L of DNA extract. Thermal cycling conditions, primers, and TaqMan probes (Life Technologies) were used as previously described (Gaines et al. 2014).

#### Epidemiological-collection associations (species, month, habitat, and trap)

To determine overall trends in the data set ANOVA, summary statistics, and pathogen abundance (overall and within each tick species) were calculated using JMP<sup>®</sup> Pro 11.1.1 (SAS Institute, Inc. Cary, NC) ( $\alpha=0.05$ ). Pathogen prevalence was calculated by dividing the number of PCR-positive samples by the total number of ticks screened.

Contingency tests ( $\chi^2$ ) were used to compare the frequencies of positive and negative ticks to determine if the tick species (3 species), collection period (3 species $\times$ 5 collection trips), habitat (3 species $\times$ 4 habitats), or trapping method (3 species $\times$ 2 trapping methods) was associated with the probability the tick would be PCR positive for either pathogen. For the contingency tests, Bonferroni's correction with 36 comparisons (species, collection period, habitat, and trapping method) was used to adjust calculated *p* values to prevent type I error caused by the various comparisons. Based on Bonferroni's correction, comparisons that yielded a *p* value  $\leq 0.0014$  were considered statistically significant. An additional analysis of logistic regression was calculated in SAS 9.4 TS1M3 for Windows 64 $\times$  (SAS Institute, Inc., Cary, NC) to determine the odds of collecting a positive tick for *Ehrlichia* or *Rickettsia* in a specific habitat and/or during a specific period. The trapping method variable was not included in the analysis because of the inherent differences in sampling methods (dragging for 5 min vs. baiting for 12<sup>+</sup> h).

## Results

### Tick collections

Three tick species were collected totaling 9450 ticks: 6900 *A. americanum* (5974 nymphs and 926 adults), 2530 *D. variabilis* (5 nymphs and 2525 adults), and 20 adult *A. maculatum* (Fig. 1b). No *I. scapularis* or *Rhipicephalus sanguineus* specimens were collected. Twenty *A. americanum* adults were saved as voucher specimens (*i.e.*, not included in the pathogen analyses). Ten of the *D. variabilis* lacked appropriate collection labels and were not included in the categorical analyses (collection month, trapping method, or habitat).

Briefly, *A. americanum* nymphs were collected significantly more in bottomland deciduous sites at the end of June (third sampling period) with a CO<sub>2</sub> trap and rarely collected in upland deciduous sites or past the August sampling event (fourth sampling period) ( $F=3.3867$ ;  $df=46$ , 1063;  $p<0.0001$ ). *A. americanum* adults were collected significantly more at coniferous sites during early May and June (first and second sampling period) with a CO<sub>2</sub> trap and rarely collected past the August sampling event (fourth sampling period) ( $F=7.6422$ ;  $df=46$ , 1063;  $p<0.0001$ ).

*D. variabilis* adults were collected significantly more from both deciduous sites during early June (second sampling period) with a CO<sub>2</sub> trap than in grassland or coniferous sites later in the sampling season ( $F=6.2883$ ;  $df=46$ , 1063;  $p<0.0001$ ). *Amblyomma maculatum* were not collected sufficiently for comparisons, but 16 of the 20 specimens were collected from grassland sites. Overall, a majority of ticks were collected from deciduous sites early in the sampling period ( $F=1073.93$ ;  $df=46$ , 1063;  $p<0.0001$ ).

### Bacterial screening of adult ixodid ticks

None of the *D. variabilis* or *A. maculatum* was *groEL* PCR positive. *Ehrlichia* or *Anaplasma* was identified in 17 (1.8%) of the 926 *A. americanum* (Table 1); 12 were 97–100% identical to *E. ewingii* (GenBank KJ907744, identified as *Ehrlichia* 1–4), 2 were 100% identical to Panola Mountain *Ehrlichia* (GenBank HQ658904, identified as *Ehrlichia*-5), 1 was 99% identical to *E. chaffeensis* (GenBank KJ907753,

TABLE 1. SIX *EHRLICHIA*, 1 *ANAPLASMA*, AND 9 *RICKETTSIA* GENOTYPES WERE IDENTIFIED FROM TICKS COLLECTED IN SOUTHWESTERN TENNESSEE AND SEQUENCES WERE UPLOADED TO GENBANK (KX158256-158278)

Genotype	Genotype frequency	Amblyomma americanum	Amblyomma maculatum	Dermacentor variabilis	% Identical (GenBank)	Publication (PubMed no.)
<i>groEL</i> amplicons for <i>Ehrlichia</i>						
<i>Ehrlichia</i> -1	9	9	0	0	100% <i>Ehrlichia ewingii</i> (KJ 907744)	Lee et al. unpublished
<i>Ehrlichia</i> -2	1	1	0	0	100% <i>E. ewingii</i> (AF195273)	Lee et al. unpublished
<i>Ehrlichia</i> -3	1	1	0	0	99% <i>E. ewingii</i> (KJ 907744)	Lee et al. unpublished
<i>Ehrlichia</i> -4	1	1	0	0	99% <i>E. ewingii</i> (KJ 907744)	Lee et al. unpublished
<i>Ehrlichia</i> -5	2	2	0	0	100% <i>Ehrlichia</i> Panola Mtn. (HQ658904)	Harmon et al. 2010 unpublished
<i>Ehrlichia</i> -6	1	1	0	0	99% <i>Ehrlichia chaffeensis</i> (KJ907753)	Lee et al. unpublished
<i>Anaplasma</i> -1	2	2	0	0	99% <i>Anaplasma odocoilei</i> (JX876642)	Tate et al. 2013 (23276749)
<i>ompA</i> amplicons for <i>Rickettsia</i>						
<i>Rickettsia</i> -1	41	41	0	0	100% <i>Ca. R. amblyommii</i> (JF694090)	Hun et al. 2011 (21612539)
<i>Rickettsia</i> -2	1	1	0	0	99% <i>Ca. R. amblyommii</i> (JF694090)	Hun et al. 2011 (21612539)
<i>Rickettsia</i> -3	1	1	0	0	99% <i>Ca. R. amblyommii</i> (JF694090)	Hun et al. 2011 (21612539)
<i>Rickettsia</i> -4	1	1	0	0	99% <i>Ca. R. amblyommii</i> (JF694090)	Hun et al. 2011 (21612539)
<i>Rickettsia</i> -5	1	1	0	0	99% <i>Ca. R. amblyommii</i> (JF694090)	Hun et al. 2011 (21612539)
<i>Rickettsia</i> -6	4	0	4	0	100% <i>R. parkeri</i> (KF782320)	Monje et al. 2014 (25113981)
<i>Rickettsia</i> -7	1	0	1	0	100% <i>R. andeanae</i> (KF179352)	Ogrzewalska et al. 2014 (24231270)
<i>Rickettsia</i> -8	1	0	0	1	99% <i>Rickettsia</i> sp. D192 (AY543683)	Ammerman and Norris 2004 (15496254)
<i>Rickettsia</i> -9	30	0	0	30	100% <i>Rickettsia</i> sp. D192 (AY543683)	Ammerman and Norris 2004 (15496254)

identified as *Ehrlichia*-6), and 2 were 99% identical to *Anaplasma odocoilei* sp. (GenBank JX876642, identified as *Ehrlichia*-7) (Table 1).

*Rickettsia* was identified in 353 (38.12%) of the 926 *A. americanum*, 92 (3.66%) of the 2515 *D. variabilis*, and 5 (25%) of the 20 *A. maculatum*. From the 45 sequenced *A. americanum ompA* amplicons, 5 different *Rickettsia* sequence variants were identified (Table 1), but all were identified as >99% identical to “*Ca. R. amblyommii*” (GenBank KT722804) (identified as *Rickettsia* 1–5, Table 1). Four of the *A. maculatum Rickettsia*-positives were 100% identical to *R. parkeri* (GenBank KF782320, identified as *Rickettsia*-6) and one was 100% identical to *R. andeanae* (GenBank KF179352, identified as *Rickettsia*-7). The 31 *Rickettsia* amplicons sequenced from *D. variabilis* were >99% identical to nonpathogenic *R. montanensis* (= *R. montana*) (GenBank AY543681); 1 was identified as *Rickettsia*-8 (Table 1) and 30 were identified as *Rickettsia*-9 (Table 1).

#### Identifying bacterial coinfections within adult ixodid ticks

Eight of the *A. americanum* specimens were coinfecting; six were coinfecting with *E. ewingii* and “*Ca. R. am-*

*blyommii*,” two with Panola Mountain *Ehrlichia* and “*Ca. R. amblyommii*.” The number of coinfecting ticks was not different from expected ( $X^2=0.07$ ;  $df=1$ ;  $p=0.2463$ ). No trend or pattern was evident in the coinfecting ticks; they were collected from eight different sites, throughout the sampling period (2 specimens in May, 5 specimens in July, and 1 specimen in August), with both trap types (2 specimens with a drag and 6 with the CO<sub>2</sub> trap). Five coinfecting ticks were collected in upland deciduous sites and three were collected in coniferous sites. Coinfections of more than one *Ehrlichia* or *Rickettsia* were not identified using the novel real-time PCR assays described in Gaines et al. (2014). The multiplex data corroborated the sequencing results—identification of *R. parkeri* in *A. maculatum* and identification of “*Ca. R. amblyommii*” in *A. americanum* (Table 1).

#### *Ehrlichia*/*Anaplasma*-infected ixodid ticks, epidemiological-collection associations

*Ehrlichia* and *Anaplasma* were only amplified from 17 adult *A. americanum* ticks, in May (5/215), July (10/242), and August (2/78) (Table 2), indicating positive ticks were collected throughout the season; but more were collected from June 25 to July 6 (4.13% positive) compared to one sampling

TABLE 2. *A. AMERICANUM* POLYMERASE CHAIN REACTION POSITIVES WITH *E. EWINGII*, *E. CHAFFEENSIS*, PANOLA MOUNTAIN *EHRLICHIA*, OR *A. ODOCOILEI* WERE COLLECTED PRIMARILY DURING JUNE AND JULY IN THE SOUTHEASTERN UNITED STATES

Trapping dates (period)	Trapping method	No. positive/no. screened (% positive)	Southeastern United States Habitat Type (no. positive/no. screened)			
			Grasslands (n = 25 sites), n (%)	Coniferous (n = 19 sites), n (%)	Bottomland deciduous (n = 19 sites), n (%)	Upland deciduous (n = 37 sites), n (%)
May 8–16 (1)	Dragging	5/164 (3.0)	2/24 (8.3)	1/38 (2.6)	0/19 (0)	2/83 (2.4)
	CO <sub>2</sub> -trapping	0/51 (0)	0/0 (0)	0/35 (0)	0/3 (0)	0/13 (0)
	Total	5/215 (2.3)	2/24 (8.3)	1/73 (1.4)	0/22 (0)	2/96 (2.1)
May 28–June 8 (2)	Dragging	0/115 (0)	0/3 (0)	0/26 (0)	0/28 (0)	0/58 (0)
	CO <sub>2</sub> -trapping	0/269 (0)	0/23 (0)	0/54 (0)	0/56 (0)	0/136 (0)
	Total	0/384 (0)	0/26 (0)	0/80 (0)	0/84 (0)	0/194 (0)
June 25–July 6 (3)	Dragging	3/42 (7.1)	1/2 (50.0)	2/10 (20.0)	0/9 (0)	0/21 (0)
	CO <sub>2</sub> -trapping	7/200 (3.5)	0/9 (0)	3/74 (4.1)	0/49 (0)	4/68 (5.9)
	Total	10/242 (4.1)	1/11 (9.1)	5/84 (6.0)	0/58 (0)	4/89 (4.5)
July 23–August 3 (4)	Dragging	0/18 (0)	0/2 (0)	0/5 (0)	0/3 (0)	0/8 (0)
	CO <sub>2</sub> -trapping	2/60 (3.3)	0/1 (0)	1/6 (16.7)	1/21 (4.8)	0/32 (0)
	Total	2/78 (2.6)	0/3 (0)	1/11 (9.1)	1/24 (4.2)	0/40 (0)
August 20–29 (5)	Dragging	0/2 (0)	0/0 (0)	0/0 (0)	0/1 (0)	0/1 (0)
	CO <sub>2</sub> -trapping	0/5 (0)	0/0 (0)	0/0 (0)	0/2 (0)	0/3 (0)
	Total	0/7 (0)	0/0 (0)	0/0 (0)	0/3 (0)	0/4 (0)
September 21–29 (6)	Dragging	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	CO <sub>2</sub> -trapping	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	Total	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
Total	Dragging	8/341 (2.3)	3/31 (9.7)	3/79 (3.8)	0/60 (0)	2/171 (1.2)
	CO <sub>2</sub> -trapping	9/585 (1.5)	0/33 (0)	4/169 (2.4)	1/131 (0.8)	4/252 (1.6)
	Total	17/926 (1.8)	3/64 (4.7)	7/248 (2.8)	1/191 (0.5)	6/423 (1.4)

trip prior (May 28–June 8; 0 positive ticks out of 384 screened) ( $X^2 = 14.9$ ;  $df = 4$ ;  $p = 0.005$ ). Slightly but not significantly more *Ehrlichia/Anaplasma*-infected *A. americanum* were collected from grassland sites (4.69% positive) compared to bottomland deciduous sites (0.52% positive) ( $F = 2.1597$ ;  $df = 3, 922$ ;  $p = 0.00913$ ) (Fig. 1c). The collection method ( $F = 0.7786$ ;  $df = 1, 924$ ;  $p = 0.3778$ ) and sex of *A. americanum* ( $F = 0.4504$ ;  $df = 2$ ;  $p = 0.6375$ ) were not significantly associated with *Ehrlichia/Anaplasma* infection. The logistic model for *Ehrlichia* indicated that period ( $\chi^2 = 0.0133$ ;  $df = 4$ ;  $p = 1.00$ ), habitat ( $\chi^2 = 0.0088$ ;  $df = 3$ ;  $p = 0.9998$ ), and their interactive effects ( $\chi^2 = 0.7275$ ;  $df = 4$ ;  $p = 1.00$ ) were not significantly associated with encountering an *Ehrlichia*-infected *A. americanum*. Prevalence rates for each habitat and period for *Ehrlichia/Anaplasma* infection are presented in Table 2.

#### *Rickettsia*-infected ixodid ticks, epidemiological-collection associations

While each tick species was positive with its own *Rickettsia*, *Rickettsia* prevalence rates (no. positive/no. screened) were significantly greater in *A. americanum* (38.12% positive) and *A. maculatum* (25.0% positive) compared to *D. variabilis* (3.64% positive) ( $F = 450.9094$ ;  $df = 2, 3468$ ;  $p < 0.0001$ ). *Rickettsia*-positive ticks were slightly but not significantly associated with habitat (Table 3,  $F = 3.5346$ ;  $df = 3, 3457$ ;  $p = 0.0142$ ) such that collections of *Rickettsia*-infected ticks were more abundant in coniferous sites (15.84% positive) compared to grassland sites (8.78% posi-

tive) (Fig. 3d). *Rickettsia* infection did not vary by month within each species: *A. americanum* ( $\chi^2 = 5.79$ ;  $df = 4$ ;  $p = 0.215$ ), *A. maculatum* ( $\chi^2 = 1.78$ ;  $df = 4$ ;  $p = 0.777$ ), and *D. variabilis* ( $\chi^2 = 2.05$ ;  $df = 4$ ;  $p = 0.726$ ); however, there was an overall significant collection effect such that more *Rickettsia*-infected ticks were collected earlier in the season (May through July) compared to later in the season (August through September) ( $F = 9.0641$ ;  $df = 5, 3465$ ;  $p < 0.0001$ ). This was confirmed when the data set as a whole was analyzed in the logistic model, which indicated that only period ( $\chi^2 = 35.5775$ ;  $df = 5$ ;  $p < 0.0001$ ) exerts significant effect on the odds of encountering a *Rickettsia*-infected tick; this was not true for habitat ( $\chi^2 = 0.6030$ ;  $df = 3$ ;  $p = 0.6030$ ) or period  $\times$  habitat ( $\chi^2 = 18.1107$ ;  $df = 12$ ;  $p = 0.1124$ ). Odds ratios are presented in Table 4. Interestingly, significantly more *Rickettsia*-infected ticks were collected via dragging (19.53% positive) compared to the CO<sub>2</sub> trap (11.29% positive) ( $F = 17.8556$ ;  $df = 2, 3468$ ;  $p < 0.0001$ ; eight ticks were collected with an unknown method).

Within species, there was no significant habitat, seasonal, or collection effect for *A. americanum* or *A. maculatum* ( $p > 0.05$ ), but there was a slight habitat effect (which was not significant after the correction) within *D. variabilis*. Contingency tests indicated that slightly but not significantly more *Rickettsia*-positive *D. variabilis* were collected in upland deciduous (4.5% positive) and coniferous sites (4.4%) compared to grassland (1.3% positive) and bottomland deciduous sites (1.8% positive) ( $\chi^2 = 12.9$ ;  $df = 3$ ;  $p = 0.005$ ). Contingency tests also indicated significantly more *Rickettsia*-positive *A. americanum* were collected in bottomland deciduous sites

TABLE 3. OF THE THREE IXODID TICKS COLLECTED IN WESTERN TENNESSEE, SIGNIFICANTLY, MORE *A. AMERICANUM* AND *A. MACULATUM* WERE *RICKETTSIA* POSITIVE COMPARED TO *D. VARIABILIS*, AND CONIFEROUS SITES EARLIER IN THE SEASON (MAY–JULY) WERE MOST LIKELY TO HARBOR THESE *RICKETTSIA*-INFECTED TICKS

		Southeastern United States Habitat Type (no. positive/no. screened)				
Trapping dates (period)	Trapping method	No. positive/no. screened (% positive)	Grasslands (n=25 sites), n (%)	Coniferous (n=19 sites), n (%)	Bottomland deciduous (n=19 sites), n (%)	Upland deciduous (n=37 sites), n (%)
<i>A. Americanum</i>						
May 8–16 (1)	Dragging	52/164 (31.7)	7/24 (29.2)	11/38 (28.9)	9/19 (47.4)	25/83 (30.1)
	CO <sub>2</sub> -trapping	15/51 (29.4)	0/0 (0)	9/35 (25.7)	0/3 (0)	6/13 (46.2)
	Total	67/215 (31.2)	7/24 (29.2)	20/73 (27.4)	9/22 (40.9)	31/96 (32.3)
May 28–June 8 (2)	Dragging	52/115 (45.2)	2/3 (66.7)	9/26 (34.6)	12/28 (42.9)	29/58 (50)
	CO <sub>2</sub> -trapping	102/269 (37.9)	9/23 (39.1)	23/54 (42.6)	20/56 (35.7)	50/136 (36.8)
	Total	154/384 (40.1)	11/26 (42.3)	32/80 (40.0)	32/84 (38.1)	79/194 (40.7)
June 25–July 6 (3)	Dragging	13/42 (31.0)	0/2 (0)	5/10 (50)	5/9 (55.6)	3/21 (14.3)
	CO <sub>2</sub> -trapping	84/200 (42)	1/9 (11.1)	34/74 (45.9)	20/49 (40.8)	29/68 (42.6)
	Total	97/242 (40.1)	1/11 (9.1)	39/84 (46.4)	25/58 (43.1)	32/89 (36.0)
July 23–August 3 (4)	Dragging	11/18 (61.1)	1/2 (50)	3/5 (60)	2/3 (66.6)	5/8 (62.5)
	CO <sub>2</sub> -trapping	21/60 (35)	1/1 (100)	2/6 (33.3)	9/21 (42.9)	9/32 (28.1)
	Total	32/78 (41.0)	2/3 (66.7)	5/11 (45.5)	11/24 (45.8)	14/40 (35.0)
August 20–29 (5)	Dragging	1/2 (50)	0/0 (0)	0/0 (0)	1/1 (100)	0/1 (0)
	CO <sub>2</sub> -trapping	2/5 (40)	0/0 (0)	0/0 (0)	1/2 (50)	1/3 (33.3)
	Total	3/7 (42.9)	0/0 (0)	0/0 (0)	2/3 (66.7)	1/4 (25.0)
September 21–29 (6)	Dragging	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	CO <sub>2</sub> -trapp	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	Total	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
Total	Dragging	129/341 (37.8)	10/31 (32.3)	28/79 (35.4)	29/60 (48.3)	62/171 (36.3)
	CO <sub>2</sub> -trapping	224/585 (38.3)	11/33 (33.3)	68/169 (40.2)	50/131 (38.2)	95/252 (37.7)
	Total	353/926 (38.1)	21/64 (32.8)	96/248 (38.7)	79/191 (41.4)	157/423 (37.1)
<i>A. maculatum</i>						
May 8–16 (1)	Dragging	1/2 (50)	1/2 (50)	0/0 (0)	0/0 (0)	0/0 (0)
	CO <sub>2</sub> -trapping	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	Total	1/2 (50)	1/2 (50)	0/0 (0)	0/0 (0)	0/0 (0)
May 28–June8 (2)	Dragging	0/1 (0)	0/1 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	CO <sub>2</sub> -trapping	2/5 (40)	1/2 (50)	0/0 (0)	1/2 (50)	0/1 (0)
	Total	2/6 (33.3)	1/3 (33.3)	0/0 (0)	1/2 (50)	0/1 (0)
June 25–July 6 (3)	Dragging	0/4 (0)	0/4 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	CO <sub>2</sub> -trapping	1/2 (50)	1/1 (100)	0/0 (0)	0/0 (0)	0/1 (0)
	Total	1/6 (16.7)	1/5 (20.0)	0/0 (0)	0/0 (0)	0/1 (0)
July 23–August 3 (4)	Dragging	0/1 (0)	0/1 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	CO <sub>2</sub> -trapping	0/1 (0)	0/1 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	Total	0/2 (0)	0/2 (0)	0/0 (0)	0/0 (0)	0/0 (0)
August 20–29 (5)	Dragging	0/2 (0)	0/2 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	CO <sub>2</sub> -trapping	1/2 (50)	1/2 (50)	0/0 (0)	0/0 (0)	0/0 (0)
	Total	1/4 (25.0)	1/4 (25.0)	0/0 (0)	0/0 (0)	0/0 (0)
September 21–29 (6)	Dragging	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	CO <sub>2</sub> -trapping	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	Total	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
Total	Dragging	1/10 (10)	1/10 (10)	0/0 (0)	0/0 (0)	0/0 (0)
	CO <sub>2</sub> -trapping	4/10 (40)	3/6 (50)	0/0 (0)	1/2 (50)	0/2 (0)
	Total	5/20 (25.0)	4/16 (25.0)	0/0 (0)	1/2 (50)	0/2 (0)
<i>D. variabilis</i>						
May 8–16 (1)	Dragging	3/62 (4.8)	0/9 (0)	0/7 (0)	1/13 (7.7)	2/33 (6.1)
	CO <sub>2</sub> -trapping	0/17 (0)	0/0 (0)	0/0 (0)	0/3 (0)	0/14 (0)
	Total	3/79 (3.8)	0/9 (0)	0/7 (0)	1/16 (6.3)	2/47 (4.3)
May 28–June8 (2)	Dragging	1/68 (1.5)	0/1 (0)	0/13 (0)	0/19 (0)	1/35 (2.9)
	CO <sub>2</sub> -trapping	28/917 (3.1)	0/126 (0)	11/106 (10.4)	0/187 (0)	17/498 (3.4)
	Total	29/985 (2.9)	0/127 (0)	11/119 (9.2)	0/206 (0)	18/533 (3.4)
June 25–July 6 (3)	Dragging	2/118 (1.7)	0/6 (0)	0/27 (0)	1/24 (4.2)	1/61 (1.6)
	CO <sub>2</sub> -trapping	31/700 (4.4)	0/47 (0)	7/188 (3.7)	6/143 (4.2)	18/322 (5.6)
	Total	33/818 (4.0)	0/53 (0)	7/215 (3.3)	7/167 (4.2)	19/383 (5.0)

(continued)

TABLE 3. (CONTINUED)

Trapping dates (period)	Trapping method	No. positive/no. screened (% positive)	Southeastern United States Habitat Type (no. positive/no. screened)			
			Grasslands (n=25 sites), n (%)	Coniferous (n=19 sites), n (%)	Bottomland deciduous (n=19 sites), n (%)	Upland deciduous (n=37 sites), n (%)
July 23–August 3 (4)	Dragging	2/96 (2.1)	1/16 (6.3)	0/34 (0)	0/19 (0)	1/27 (3.7)
	CO <sub>2</sub> -trapping	21/466 (4.5)	1/31 (3.2)	4/108 (3.7)	1/76 (1.3)	15/251 (6.0)
	Total	23/562 (4.1)	2/47 (4.3)	4/142 (2.8)	1/95 (1.1)	16/278 (5.8)
August 20–29 (5)	Dragging	2/20 (10)	1/2 (50)	0/2 (0)	0/5 (0)	1/11 (9.1)
	CO <sub>2</sub> -trapping	2/49 (4.1)	0/1 (0)	0/12 (0)	0/9 (0)	2/27 (7.4)
	Total	4/69 (5.8)	1/3 (33.3)	0/14 (0)	0/14 (0)	3/38 (7.9)
September 21–29 (6)	Dragging	0/2 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/2 (0)
	CO <sub>2</sub> -trapping	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	Total	0/2 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/2 (0)
Total	Dragging	10/366 (2.7)	2/34 (5.9)	0/83 (0)	2/80 (2.5)	6/169 (3.6)
	CO <sub>2</sub> -trapping	82/2149 (3.8)	1/205 (0.5)	22/414 (5.3)	7/418 (1.7)	52/1112 (4.7)
	Total	92/2515 (3.7)	3/239 (1.3)	22/497 (4.4)	9/498 (1.8)	58/1281 (4.5)
All ticks collected						
May 8–16 (1)	Dragging	56/228 (24.6)	8/35 (22.9)	11/45 (24.4)	10/32 (31.3)	27/116 (23.3)
	CO <sub>2</sub> -trapping	15/68 (22.1)	0/0 (0)	9/35 (25.7)	0/6 (0)	6/27 (22.2)
	Total	71/296 (24.0)	8/35 (22.9)	20/80 (25.0)	10/38 (26.3)	33/143 (23.1)
May 28–June 8 (2)	Dragging	53/184 (28.8)	2/5 (40)	9/39 (23.1)	12/47 (25.5)	30/93 (32.3)
	CO <sub>2</sub> -trapping	132/1191 (11.1)	10/151 (6.6)	34/160 (21.3)	21/245 (8.6)	67/635 (10.6)
	Total	185/1375 (13.5)	12/156 (7.7)	43/199 (21.6)	33/292 (11.3)	97/728 (13.3)
June 25–July 6 (3)	Dragging	15/164 (9.1)	0/12 (0)	5/37 (13.5)	6/33 (1.8)	4/82 (4.9)
	CO <sub>2</sub> -trapping	116/902 (12.9)	2/57 (3.5)	41/262 (15.6)	26/192 (13.5)	47/391 (12.0)
	Total	131/1066 (12.3)	2/69 (2.9)	46/299 (15.4)	32/225 (14.2)	51/473 (10.8)
July 23–August 3 (4)	Dragging	13/115 (11.3)	2/19 (10.5)	3/39 (7.7)	2/22 (9.1)	6/35 (17.1)
	CO <sub>2</sub> -trapping	42/527 (8.0)	2/33 (6.1)	6/114 (5.3)	10/97 (10.3)	24/283 (8.5)
	Total	55/642 (8.6)	4/52 (7.7)	9/153 (5.9)	12/119 (10.1)	30/318 (9.4)
August 20–29 (5)	Dragging	3/24 (12.5)	1/4 (25)	0/2 (0)	1/6 (16.7)	1/12 (8.3)
	CO <sub>2</sub> -trapping	5/56 (8.9)	1/3 (33.3)	0/12 (0)	1/11 (9.1)	3/30 (10)
	Total	8/80 (10.0)	2/7 (28.6)	0/14 (0)	2/17 (11.8)	4/42 (9.5)
September 21–29 (6)	Dragging	0/2 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/2 (0)
	CO <sub>2</sub> -trapping	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/0 (0)
	Total	0/2 (0)	0/0 (0)	0/0 (0)	0/0 (0)	0/2 (0)
Total	Dragging	140/717 (19.5)	13/75 (17.3)	28/162 (17.3)	31/140 (22.1)	68/340 (20)
	CO <sub>2</sub> -trapping	310/2744 (11.3)	15/244 (6.1)	90/583 (15.4)	58/551 (10.5)	147/1366 (10.8)
	Total	450/3461 (13.0)	28/319 (8.8)	118/745 (15.8)	89/691 (12.9)	215/1706 (12.6)

TABLE 4. THE ODDS OF ENCOUNTERING *RICKETTSIA*-INFECTED TICKS WERE GREATER EARLIER IN THE SEASON (DURING PERIODS 1 AND 2) COMPARED TO LATER IN THE SEASON, REGARDLESS OF HABITAT

Period	Odds ratio (wald 95% CI)			
	Grasslands	Coniferous	Bottomland deciduous	Upland deciduous
1 vs. 2	<b>3.556 (1.328–9.516)</b>	1.209 (0.658–2.222)	<b>2.803 (1.250–6.287)</b>	<b>1.952 (1.252–3.042)</b>
1 vs. 3	<b>9.926 (1.979–49.787)</b>	<b>1.833 (1.011–3.326)</b>	2.154 (0.955–4.858)	<b>2.482 (1.528–4.034)</b>
1 vs. 4	3.556 (0.979–12.910)	<b>5.333 (2.297–12.383)</b>	<b>3.185 (1.248–8.125)</b>	<b>2.880 (1.677–4.947)</b>
1 vs. 5	0.741 (0.120–4.571)	>999 (<0.001 to >999)	2.679 (0.518–13.845)	2.850 (0.947–8.573)
2 vs. 3	2.792 (0.608–12.825)	1.516 (0.956–2.404)	0.768 (0.457–1.294)	1.272 (0.887–1.824)
2 vs. 4	1.000 (0.308–3.247)	<b>4.410 (2.076–9.367)</b>	1.136 (0.565–2.283)	1.476 (0.958–2.274)
2 vs. 5	0.208 (0.036–1.190)	>999 (<0.001 to >999)	0.956 (0.209–4.366)	1.460 (0.510–4.182)
3 vs. 4	0.358 (0.063–2.035)	<b>2.909 (1.384–6.116)</b>	1.478 (0.731–2.990)	1.160 (0.721–1.866)
3 vs. 5	<b>0.075 (0.009–0.647)</b>	>999 (<0.001 to >999)	1.244 (0.271–5.698)	1.148 (0.394–3.349)
4 vs. 5	0.208 (0.030–1.437)	>999 (<0.001 to >999)	0.841 (0.171–4.131)	0.990 (0.330–2.963)

*Bolded* values are significant and period 6 was not included in the table; all periods versus period 6 had odds ratio >999 (<0.001 to >999).



(41.4% positive) compared to grassland sites (32.8% positive) ( $\chi^2 = 43.6$ ;  $df = 3$ ;  $p < 0.0001$ ). The logistic model confirmed these findings. The logistic model indicated there was no significant risk of encountering *Rickettsia*-infected *A. americanum* by period ( $\chi^2 = 5.3804$ ;  $df = 4$ ;  $p = 0.2504$ ), habitat ( $\chi^2 = 3.0887$ ;  $df = 3$ ;  $p = 0.3782$ ), or their interactive effect ( $\chi^2 = 9.2827$ ;  $df = 10$ ;  $p = 0.5055$ ). The logistic model also indicated there was no significant risk of encountering *Rickettsia*-infected *A. maculatum* by period ( $\chi^2 = 0.6562$ ;  $df = 4$ ;  $p = 0.9566$ ), habitat ( $\chi^2 = 0.0043$ ;  $df = 2$ ;  $p = 0.9978$ ), or their interactive effect ( $\chi^2 = 0.000$ ;  $df = 1$ ;  $p = 0.9980$ ). The logistic model also indicated there was no significant risk of encountering *Rickettsia*-infected *D. variabilis* by period ( $\chi^2 = 0.0194$ ;  $df = 5$ ;  $p = 1.0$ ), habitat ( $\chi^2 = 1.6862$ ;  $df = 3$ ;  $p = 0.64$ ), or their interactive effect ( $\chi^2 = 13.1052$ ;  $df = 12$ ;  $p = 0.3614$ ). Interestingly, contingency tests indicated CO<sub>2</sub> trapping collected significantly fewer *Rickettsia*-infected *D. variabilis* in grassland sites (0.5% positive) compared to upland deciduous (4.7% positive) and coniferous (5.3% positive) sites ( $\chi^2 = 16.2$ ;  $df = 3$ ;  $p = 0.001$ ).

## Discussion

The confirmed identification of pathogenic *Ehrlichia* and *Anaplasma* in *A. americanum* and pathogenic *Rickettsia* in *A. maculatum* warrants additional research in the southeast and at this study site. This study also addressed bacterial associations with ticks collected concurrently from a site in a known RMSF hotspot environment. Adults were used in this study; however, investigations into nymph collections from different habitats and seasons may be more informative as they have had only one blood meal, are numerous (representing 63% of our collection, primarily *A. americanum*), and more difficult to detect making them more likely to stay on a host and transmit a pathogen (Falco et al. 1999). A majority of questing-infected ticks were collected May–July (period 1–3) regardless of habitats, during the same period when Tennesseans are diagnosed with rickettsiosis and ehrlichiosis. Additional investigations should be conducted just before determining when these infected ticks begin to quest for a host, and during this period to identify habitats, trapping methods, and evaluate methods for control.

Our results were similar to previous investigations as nonpathogenic *Rickettsia*, pathogenic *R. parkeri*, and several pathogenic *Ehrlichia*/*Anaplasma* were identified, while *R. rickettsii* was absent (Cohen et al. 2010, Moncayo et al. 2010, Harmon 2010, Fritzen et al. 2011, Stromdahl et al. 2011, Pagac et al. 2014). Our 38% *Rickettsia*-positive rate in *A. americanum* adults is important to consider as this species is most likely to bite humans in the southeastern United States and are typically faster and more aggressive than the other species collected at the site. While in the field working on this study, we encountered 167 ticks of which 158 were *A. americanum*, 7 were *D. variabilis*, and 2 were *A. maculatum*. We also encountered a number of larval ticks, but they were neither identified nor counted. All of the nonpathogenic *Rickettsia* identified within this study are cross-reactive with pathogenic *R. rickettsii* in diagnostic tests (McCall et al. 2001, Paddock et al. 2004, Paddock 2005, Paddock et al. 2005, Parola et al. 2005, Ereemeeva et al. 2006a, b, Whitman et al. 2007, Apperson et al. 2008, Raoult and Parola 2008). Other reasons that *R. rickettsii* may not

have been recovered include (1) *R. rickettsii* may have been in the positive nonsequenced *A. americanum* ( $n = 308$ ), (2) *Rickettsia* coinfections such that the most abundant *Rickettsia* were identified with PCR and *R. rickettsii* was not amplified, (3) *R. rickettsii* may be in the substantially greater nymph population, which was not screened (5979 nymphs out of 9450 ticks, 63% of collection), or (4) that the site is not a true RMSF hotspot and the disease of concern is misdiagnosed as RMSF. In addition, the use of a nested PCR for *Rickettsia* may have amplified additional positive samples, some of which may have been *R. rickettsii*.

The few *Ehrlichia*/*Anaplasma*-infected ticks made it difficult to identify preferred habitats and trapping methods (Table 2); however, it was clear that *A. americanum* harbors several different *Ehrlichia*/*Anaplasma* species; collected with each trapping method and from each habitat, suggesting habitat might not play a pivotal role in *Ehrlichia*/*Anaplasma* transmission. In Tennessee, ehrlichiosis cases begin in May and steadily increase with a peak of cases in September (TNDOH 2013), which aligns with *A. americanum* adult and nymph activity demonstrated here. Previously, we identified the entire study site as a suitable habitat for *A. americanum* and speculated that host movement may be the best indicator for this species (Trout Fryxell et al. 2015a). White-tailed deer are primary hosts for *A. americanum*, move widely across a variety of habitats, and are known *Ehrlichia* and *Anaplasma* reservoirs; they are likely fundamental in transmission (Childs and Paddock 2003, Paddock and Yabsley 2007, Harmon et al. 2015). Because *A. americanum* was the only tick species infected with *Ehrlichia*/*Anaplasma*, studying the dispersal patterns along with movement patterns of their infected reservoirs should provide a mechanism for pathogen surveillance and vector control.

Both *A. americanum* and *A. maculatum* had significantly higher *Rickettsia* infection rates compared to *D. variabilis*, which allowed comparison of trapping success and prevalence between habitat types. *Rickettsia* infection was not specific to habitat, collection period, or trapping method; however, data indicate that ticks encountered earlier in the season (May, June, and July) have a greater chance of being *Rickettsia* infected regardless of habitat (Tables 3 and 4). In this study, our data indicate that the risk of encountering a *Rickettsia*-positive tick is nearly three times greater in periods 1 (May 8–16), 2 (May 28–June 8), and 3 (June 25–July 6), compared to periods 4–6. The risk for encountering a *Rickettsia*-positive tick in period 6 (September 21–29) was minimal compared to the other periods. In addition, dragging (19.5% positive) was significantly more effective at collecting *Rickettsia*-infected ticks compared to the CO<sub>2</sub> trap (11.3% positive), indicating dragging is sufficient to obtain a representative of the questing and infected tick populations. Interestingly, the epidemiology of RMSF in human cases in Tennessee is bimodal with peaks in April and a second larger peak in July/August (TNDOH 2013), which corresponded with both adult and nymph *A. americanum* and *A. maculatum* adult abundance. Since *A. maculatum* are rare at the site, it would be beneficial to test the hypothesis that *A. americanum* nymphs are responsible for the second RMSF peak.

In this study, we found questing-infected ticks just before peak ehrlichiosis and RMSF human diagnoses in Tennessee (June–August), and this helps to explain the epidemiology of southern TBD cases. Tennessee TBD cases are likely

acquired (person was likely bit by an infected tick) during May–July (or earlier) and a majority of cases are diagnosed 1–2 weeks after (indicating immune response and bacterial pathogenesis occur shortly after bite) (TNDOH 2013). The identification of habitats and temporal periods of questing-infected ticks provides public health professionals with information to prevent transmission. For example, signs can be posted outside of state parks with dense conifer vegetation in June to remind visitors to avoid the area or take extra precautions against tick bites (*e.g.*, Deet on self, permethrin on clothing, tick checks). Similarly, this information can help physicians diagnose patients with a tick-borne illness if the patient indicates he/she/they visited a coniferous site in late June. Surveillance efforts can be concentrated during this period to identify specific land use areas with questing-infected ticks. Moreover, this information can also be used to develop and test management methods for the control of southern ticks as has been demonstrated for *I. scapularis* in the northeast (Ginsberg and Stafford 2005). Tick management methods include altering habitats conducive to infected ticks by applying acaricides, preventing hosts from entering the habitat, or any of the many other vegetation management options such as mowing, burning, and/or removing leaf litter (Ginsberg and Stafford 2005). Future research should focus on habitat/vector associations for each tick species, developing effective methods for collecting typically underrepresented tick species (such as *A. maculatum*), and creating surveillance programs for each species and pathogen.

Southeastern ehrlichiosis and rickettsiosis cases continue to be diagnosed and their vector distributions are expanding. In addition, the recent discovery of tick-borne viruses transmitted by *A. americanum* within the region, including Heartland virus (Savage et al. 2013) and Bourbon virus (Kosoy et al. 2015), warrants future studies to identify the true prevalence of these viruses along with their interactive effects with bacteria. Additional research needs to address the reasons for rickettsiosis cases in the southeast and identify the cause of increased fatalities in the Tennessee Valley compared to other regions with diagnoses but without high mortality (Adjemian et al. 2009). These reasons could be entomological (vector populations, frequencies, and/or densities), microbial (multiple *Rickettsia* coinfections, synergistic bacteria, and/or antagonistic bacteria), environmental (microbial, soil, host, abiotic, and/or landscape), and/or human mediated (medical doctors, patient behavior). Systematic studies examining biotic and abiotic parameters associated with southeastern ticks and TBDs should be conducted to develop a firm understanding of the interactions and dependencies between pathogens and hosts, the role of genetics and microbiomes, and how the ticks/pathogens/hosts partition or coexist in representative habitats. Field studies should be conducted at regional scales, allowing meaningful extrapolation to the southeastern ecological systems and human occupation patterns. Until *R. rickettsii* is identified in a tick vector or another pathogen is shown as a causative agent, the enigma surrounding TBDs in the south will remain (Dantas-Torres 2007, Cunha 2008, Raoult and Parola 2008, Adjemian et al. 2009).

## Conclusions

Several bacteria causing ehrlichiosis were identified in *A. americanum*, and SFGR bacteria of unknown or varying

pathogenicity were identified in *A. americanum* (“*Ca. R. amblyommii*”), *A. maculatum* (*R. parkeri* and *R. andersoniae*), and *D. variabilis* (*R. montanensis*). These tick species are encountered commonly, have overlapping distributions, and positive specimens tend to be collected May–July, which correlate with ehrlichiosis and rickettsiosis cases in Tennessee. This period should be used as a time for increased awareness, self-protection, and avoidance by the public; surveillance by public health professionals and physicians, and habitat and host management by land managers. In this study, we report several pathogenic *Ehrlichia/Anaplasma* species amplified from *A. americanum* and *R. parkeri* amplified from *A. maculatum*, the absence of *R. rickettsii*, and identify May–July as peak tick transmission in Tennessee.

## Acknowledgments

We thank Dr. Abelardo Moncayo with the Tennessee Department of Health for his contribution of *E. chaffeensis* and “*Ca. Rickettsia amblyommii*”-positive samples used as positive controls during the pathogen screening portion of our experiment. We also want to thank the reviewers for improving this manuscript, and Xiaocun Sun for help with additional statistical tests. We thank the board of trustees and the employees of Ames Plantation for accommodations, travel support, and so much more. In particular, we thank Larry Teague and James Morrow for their assistance with tick collection and vehicle extractions. Funding for this project was provided by the American Kennel Club Canine Health Foundation Acorn Grant (01894-A), Ames Foundation, and USDA Tennessee Hatch Project (TN00433). Fellowship support for B. Hendricks was provided by the University of Tennessee Institute of Agriculture’s Department of Entomology and Plant Pathology, for S. Mays by the J. Wallace and Katie Dean Graduate Fellowship at the University of Tennessee, and for K. Pompo by Merial Veterinary Scholar Program at the University of Tennessee College of Veterinary Medicine.

## Author Disclosure Statement

No competing financial interests exist.

## References

- Adjemian JZ, Krebs J, Mandel E, McQuiston J. Spatial clustering by disease severity among reported Rocky Mountain spotted fever cases in the United States, 2001–2005. *Am J Trop Med Hyg* 2009; 80:72–77.
- Ammerman NC, Swanson KI, Anderson JM, Schwartz TR, et al. Spotted-fever group *Rickettsia* in *Dermacentor variabilis*, Maryland. *Emerg Infect Dis* 2004; 10:1478–1481.
- Anderson BE, Dawson JE, Jones DC, Wilson KH. *Ehrlichia chaffeensis*, a new species associated with human ehrlichiosis. *J Clin Microbiol* 1991; 29:2838–2842.
- Anderson BE, Sims KG, Olson JG, Childs JE, et al. *Amblyomma americanum*: A potential vector of human ehrlichiosis. *Am J Trop Med Hyg* 1993; 49:239–244.
- Apperson CS, Engber B, Nicholson WL, Mead DG, et al. Tick-borne diseases in North Carolina: Is “*Rickettsia amblyommii*” a possible cause of rickettsiosis reported as Rocky Mountain spotted fever? *Vector-Borne Zoonotic Dis* 2008; 8:1–10.
- Berrada ZL, Goethert HK, Cunningham J, Telford SR III. *Rickettsia rickettsii* (Rickettsiales: Rickettsiaceae) in

- Amblyomma americanum* (Acari:Ixodidae) from Kansas. J Med Entomol 2011; 48:461–467.
- Breitschwerdt EB, Hegarty BC, Maggi RG, Lantos PM. *Rickettsia rickettsii* transmission by a lone star tick, North Carolina. Emerg Infect Dis 2011; 17:873–875.
- Buller RS, Arens M, Hmiel SP, Paddock CD, et al. *Ehrlichia ewingii*, a newly recognized agent of human ehrlichiosis. N Engl J Med 1999; 341:148–155.
- Carmichael JR, Fuerst PA. Molecular detection of *Rickettsia bellii*, *Rickettsia montanensis*, and *Rickettsia rickettsii* in *Dermacentor variabilis* ticks from nature. Vector-Borne Zoonotic Dis 2010; 10:111–115.
- (CDC) Ehrlichiosis. (2013a) Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases (NCEZID), Division of Vector-Borne Diseases. www.cdc.gov/ehrlichiosis Accessed May 31, 2017.
- (CDC) Rocky Mountain spotted fever. (2013b) Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases (NCEZID), Division of Vector-borne Diseases. www.cdc.gov/rmsf Accessed May 31, 2017.
- Chapman AS in collaboration with the Tickborne Rickettsial Diseases Working Group. Diagnosis and management of tick-borne rickettsial diseases: Rocky Mountain spotted fever, ehrlichiosis, and anaplasmosis—United States. CDC MMWR 2006; 55:1–27.
- Childs JE, Paddock CD. The ascendancy of *Amblyomma americanum* as a vector of pathogens affecting humans in the United States. Ann Rev Entomol 2003; 48:307–337.
- Cohen SB, Yabsley MJ, Freye JD, Dunlap BG, et al. Prevalence of *Ehrlichia chaffeensis* and *Ehrlichia ewingii* in ticks from Tennessee. Vector-Borne Zoonotic Dis 2010; 10:435–440.
- Cooley RA, Kohls GM. The genus *Amblyomma* (Ixodidae) in the United States. J Parasitol 1944; 30:77–111.
- Cunha BA. Clinical features of Rocky Mountain spotted fever. Lancet Infect Dis 2008; 8:143–144.
- Dantas-Torres F. Rocky Mountain spotted fever. Lancet Infect Dis 2007; 7:724–732.
- Eremeeva M, Yu X, Raoult D. Differentiation among spotted fever group rickettsiae species by analysis of fragment length polymorphism of PCR-amplified DNA. J Clin Microbiol 1994; 32:803.
- Eremeeva ME, Bosserman EA, Demma LJ, Zambrano ML, et al. Isolation and identification of *Rickettsia massiliae* from *Rhipicephalus sanguineus* ticks collected in Arizona. Appl Environ Microbiol 2006a; 72:85569–85577.
- Eremeeva ME, Bosserman E, Zambrano M, Demma L, et al. Molecular typing of novel *Rickettsia rickettsii* isolates from Arizona. Ann NY Acad Sci 2006b; 1078:573–577.
- Ewing SA, Dawson JE, Kocan AA, Barker RW, et al. Experimental transmission of *Ehrlichia chaffeensis* (Rickettsiales: Ehrlichiae) among white-tailed deer by *Amblyomma americanum* (Acari: Ixodidae). J Med Entomol 1995; 32:368–374.
- Falco RC, Fish D. A comparison of methods for sampling the deer tick, *Ixodes dammini*, in a Lyme disease endemic area. Experim App Acarol 1992; 14:165–173.
- Falco RC, McKenna DF, Daniels TJ, Nadelman, et al. Temporal relation between *Ixodes scapularis* abundance and risk for Lyme disease associated with erythema migrans. Am J Epidemiol 1999; 149:771–776.
- Fritzen CM, Huang J, Westby K, Freye JD, et al. Infection prevalence's of common tick-borne pathogens in adult lone star ticks (*Amblyomma americanum*) and American dog ticks (*Dermacentor variabilis*) in Kentucky. Am J Trop Med Hyg 2011; 85:718–723.
- Gaines D, Operario DJ, Stroup S, Stromdahl E, et al. *Ehrlichia* and spotted fever group rickettsiae surveillance in *Amblyomma americanum* in Virginia through use of a novel six-plex real-time PCR assay. Vector-Borne Zoonotic Dis 2014; 14:307–316.
- Ginsberg HS, Stafford KC. Management of ticks and tick-borne diseases. In: Goodman JL, Dennis DT, Sonenshine DE, ed. *Tick-borne Diseases of Humans*. Washington, DC: ASM Press, 2005: 65–86.
- Goddard J, Norment BR. Spotted fever group rickettsiae in the lone star tick, *Amblyomma americanum*. J Med Entomol 1986; 23:465–472.
- Goddard J, Varela-Stokes AS. Role of the lone star tick, *Amblyomma americanum* (L.) in human and animal diseases. Vet Parasitol 2009; 160:1–12.
- Harmon JR. Management of ticks and tick-borne disease in a Tennessee retirement community. M.Sci. Thesis, Institute of Agriculture, University of Tennessee, Knoxville. 2010.
- Harmon JR, Scott MC, Baker EM, Jones CJ, et al. Molecular identification of *Ehrlichia* species and host blood meal source in *Amblyomma americanum* L. from two locations in Tennessee, United States. Ticks Tick-Borne Dis 2015; 6: 246–252.
- Heise SR, Elshahed MS, Little SE. Bacterial diversity in *Amblyomma americanum* (Acari: Ixodidae) with a focus on members of the genus *Rickettsia*. J Med Entomol 2010; 47: 258–268.
- Hendricks B. Identification and Characterization of Peak Activity, Environmental Variables and Bacterial Pathogens in *A. americanum* L. at Ames Plantation, West Tennessee. Knoxville: M.Sci. Thesis, Institute of Agriculture, University of Tennessee, 2013.
- Henning TC, Orr JM, Smith JD, Arias JR, et al. Spotted fever group rickettsiae in multiple hard tick species from Fairfax county, Virginia. Vector-Borne Zoonotic Dis 2014; 14:482–485.
- Holland CJ, Ristic M, Huxsoll DL, Cole AI, et al. Adaptation of *Ehrlichia sennetsu* to canine blood monocytes: Preliminary structural and serological studies with cell culture-derived *Ehrlichia sennetsu*. Infect Immun 1985; 48:366–371.
- Hun L, Troyo A, Taylor L, Barbieri AM, et al. First report of the isolation and molecular characterization of *Rickettsia amblyommii* and *Rickettsia felis* in Central America. Vector-borne Zoonotic Dis 2011; 10:1395–1397.
- Keirans JE, Durden LA. Illustrated key to nymphs of the tick genus *Amblyomma* (Acari: Ixodidae) found in the United States. J Med Entomol 1998; 35:489–495.
- Keirans JE, Litwak TR. Pictorial key to the adults of hard ticks, family Ixodidae (Ixodida: Ixodoidea), East of the Mississippi river. J Med Entomol 1989; 26:435–448.
- Kosoy OI, Lambert J, Hawkinson DJ, Pastula DM, et al. Novel *Thogotovirus* species associated with febrile illness and death, United States, 2014. Emerg Infect Dis 2015; 21:760–764.
- Loftis AD, Mixson TR, Stromdahl EY, Yabsley MJ, et al. Geographic distribution and genetic diversity of the *Ehrlichia* sp. from Panola Mountain in *Amblyomma americanum*. BMC Infect Dis 2008; 8:54–61.
- Mays SE, Houston AE, Trout Fryxell RT. Comparison of novel and conventional methods of trapping ixodid ticks in the southeastern USA. Med and Vet Entomol 2016; 30:123–134.
- McCall CL, Curns AT, Rotz LD, Singleton Jr JA, et al. Fort Chaffee revisited: The epidemiology of tick-borne rickettsial and ehrlichial diseases at a natural focus. Vector Borne Zoonotic Dis 2001; 1:119–127.

- McDade JE, Newhouse VF. Natural history of *Rickettsia rickettsii*. *Ann Rev Microbiol* 1986; 40:287–309.
- Mixson TR, Campbell SR, Gill JG, Ginsberg HS, et al. Prevalence of *Ehrlichia*, *Borrelia*, and *Rickettsial* agents in *Amblyomma americanum* (Acari: Ixodidae) collected from nine states. *J Med Entomol* 2006; 43:1261–1268.
- Moncayo AC, Cohen SB, Fritzen CB, Huang E, et al. Absence of *Rickettsia rickettsii* and occurrence of other spotted fever group rickettsiae in ticks from Tennessee. *Am J Trop Med Hyg* 2010; 83:653–657.
- Monje LD, Nava S, Antoniazzi LR, Colombo VC, et al. In vitro isolation and infection intensity of *Rickettsia parkeri* in *Amblyomma triste* ticks from the Paraná River Delta region, Argentina. *Ticks Tick Borne Dis* 2014; 5:924–927.
- Ogrzewalska M, Literak I, Martins TF, Labruna MB. Rickettsial infections in ticks from wild birds in Paraguay. *Ticks Tick Borne Dis* 2014; 5:83–89.
- Paddock CD. *Rickettsia parkeri* as a paradigm for multiple causes of tick-borne spotted fever in the western hemisphere. *Ann NY Acad Sci* 2005; 1063:315–326.
- Paddock CD, Liddell AM, Storch GA. Other causes of tick-borne ehrlichiosis, including *Ehrlichia ewingii*. In: Goodman JL, Dennis DT, Sonenshine DE, ed. *Tick-Borne Diseases of Humans*. Washington, DC: ASM Press, 2005: 258–267.
- Paddock CD, Yabsley MJ. Ecological havoc, the rise of the white-tailed deer, and the emergence of *Amblyomma americanum*-associated zoonoses in the United States. *CTMI* 2007; 315:289–324.
- Paddock CD, Sumner JW, Comer JA, Zaki SR. *Rickettsia parkeri*: A newly recognized cause of spotted fever Rickettsiosis in the United States. *Clin Infect Dis* 2004; 38:812–813.
- Pagac BB, Miller MK, Mazzei MC, Nielsen DH, et al. *Rickettsia parkeri* and *Rickettsia montanensis*, Kentucky and Tennessee, USA. *Emerg Infect Dis* 2014; 20:1750–1752.
- Parola P, Paddock CD, Raoult D. Tick-borne rickettsioses around the world: Emerging diseases challenging old concepts. *Clin Microbiol Rev* 2005; 18:719–756.
- Raoult D, Parola P. Rocky Mountain spotted fever in the USA: A benign disease or a common diagnostic error? *Lancet Infect Dis* 2008; 8:587–589.
- Ricketts HT. The role of the wood-tick (*Dermacentor occidentalis*) in Rocky Mountain Spotted fever. *J Am Med Assoc* 1906; 49:24.
- Savage HM, Godsey Jr MS, Lambert A, Panella NA, et al. First detection of heartland virus (Bunyaviridae: *Phlebovirus*) from field collected arthropods. *Am J Trop Med Hyg* 2013; 89: 445–452.
- Sonenshine DE. The biology of tick vectors of human disease. In: Goodman JL, Dennis DT, Sonenshine DE, ed. *Tick-Borne Diseases of Humans*. Washington, DC: ASM Press, 2005: 12–36.
- Stromdahl EY, Hickling GJ. Beyond Lyme: Etiology of tick-borne human diseases with emphasis on the south-eastern United States. *Zoonoses Public Health* 2012; 59:48–64.
- Stromdahl EY, Jiang J, Vince M, Richards S. Infrequency of *Rickettsia rickettsii* in *Dermacentor variabilis* removed from humans, with comments on the role of other human-biting ticks associated with Spotted fever group Rickettsiae in the United States. *Vector-Borne Zoonotic Dis* 2011; 11:969–977.
- Sumner JW, Durden LA, Goddard J, Stromdahl EY, et al. Gulf Coast ticks (*Amblyomma maculatum*) and *Rickettsia parkeri*, United States. *Emerg Infect Dis* 2007; 13:751–753.
- Tabara K, Arai S, Kawabuchi T, Itagaki A, et al. Molecular survey of *Babesia microti*, *Ehrlichia* species, and *Candidatus Neorickettsia mikurensis* in wild rodents from Shimane Prefecture, Japan. *Microbiol Immunol* 2007; 51:359–367.
- Takano A, Ando S, Kishimoto T, Fujita H, et al. Presence of a novel *Ehrlichia* sp. in *Ixodes granulatus* found in Okinawa, Japan. *Microbiol Immunol* 2009; 53:101–106.
- Tate CM, Howerth EW, Mead DG, Dugan VG, et al. *Anaplasma odocoilei* sp. Nov. (family Anaplasmataceae) from white-tailed deer (*Odocoileus virginianus*). *Ticks Tick Borne Dis* 2013; 4:110–119.
- (TNDOH) Tennessee Department of Health. 2013. Communicable Disease Interactive Datae [Cited 09/02/13]. <http://health.state.tn.us/Ceds/WebAim> Accessed May 31, 2017.
- Trout RT, Steelman CD, Szalanski AL, Steelman CD. Rickettsiae in Gulf Coast ticks, Arkansas, USA. *Emerg Infect Dis* 2010; 16:830–832.
- Trout Fryxell RT, Moore JE, Collins MD, Kwon Y, et al. Habitat and vegetation variabilis are not enough when predicting tick populations in the southeastern United States. *PLoS One* 2015a; 12:e0144092.
- Trout Fryxell RT, Steelman CD, Szalanski AL, Billingsley PM, et al. Molecular detection of *Rickettsia* species within ticks (Acari: Ixodidae) collected from Arkansas United States. *J Med Entomol* 2015b; 52:500–508.
- Whitman TJ, Richards AL, Paddock CD, Tamminga CL, et al. *Rickettsia parkeri* infection after tick bite, Virginia. *Emerg Infect Dis* 2007; 13:334–336.
- Wilson JG, Kinzer DR, Sauer JR, Hair JA. Chemo-attraction in the lone star tick (Acarina: Ixodidae) response of different developmental stages to carbon dioxide administered via traps. *J Med Entomol* 1972; 9:245–252.

Address correspondence to:

Rebecca T. Trout Fryxell

Department of Entomology and Plant Pathology

University of Tennessee

370 Plant Biotechnology Building

2505 EJ Chapman Drive

Knoxville, TN 37996-4560

E-mail: rfryxell@utk.edu