

Types of Matrix :

Zero matrix =  $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}_{3 \times 3}$

Identity matrix =  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}_{3 \times 3}$   
 $n \times n$

Symmetric matrix =  $\begin{bmatrix} 2 & 3 & -1 \\ 3 & 0 & 6 \\ -1 & 6 & 5 \end{bmatrix}$

Diagonal matrix =  $\begin{bmatrix} 6 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 2 \end{bmatrix}$

Upper Triangular matrix =  $\begin{bmatrix} 6 & -1 & 5 \\ 0 & 4 & 2 \\ 0 & 0 & 2 \end{bmatrix}$

Lower Triangular matrix =  $\begin{bmatrix} 2 & 0 & 0 \\ 6 & 4 & 0 \\ -1 & 5 & 2 \end{bmatrix}$



Inverse of Matrix =  $A A^{-1} = A^{-1} A = I$  (Identity matrix)

Note: Determinant of square matrix should not be zero in inverse or else inverse doesn't exist.

Inverse is always for square matrix.

Transpose of matrix = for matrix A, transpose is " $A^T$ "

$$A \neq A^T$$

$$A = \begin{bmatrix} 6 & 2 & 1 \\ 3 & 5 & 9 \\ 7 & 4 & 8 \end{bmatrix}; \quad A^T = \begin{bmatrix} 6 & 3 & 7 \\ 2 & 5 & 4 \\ 1 & 9 & 8 \end{bmatrix}$$

If matrix is Symmetric & square then

$$\underline{\underline{A = A^T}}$$



# Determinants

$$A = \begin{bmatrix} m & n & o & p \\ q & r & s & t \\ u & v & w & x \\ y & z & a & b \end{bmatrix} \Rightarrow \text{Matrix}_{4 \times 4}$$

$$|A| = \begin{vmatrix} m & n & o & p \\ q & r & s & t \\ u & v & w & x \\ y & z & a & b \end{vmatrix} \Rightarrow \text{Determinant}_{4 \times 4}$$

~~$$= m \begin{vmatrix} r & s & t \\ u & v & w \end{vmatrix} - n \begin{vmatrix} q & s & t \\ u & w & x \end{vmatrix} + o \begin{vmatrix} q & r & t \\ u & v & x \end{vmatrix} - p \begin{vmatrix} q & r & s \\ u & v & w \end{vmatrix}$$~~

~~First method~~

$$= m \begin{vmatrix} r & s & t \\ u & v & w \\ z & a & b \end{vmatrix} - n \begin{vmatrix} q & s & t \\ u & w & x \\ y & a & b \end{vmatrix} + o \begin{vmatrix} q & r & t \\ u & v & x \\ y & z & b \end{vmatrix} - p \begin{vmatrix} q & r & s \\ u & v & w \\ y & z & a \end{vmatrix} = |A|$$

$$+ o \begin{vmatrix} q & r & t \\ u & v & x \\ y & z & b \end{vmatrix} - p \begin{vmatrix} q & r & s \\ u & v & w \\ y & z & a \end{vmatrix}$$



~~Second method~~

Finding determinant of matrix;

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}_{2 \times 2} \quad \text{for } 2 \times 2 \text{ matrix}$$

$$|A| = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$$

$$|A| = a_{11} \cdot a_{22} - a_{21} \cdot a_{12}$$

$$|A| = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}_{3 \times 3}$$

$$|A| = a_{11} \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$$



Find Inverse of matrix

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

method 1

$$A^{-1} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1}$$

$$A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

or

$$A^{-1} = \frac{\text{adj}(A)}{\det(A)} = \frac{\text{adj} \begin{bmatrix} 33 & 12 & 3 \\ -6 & 12 & -6 \\ 27 & -12 & 3 \end{bmatrix}}{\det(A)}$$

→ Transpose of minors matrix

$$\text{adj}(A) = \det \begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 2 & 9 \end{vmatrix} = -36$$

$$\therefore M_{1,1} = 33 \quad M_{3,1} = -3$$

$$\therefore M_{1,2} = -6 \quad M_{3,2} = -6$$

$$\therefore M_{1,3} = -27 \quad M_{3,3} = -3$$

$$\therefore M_{2,1} = 12$$

$$M_{2,2} = -12$$

$$M_{2,3} = -12$$

$$\therefore \begin{bmatrix} 33 & -6 & -27 \\ 12 & -12 & -12 \\ -3 & -6 & -3 \end{bmatrix}$$

Transpose this and result is  $\text{adj}(A)$



$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

Inverse is  $\Rightarrow$

$$A^{-1} = \frac{1}{|A|} \begin{bmatrix} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} & \begin{vmatrix} a_{13} & a_{12} \\ a_{33} & a_{32} \end{vmatrix} & \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix} \\ \begin{vmatrix} a_{23} & a_{21} \\ a_{33} & a_{31} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} & \begin{vmatrix} a_{13} & a_{11} \\ a_{23} & a_{21} \end{vmatrix} \\ \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} & \begin{vmatrix} a_{12} & a_{11} \\ a_{32} & a_{31} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} \end{bmatrix}$$

or

$$A = \begin{bmatrix} 2 & 1 \\ 7 & 4 \end{bmatrix}, \text{ Let } A = I A$$

$$\therefore \begin{bmatrix} 2 & 1 \\ 7 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} A$$

$$\therefore R_1 \rightarrow \left(\frac{1}{2}\right) R_1,$$

$$\begin{bmatrix} 1 & \frac{1}{2} \\ 7 & 4 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix} A$$



$$R_2 \rightarrow R_2 - 7R_1;$$

$$\begin{bmatrix} 1 & 1/2 \\ 0 & 1/2 \end{bmatrix} = \begin{bmatrix} 1/2 & 0 \\ -7/2 & 1 \end{bmatrix} A$$

$$R_2 \rightarrow 2R_2;$$

$$\begin{bmatrix} 1 & 1/2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1/2 & 0 \\ -7 & 2 \end{bmatrix} A$$

$$R_1 \rightarrow R_1 - (1/2)R_2;$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 4 & -1 \\ -7 & 2 \end{bmatrix} A$$

$$\therefore I = A^{-1}A$$

$$\therefore A^{-1} = \begin{bmatrix} 4 & -1 \\ -7 & 2 \end{bmatrix}$$

$$\text{Cofactor} = C_{ij} = (-1)^{i+j} M_{ij}$$

$M_{ij}$  = Minor, Transpose means rows converted to columns.



Properties:

$$\lambda(A+B) = \lambda A + \lambda B$$

$$\lambda A = A \lambda$$

$$AB \neq BA$$

$$(AB)C = A(BC)$$

$$A I_n = A = I_n A$$

for every  $A$  ( $|A| \neq 0$ ), there exists a  $B$

where  $AB = I_n = BA$ , i.e.  $B = A^{-1}$  or  $A = B^{-1}$ .

$$A' = A = A^T$$

$$(AB)' = A'B'$$

if  $A' = A$ , it is Symmetric

if  $A' = -A$ , it is skew Symmetric

$$|A| = |A'|$$

if two rows or columns are equal  $|A| = 0$