

The background of the cover features a large, orange wireframe sphere and a smaller wireframe torus (doughnut shape) positioned below it. These geometric shapes are superimposed on a photograph of a blue sea under a blue sky with scattered white clouds. The text is overlaid on semi-transparent colored rectangles: a dark green rectangle for the title, a dark red rectangle for the edition, and an orange rectangle for the authors.

4th Edition

Physical Chemistry

Silbey
Alberty
Bawendi

Physical Chemistry
Fourth Edition

Robert J. Silbey

*Class of 1942 Professor of Chemistry
Massachusetts Institute of Technology*

Robert A. Alberty

*Professor Emeritus of Chemistry
Massachusetts Institute of Technology*

Moungi G. Bawendi

*Professor of Chemistry
Massachusetts Institute of Technology*



John Wiley & Sons, Inc.

ACQUISITIONS EDITOR	Deborah Brennan
SENIOR PRODUCTION EDITOR	Patricia McFadden
SENIOR MARKETING MANAGER	Robert Smith
SENIOR DESIGNER	Kevin Murphy
NEW MEDIA EDITOR	Martin Batey

This book was set in 10/12 Times Roman by Publication Services, Inc. and printed and bound by Hamilton Printing. The cover was printed by Lehigh Press, Inc.
This book is printed on acid-free paper.Ⓢ

Copyright 2005 © John Wiley & Sons, Inc. All rights reserved.

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning or otherwise, except as permitted under Sections 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center Inc. 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 646-8600. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030-5774, (201) 748-6011, fax (201) 748-6008.

To order books or for customer service, call 1(800)-CALL-WILEY (225-5945).

PREFACE

The objective of this book is to make the concepts and methods of physical chemistry clear and interesting to students who have had a year of calculus and a year of physics. The underlying theory of chemical phenomena is complicated, and so it is a challenge to make the most important concepts and methods understandable to undergraduate students. However, these basic ideas are accessible to students, and they will find them useful whether they are chemistry majors, biologists, engineers, or earth scientists. The basic theory of chemistry is presented from the viewpoint of academic physical chemists, but many applications of physical chemistry to practical problems are described.

One of the important objectives of a course in physical chemistry is to learn how to solve numerical problems. The problems in physical chemistry help emphasize features in the underlying theory, and they illustrate practical applications.

There are two types of problems: problems that can be solved with a hand-held calculator and **COMPUTER PROBLEMS** that require a personal computer with a mathematical application installed. There are two sets of problems of the first type. The answers to problems in the first set are given in the back of the textbook, and worked-out solutions to these problems are given in the *Solutions Manual for Physical Chemistry*. The answers for the second set of problems are given in the *Solutions Manual*. In the two sets of problems that can be solved using hand-held calculators, some problems are marked with an icon to indicate that they may be more conveniently solved on a personal computer with a mathematical program. There are 170 **COMPUTER PROBLEMS** that require a personal computer with a mathematical application such as MathematicaTM, MathCadTM, MATLABTM, or MAPLETM installed. The recent development of these mathematical applications makes it possible to undertake problems that were previously too difficult or too time consuming. This is particularly true for two- and three-dimensional plots, integration and differentiation of complicated functions, and solving differential equations. The *Solutions Manual for Physical Chemistry* provides MathematicaTM programs and printouts for the **COMPUTER PROBLEMS**.

The MathematicaTM solutions of the 170 **COMPUTER PROBLEMS** in digital form are available on the web at <http://www.wiley.com/college/silbey>. They can be downloaded into a personal computer with MathematicaTM installed. Students

can obtain Mathematica at a reduced price from Wolfram Research, 100 Trade Center Drive, Champaign, Illinois, 61820-7237. A password is required and will be available in the *Solutions Manual*, along with further information about how to access the Mathematica solutions in digital form. Emphasis in the COMPUTER PROBLEMS has been put on problems that do not require complicated programming, but do make it possible for students to explore important topics more deeply. Suggestions are made as to how to vary parameters and how to apply these programs to other substances and systems. As an aid to showing how commands are used, there is an index in the *Solutions Manual* of the major commands used.

Mathematica™ plots are used in some 60 figures in the textbook. The legends for these figures indicate the COMPUTER PROBLEM where the program is given. These programs make it possible for students to explore changes in the ranges of variables in plots and to make calculations on other substances and systems.

One of the significant changes in the fourth edition is increased emphasis on the thermodynamics and kinetics of biochemical reactions, including the denaturation of proteins and nucleic acids. In this edition there is more discussion of the uses of statistical mechanics, nuclear magnetic relaxation, nano science, and oscillating chemical reactions.

This edition has 32 new problems that can be solved with a hand-held calculator and 35 new problems that require a computer with a mathematical application. There are 34 new figures and eight new tables.

Because the number of credits in physical chemistry courses, and therefore the need for more advanced material, varies at different universities and colleges, more topics have been included in this edition than can be covered in most courses.

The Appendix provides an alphabetical list of symbols for physical quantities and their units. The use of nomenclature and units is uniform throughout the book. SI (Système International d'Unités) units are used because of their advantage as a coherent system of units. That means that when SI units are used with all of the physical quantities in a calculation, the result comes out in SI units without having to introduce numerical factors. The underlying unity of science is emphasized by the use of seven base units to represent all physical quantities.

HISTORY

Outlines of Theoretical Chemistry, as it was then entitled, was written in 1913 by Frederick Getman, who carried it through 1927 in four editions. The next four editions were written by Farrington Daniels. In 1955, Robert Alberty joined Farrington Daniels. At that time, the name of the book was changed to *Physical Chemistry*, and the numbering of the editions was started over. The collaboration ended in 1972 when Farrington Daniels died. It is remarkable that this textbook traces its origins back 91 years.

Over the years this book has profited tremendously from the advice of physical chemists all over the world. Many physical chemists who care how their subject is presented have written to us with their comments, and we hope that will continue. We are especially indebted to colleagues at MIT who have reviewed various sections and given us the benefit of advice. These include Sylvia T. Ceyer, Robert W. Field, Carl W. Garland, Mario Molina, Keith Nelson, and Irwin Oppenheim.

The following individuals made very useful suggestions as to how to improve this fourth edition: Kenneth G. Brown (Old Dominion University), Thandi Buthelez (Western Kentucky University), Susan Collins (California State University Northridge), John Gold (East Straudsbury University), Keith J. Stine (University of Missouri–St. Louis), Ronald J. Terry (Western Illinois University), and Worth E. Vaughan (University of Wisconsin, Madison). We are also indebted to reviewers of earlier editions and to people who wrote us about the third edition.

The following individuals made very useful suggestions as to how to improve the Mathematica™ solutions to COMPUTER PROBLEMS: Ian Brooks (Wolfram Research), Carl W. David (U. Connecticut), Robert N. Goldberg (NIST), Mark R. Hoffmann (University of North Dakota), Andre Kuzniarek (Wolfram Research), W. Martin McClain (Wayne State University), Kathryn Tomasson (University of North Dakota), and Worth E. Vaughan (University of Wisconsin, Madison).

We are indebted to our editor Deborah Brennan and to Catherine Donovan and Jennifer Yee at Wiley for their help in the production of the book and the solutions manual. We are also indebted to Martin Batey for making available the web site, and to many others at Wiley who were involved in the production of this fourth edition.

Cambridge, Massachusetts
January 2004

Robert J. Silbey
Robert A. Alberty
Moungi G. Bawendi

CONTENTS

PART ONE **THERMODYNAMICS**

1. Zeroth Law of Thermodynamics and Equations of State 3
2. First Law of Thermodynamics 31
3. Second and Third Laws of Thermodynamics 74
4. Fundamental Equations of Thermodynamics 102
5. Chemical Equilibrium 132
6. Phase Equilibrium 177
7. Electrochemical Equilibrium 218
8. Thermodynamics of Biochemical Reactions 254

PART TWO **QUANTUM CHEMISTRY**

9. Quantum Theory 295
10. Atomic Structure 348
11. Molecular Electronic Structure 396
12. Symmetry 437
13. Rotational and Vibrational Spectroscopy 458
14. Electronic Spectroscopy of Molecules 502
15. Magnetic Resonance Spectroscopy 537
16. Statistical Mechanics 568

PART THREE
KINETICS

- 17. Kinetic Theory of Gases 613
- 18. Experimental Kinetics and Gas Reactions 641
- 19. Chemical Dynamics and Photochemistry 686
- 20. Kinetics in the Liquid Phase 724

PART FOUR
MACROSCOPIC AND MICROSCOPIC STRUCTURES

- 21. Macromolecules 763
- 22. Electric and Magnetic Properties of Molecules 786
- 23. Solid-State Chemistry 803
- 24. Surface Dynamics 840

APPENDIX

- A. Physical Quantities and Units 863
- B. Values of Physical Constants 867
- C. Tables of Physical Chemical Data 868
- D. Mathematical Relations 884
- E. Greek Alphabet 897
- F. Useful Information on the Web 898
- G. Symbols for Physical Quantities and Their SI Units 899
- H. Answers to the First Set of Problems 912

INDEX 933

This page intentionally left blank

Thermodynamics

Thermodynamics deals with the interconversion of various kinds of energy and the changes in physical properties that are involved. Thermodynamics is concerned with equilibrium states of matter and has nothing to do with time. Even so, it is one of the most powerful tools of physical chemistry; because of its importance, the first part of this book is devoted to it. The first law of thermodynamics deals with the amount of work that can be done by a chemical or physical process and the amount of heat that is absorbed or evolved. On the basis of the first law it is possible to build up tables of enthalpies of formation that may be used to calculate enthalpy changes for reactions that have not yet been studied. With information on heat capacities of reactants and products also available, it is possible to calculate the heat of a reaction at a temperature where it has not previously been studied.

The second law of thermodynamics deals with the natural direction of processes and the question of whether a given chemical reaction can occur by itself. The second law was formulated initially in terms of the efficiencies of heat engines, but it also leads to the definition of entropy, which is important in determining the direction of chemical change. The second law provides the basis for the definition of the equilibrium constant for a chemical reaction. It provides an answer to the question, "To what extent will this particular reaction go before equilibrium is reached?" It also provides the basis for reliable predictions of the effects of temperature, pressure, and concentration on chemical and physical equilibrium. The third law provides the basis for calculating equilibrium constants from calorimetric measurements only. This is an illustration of the way in which thermodynamics interrelates apparently unrelated measurements on systems at equilibrium.

After discussing the laws of thermodynamics and the various physical quantities involved, our first applications will be to the quantitative treatment of chemical equilibria. These methods are then applied to equilibria between different phases. This provides the basis for the quantitative treatment of distillation and for the interpretation of phase changes in mixtures of solids. Then thermodynamics is applied to electrochemical cells and biochemical reactions.

This page intentionally left blank



Zeroth Law of Thermodynamics and Equations of State

- 1.1** State of a System
- 1.2** The Zeroth Law of Thermodynamics
- 1.3** The Ideal Gas Temperature Scale
- 1.4** Ideal Gas Mixtures and Dalton's Law
- 1.5** Real Gases and the Virial Equation
- 1.6** P - \bar{V} - T Surface for a One-Component System
- 1.7** Critical Phenomena
- 1.8** The van der Waals Equation
- 1.9** Description of the State of a System
without Chemical Reactions
- 1.10** Partial Molar Properties
- 1.11** Special Topic: Barometric Formula

Physical chemistry is concerned with understanding the quantitative aspects of chemical phenomena. To introduce physical chemistry we will start with the most accessible properties of matter—those that can readily be measured in the laboratory. The simplest of these are the properties of matter at equilibrium. Thermodynamics deals with the properties of systems at equilibrium, such as temperature, pressure, volume, and amounts of species; but it also deals with work done on a system and heat absorbed by a system, which are not properties of the system but measures of changes. The amazing thing is that the thermodynamic properties of systems at equilibrium obey all the rules of calculus and are therefore interrelated. The principle involved in defining temperature was not recognized until the establishment of the first and second laws of thermodynamics, and so it is referred to as the zeroth law. This leads to a discussion of the thermodynamic properties of gases and liquids. After discussing the ideal gas, we consider the behavior of real gases. The thermodynamic properties of a gas or liquid are represented by an equation of state, such as the virial equation or the van der Waals equation. The latter has the advantage that it provides a description of the critical region, but much more complicated equations are required to provide an accurate quantitative description.

1.1 STATE OF A SYSTEM

A thermodynamic system is that part of the physical universe that is under consideration. A system is separated from the rest of the universe by a real or idealized **boundary**. The part of the universe outside the boundary of the system is referred to as the **surroundings**, as illustrated in Fig. 1.1. The boundary between the system and its surroundings may have certain real or idealized characteristics. For example, the boundary may conduct heat or be a perfect insulator. The boundary may be rigid or it may be movable so that it can be used to apply a specified pressure. The boundary may be impermeable to the transfer of matter between the system and its surroundings, or it may be permeable to a specified species. In other words, matter and heat may be transferred between system and surroundings, and the surroundings may do work on the system, or vice versa. If the boundary around a system prevents interaction of the system with its surroundings, the system is called an **isolated** system.

If matter can be transferred from the surroundings to the system, or vice versa, the system is referred to as an **open** system; otherwise, it is a **closed** system.

When a system is under discussion it must be described precisely. A system is **homogeneous** if its properties are uniform throughout; such a system consists of a single phase. If a system contains more than one phase, it is **heterogeneous**. A simple example of a two-phase system is liquid water in equilibrium with ice. Water can also exist as a three-phase system: liquid, ice, and vapor, all in equilibrium.

Experience has shown that the macroscopic state of a system at equilibrium can be specified by the values of a small number of macroscopic variables. These variables, which include, for example, temperature T , pressure P , and volume V , are referred to as **state variables** or **thermodynamic variables**. They are called state variables because they specify the state of a system. Two samples of a substance that have the same state variables are said to be in the same state. It is remarkable that the state of a homogeneous system at equilibrium can be specified by so few variables. When a sufficient number of state variables are specified, all of the other properties of the system are fixed. It is even more remarkable that these state variables follow all of the rules of calculus; that is, they can be treated as mathematical functions that can be differentiated and integrated. Thermodynamics leads to the definition of additional properties, such as internal energy and entropy, that can also be used to describe the state of a system, and are themselves state variables.

The thermodynamic state of a specified amount of a pure substance in the fluid state can be described by specifying properties such as temperature T , pressure P , and volume V . But experience has shown that only two of these three properties have to be specified when the amount of pure substance is fixed. If T and P , or P and V , or T and V are specified, all the other thermodynamic properties (including those that will be introduced later) are fixed and the system is at equilibrium. More properties have to be specified to describe the thermodynamic state of a homogeneous mixture of different species.

Note that the description of the microscopic state of a system containing many molecules requires the specification of a very large number of variables. For example, to describe the microscopic state of a system using classical mechanics, we would have to give the three coordinates and three components of the momentum of each molecule, plus information about its vibrational and rotational motion. For one mole of gas molecules, this would mean more than 6×10^{23} numbers. An




Figure 1.1 (a) A system is separated from its surroundings by a boundary, real or idealized. (b) As a simplification we can imagine the system to be separated from the surroundings by a single wall that may be an insulator or a heat conductor. Later, in Section 6.7 and Section 8.3 (see Fig. 8.6), we will consider semipermeable boundaries so that the system is open to the transfer of matter.

important thing to notice is that we can use a small number of state variables to describe the equilibrium thermodynamic state of a system that is too complicated to describe in a microscopic way.

Thermodynamic variables are either intensive or extensive. **Intensive variables** are independent of the size of the system; examples are pressure, density, and temperature. **Extensive variables** do depend on the size of the system and double if the system is duplicated and added to itself; examples are volume, mass, internal energy, and entropy. Note that the ratio of two extensive variables is an intensive variable; density is an example. Thus we can talk about the **intensive state of the system**, which is described by intensive variables, or the **extensive state of a system**, which is described by intensive variables plus at least one extensive variable. The intensive state of the gas helium is described by specifying its pressure and density. The extensive state of a certain amount of helium is described by specifying the amount, the pressure, and the density; the extensive state of one mole of helium might be represented by $1 \text{ mol He}(P, \rho)$, where P and ρ represent the pressure and density, respectively. We can generalize this by saying that the intensive state of a pure substance in the fluid state is specified by $N_s + 1$ variables, where N_s is the number of different kinds of species in the system. The extensive state is specified by $N_s + 2$ variables, one of which has to be extensive.

In chemistry it is generally more useful to express the size of a system in terms of the amount of substance it contains, rather than its mass. **The amount of substance n is the number of entities (atoms, molecules, ions, electrons, or specified groups of such particles) expressed in terms of moles.** If a system contains N molecules, the amount of substance $n = N/N_A$, where N_A is the Avogadro constant ($6.022 \times 10^{23} \text{ mol}^{-1}$). The ratio of the volume V to the amount of substance is referred to as the molar volume: $\bar{V} = V/n$. The volume V is expressed in SI units of m^3 , and the molar volume \bar{V} is expressed in SI units of $\text{m}^3 \text{ mol}^{-1}$. We will use the overbar regularly to indicate molar thermodynamic quantities.

Comment:

Since this is our first use of physical quantities, we should note that the value of a physical quantity is equal to the product of a numerical factor and a unit:

$$\text{physical quantity} = \text{numerical value} \times \text{unit}$$

The values of all physical quantities can be expressed in terms of SI base units (see Appendix A). However, some physical quantities are dimensionless, and so the symbol for the SI unit is taken as 1 because this is what you get when units cancel. Note that, in print, physical quantities are represented by italic type and units are represented by roman type.

When a system is in a certain state with its properties independent of time and having no fluxes (e.g., no heat flowing through the system), then the system is said to be at **equilibrium**. When a thermodynamic system is at equilibrium its state is defined entirely by the state variables, and **not by the history of the system**. By history of the system, we mean the previous conditions under which it has existed.

Since the state of a system at equilibrium can be specified by a small number of state variables, it should be possible to express the value of a variable that has not been specified as a function of the values of other variables that have been specified. The simplest example of this is the ideal gas law.




Figure 1.2 (a) The pressure exerted by the atmosphere on the surface of mercury in a cup is given by $P = h\rho g$ (see Example 1.1). (b) The pressure of a system is given by the same equation when a closed-end manometer is used.

For some systems, more than two intensive variables must be stated to specify the state of the system. If there is more than one species, the composition has to be given. If a liquid system is in the form of small droplets, the surface area has to be given. If the system is in an electric or magnetic field, this may have an effect on its properties, and then the electric field strength and magnetic field strength become state variables. We will generally ignore the effect of the earth's gravitational field on a system, although this can be important, as we will see in the special topic at the end of this chapter. Note that the properties used to describe the state of a system must be independent; otherwise they are redundant. Independent properties are separately controllable by the investigator.

The pressure of the atmosphere is measured with a barometer, as shown in Fig. 1.2a, and the pressure of a gaseous system is measured with a closed-end manometer, as shown in Fig. 1.2b.

1.2 THE ZEROTH LAW OF THERMODYNAMICS

Although we all have a commonsense notion of what temperature is, we must define it very carefully so that it is a useful concept in thermodynamics. If two closed systems with fixed volumes are brought together so that they are in thermal contact, changes may take place in the properties of both. Eventually a state is reached in which there is no further change, and this is the state of **thermal equilibrium**. In this state, the two systems have the same temperature. Thus, we can readily determine whether two systems are at the same temperature by bringing them into thermal contact and seeing whether observable changes take place in the properties of either system. If no change occurs, the systems are at the same temperature.

Now let us consider three systems, A, B, and C, as shown in Fig. 1.3. It is an experimental fact that if system A is in thermal equilibrium with system C, and system B is also in thermal equilibrium with system C, then A and B are in thermal equilibrium with each other. It is not obvious that this should be true, and so this empirical fact is referred to as the **zeroth law of thermodynamics**.

To see how the zeroth law leads to the definition of a temperature scale, we need to consider thermal equilibrium between systems A, B, and C in more detail. Assume that A, B, and C each consist of a certain mass of a different fluid. We use the word **fluid** to mean either a gas or a compressible liquid. Our experience is that if the volume of one of these systems is held constant, its pressure may vary over a range of values, and if the pressure is held constant, its volume may vary over a range of values. Thus, the pressure and the volume are independent thermodynamic variables. Furthermore, suppose that the experience with these systems is that their intensive states are specified completely when the pressure and volume are specified. That is, when one of the systems reaches equilibrium at a certain pressure and volume, all of its macroscopic properties have certain characteristic values. It is quite remarkable and fortunate that the macroscopic state of a given mass of fluid of a given composition can be fixed by specifying only the pressure and the volume.*

If there are further constraints on the system, there will be a smaller number of independent variables. An example of an additional constraint is thermal

*This is not true for water in the neighborhood of 4 °C, but the state is specified by giving the temperature and the volume or the temperature and the pressure. See Section 6.1.

equilibrium with another system. Experience shows that if a fluid is in thermal equilibrium with another system, it has only one independent variable. In other words, if we set the pressure of system A at a particular value P_A , we find that there is thermal equilibrium with system C, in a specified state, only at a particular value of V_A . Thus, system A in thermal equilibrium with system C is characterized by a **single** independent variable, pressure or volume; one or the other can be set arbitrarily, but not both. The plot of all the values of P_A and V_A for which there is equilibrium with system C is called an **isotherm**. Figure 1.4 gives this isotherm, which we label Θ_1 . Since system A is in thermal equilibrium with system C at any P_A , V_A on the isotherm, we can say that each of the pairs P_A , V_A on this isotherm corresponds with the same temperature Θ_1 .

When heat is added to system C and the experiment is repeated, a different isotherm is obtained for system A. In Fig. 1.4, the isotherm for the second experiment is labeled Θ_2 . If still more heat is added to system C and the experiment is repeated again, the isotherm labeled Θ_3 is obtained.

Figure 1.4 illustrates Boyle's law, which states that $PV = \text{constant}$ for a specified amount of gas at a specified temperature. Experimentally, this is strictly true only in the limit of zero pressure. Charles and Gay-Lussac found that the volume of a gas varies linearly with the temperature at specified pressure when the temperature is measured with a mercury in glass thermometer, for example. Since it would be preferable to have a temperature scale that is independent of the properties of particular materials like mercury and glass, it is better to say that the ratio of the P_2V_2 product at temperature Θ_2 to P_1V_1 at temperature Θ_1 depends only on the two temperatures:

$$\frac{P_2V_2}{P_1V_1} = \phi(\Theta_1, \Theta_2) \quad (1.1)$$

where ϕ is an unspecified function. The simplest thing to do is to take the ratio of the PV products to be equal to the ratio of the temperatures, thus defining






Figure 1.4 Isotherms for fluid A. This plot, which is for a hypothetical fluid, might look quite different for some other fluid.



If A and C are in thermal equilibrium, and B and C are in thermal equilibrium, then



A and B will be found to be in thermal equilibrium when connected by a heat conductor.

Figure 1.3 The zeroth law of thermodynamics is concerned with thermal equilibrium between three bodies.

a temperature scale:

$$\frac{P_2 V_2}{P_1 V_1} = \frac{T_2}{T_1} \quad \text{or} \quad \frac{P_2 V_2}{T_2} = \frac{P_1 V_1}{T_1} \quad (1.2)$$

Here we have introduced a new symbol T for the temperature because we have made a specific assumption about the function ϕ . Equations 1.1 and 1.2 are exact only in the limit of zero pressure, and so T is referred to as the ideal gas temperature.

Since, according to equation 1.2, PV/T is a constant for a fixed mass of gas and since V is an extensive property,

$$PV/T = nR \quad (1.3)$$

where n is the amount of gas and R is referred to as the **gas constant**. Equation 1.3 is called the ideal gas **equation of state**. An equation of state is a relation between the thermodynamic properties of a substance at equilibrium.

1.3 THE IDEAL GAS TEMPERATURE SCALE

The ideal gas temperature scale can be defined more carefully by taking the temperature T to be proportional to $P\bar{V} = PV/n$ in the limit of zero pressure. Since different gases give slightly different scales when the pressure is about one bar ($1 \text{ bar} = 10^5 \text{ pascal} = 10^5 \text{ Pa} = 10^5 \text{ N m}^{-2}$), it is necessary to use the limit of the $P\bar{V}$ product as the pressure approaches zero. When this is done, all gases yield the same temperature scale. We speak of gases under this limiting condition as **ideal**. Thus, the **ideal gas temperature** T is defined by

$$T = \lim_{P \rightarrow 0} (P\bar{V}/R) \quad (1.4)$$

The proportionality constant is called the gas constant R . The unit of thermodynamic temperature, 1 kelvin or 1 K, is defined as the fraction $1/273.16$ of the temperature of the triple point of water.* Thus, the temperature of an equilibrium system consisting of liquid water, ice, and water vapor is 273.16 K. The temperature 0 K is called absolute zero. According to the current best measurements, the freezing point of water at 1 atmosphere (101 325 Pa; see below) is 273.15 K, and the boiling point at 1 atmosphere is 373.12 K; however, these are experimental values and may be determined more accurately in the future. The Celsius scale t is formally defined by

$$t/^{\circ}\text{C} = T/\text{K} - 273.15 \quad (1.5)$$

The reason for writing the equation in this way is that temperature T on the Kelvin scale has the unit K, and temperature t on the Celsius scale has the unit $^{\circ}\text{C}$, which need to be divided out before temperatures on the two scales are compared. In Fig. 1.5, the molar volume of an ideal gas is plotted versus the Celsius temperature t at two pressures.




Figure 1.5 Plots of \bar{V} versus temperature for a given amount of a real gas at two low pressures P_1 and P_2 , as given by Gay-Lussac's law.

*The triple point of water is the temperature and pressure at which ice, liquid, and vapor are in equilibrium with each other in the absence of air. The pressure at the triple point is 611 Pa. The freezing point in the presence of air at 1 atm is 0.0100°C lower because (1) the solubility of air in liquid water at 1 atm (101 325 Pa) is sufficient to lower the freezing point 0.0024°C (Section 6.7), and (2) the increase of pressure from 611 to 101 325 Pa lowers the freezing point 0.0075°C , as shown in Example 6.2. Thus, the ice point is at 273.15 K.

We will find later that the ideal gas temperature scale is identical with one based on the second law of thermodynamics, which is independent of the properties of any particular substance (see Section 3.9). In Chapter 16 the ideal gas temperature scale will be identified with that which arises in statistical mechanics.

The gas constant R can be expressed in various units, but we will emphasize the use of SI units. The SI unit of **pressure** (P) is the pascal, Pa, which is the pressure produced by a force of 1 N on an area of 1 m². In addition to using the prefixes listed in the back cover of the book to express larger and smaller pressures, it is convenient to have a unit that is approximately equal to the atmospheric pressure. This unit is the bar, which is 10⁵ Pa. Earlier the atmosphere, which is defined as 101 325 Pa, had been used as a unit of pressure.

Example 1.1 *Express one atmosphere pressure in SI units*

Calculate the pressure of the earth's atmosphere at a point where the barometer reads 76 cm of mercury at 0°C and the acceleration of gravity g is 9.806 65 m s⁻². The density of mercury at 0°C is 13.5951 g cm⁻³, or 13.5951 × 10³ kg m⁻³.

Pressure P is force f divided by area A :

$$P = f/A$$

The force exerted by a column of air over an area A is equal to the mass m of mercury in a vertical column with a cross section A times the acceleration of gravity g :

$$f = mg$$

The mass of mercury raised above the flat surface in Fig. 1.2a is ρAh so that

$$f = \rho Ahg$$

Thus, the pressure of the atmosphere is

$$P = h\rho g$$

If h , ρ , and g are expressed in SI units, the pressure P is expressed in pascals. Thus, the pressure of a standard atmosphere may be expressed in SI units as follows:

$$\begin{aligned} 1 \text{ atm} &= (0.76 \text{ m})(13.5951 \times 10^3 \text{ kg m}^{-3})(9.806 65 \text{ m s}^{-2}) \\ &= 101 325 \text{ N m}^{-2} = 101 325 \text{ Pa} = 1.013 25 \text{ bar} \end{aligned}$$

This equality is expressed by the conversion factor 1.013 25 bar atm⁻¹.

To determine the value of the gas constant we also need the definition of a mole. A **mole** is the amount of substance that has as many atoms or molecules as 0.012 kg (exactly) of ¹²C. The **molar mass** M of a substance is the mass divided by the amount of substance n , and so its SI unit is kg mol⁻¹. Molar masses can also be expressed in g mol⁻¹, but it is important to remember that in making calculations in which all other quantities are expressed in SI units, the molar mass must be expressed in kg mol⁻¹. The molar mass M is related to the molecular mass m by $M = N_A m$, where N_A is the **Avogadro constant** and m is the mass of a single molecule.

Until 1986 the recommended value of the gas constant was based on measurements of the molar volumes of oxygen and nitrogen at low pressures. The accuracy

of such measurements is limited by problems of sorption of gas on the walls of the glass vessels used. In 1986 the recommended value* of the gas constant

$$R = 8.314\,51\,\text{J K}^{-1}\,\text{mol}^{-1} \quad (1.6)$$

was based on measurements of the speed of sound in argon. The equation used is discussed in Section 17.4. Since pressure is force per unit area, the product of pressure and volume has the dimensions of force times distance, which is work or energy. Thus, the gas constant is obtained in joules if pressure and volume are expressed in pascals and cubic meters; note that $1\,\text{J} = 1\,\text{Pa m}^3$.

Example 1.2 Express the gas constant in various units

Calculate the value of R in $\text{cal K}^{-1}\,\text{mol}^{-1}$, $\text{L bar K}^{-1}\,\text{mol}^{-1}$, and $\text{L atm K}^{-1}\,\text{mol}^{-1}$.

Since the calorie is defined as $4.184\,\text{J}$,

$$\begin{aligned} R &= 8.314\,51\,\text{J K}^{-1}\,\text{mol}^{-1}/4.184\,\text{J cal}^{-1} \\ &= 1.987\,22\,\text{cal K}^{-1}\,\text{mol}^{-1} \end{aligned}$$

Since the liter is $10^{-3}\,\text{m}^3$ and the bar is $10^5\,\text{Pa}$,

$$\begin{aligned} R &= (8.314\,51\,\text{Pa m}^3\,\text{K}^{-1}\,\text{mol}^{-1})(10^3\,\text{L m}^{-3})(10^{-5}\,\text{bar Pa}^{-1}) \\ &= 0.083\,145\,1\,\text{L bar K}^{-1}\,\text{mol}^{-1} \end{aligned}$$

Since $1\,\text{atm}$ is $1.013\,25\,\text{bar}$,

$$\begin{aligned} R &= (0.083\,145\,1\,\text{L bar K}^{-1}\,\text{mol}^{-1})/(1.013\,25\,\text{bar atm}^{-1}) \\ &= 0.082\,057\,8\,\text{L atm K}^{-1}\,\text{mol}^{-1} \end{aligned}$$

1.4 IDEAL GAS MIXTURES AND DALTON'S LAW

Equation 1.3 applies to a mixture of ideal gases as well as a pure gas, when n is the total amount of gas. Since $n = n_1 + n_2 + \cdots$, then

$$\begin{aligned} P &= (n_1 + n_2 + \cdots)RT/V \\ &= n_1RT/V + n_2RT/V + \cdots \\ &= P_1 + P_2 + \cdots = \sum_i P_i \end{aligned} \quad (1.7)$$

where P_1 is the partial pressure of species 1. Thus, the total pressure of an ideal gas mixture is equal to the sum of the partial pressures of the individual gases; this is **Dalton's law**. The partial pressure of a gas in an ideal gas mixture is the pressure that it would exert alone in the total volume at the temperature of the mixture:

$$P_i = n_i RT/V \quad (1.8)$$

A useful form of this equation is obtained by replacing RT/V by P/n :

$$P_i = n_i P/n = y_i P \quad (1.9)$$

*E. R. Cohen and B. N. Taylor, The 1986 Adjustment of the Fundamental Physical Constants, *CODATA Bull.* **63**:1 (1986); *J. Phys. Chem. Ref. Data* **17**:1795 (1988).