



Name **Mr. MAHESH KAVITISHETTY**
Age / Gender : 29.2 Year(s)/ Male
Contact No. : +919967833785
Address JJ H Mumbai..
Pin code : 400008

VID No. : 250138509972839
PID No. : P14525547224709
Referred by : DR.JJ HOSPITAL
Registered On : 17/04/2026 4:03 PM
Collected On : 17/04/2026 4:15 PM
Reported On : 17/04/2026 11:11 PM

SUMMARY REPORT

Investigation Outside Reference Range (Abnormal)

Investigation	Observed Value	Unit	Biological Reference Interval
<u>PT (Prothrombin Time)</u>			
(Citratd Plasma)			
Prothrombin Time	38.1	C sec	9.4-12.5 Kindly note change in reference ranges
Control (MNPT)	11.2	sec	= 11.0
PT(INR) Value	3.44		Normal Population : 0.8 - 1.2 Standard Therapy: 2.0-3.0 High Dose Therapy: 3.0-4.5
<u>Lipid Profile - 2 (Mini - Fasting)</u>			
(Serum Homogeneous Enzymatic colorimetric Assay)			
HDL Cholesterol Direct	35.2	mg/dL	> 40

MEDICAL LABORATORY REPORT

"C" represents critical results

Page 1 of 4



Name : Mr. MAHESH KAVITISHETTY
Age / Gender : 29.2 Year(s)/ Male
Contact No. : +919967833785
Address : JJ H Mumbai..
Pin code : 400008

VID No. : 250138509972839
PID No. : P14525547224709
Referred by : DR.JJ HOSPITAL
Registered On : 17/04/2026 4:03PM
Collected On : 17/04/2026 4:15PM
Reported On : 17/04/2026 11:11P

Investigation	Observed Value	Unit	Biological Reference Interval
<u>Lipid Profile - 2 (Mini - Fasting)</u>			
Cholesterol Total, Serum (Serum,Enzymatic colorimetric Assay)	120	mg/dL	Desirable: < 200 Borderline High: 200-240 High: >= 240
Triglycerides, Serum (Serum,Enzymatic)	71.8	mg/dL	Normal: < 150 Borderline High: 150-199 High: 200-499 Very High: >= 500
HDL Cholesterol Direct (Serum,Homogeneous Enzymatic colorimetric Assay)	35.2	mg/dL	> 40
Non HDL Cholesterol (Serum,Calculated)	84.8	mg/dL	Optimal:<130 ,Desirable:130-159 ,Borderline high:159-189 ,High:189-220 ,Very High:>=220
LDL Cholesterol (Serum,Calculated)	70	mg/dL	Optimal:<100 ,Near Optimal:100-129 ,Borderline high:130-159 ,High:160-189 ,Very High:>=190
VLDL Cholesterol (Serum,Calculated)	14	mg/dL	6-38
LDL/HDL Ratio (Serum,Calculated)	2		2.5-3.5
Cholestrol / HDL Ratio (Serum,Calculated)	3		3.5-5.0

Note: Reference Interval as per National Cholesterol Education Program (NCEP) Adult Treatment Panel III Report.
 VLDL,CHOL/HDL RATIO,LDL/HDL RATIO,LDL Cholesterol,serum,Non HDL Colesterol are calculated parameters

Dr. ANISH BHATT
 MD (Pathology)
 Consultant Pathologist
 Reg No. 2014041630

MEDICAL LABORATORY REPORT



Name **Mr. MAHESH KAVITISHETTY**
 Age / Gender : 29.2 Year(s)/ Male
 Contact No. : +919967833785
 Address JJ H Mumbai..
 Pin code : 400008

VID No. : 250138509972839
 PID No. : P14525547224709
 Referred by : DR.JJ HOSPITAL
 Registered On : 17/04/2026 4:03PM
 Collected On : 17/04/2026 4:15PM
 Reported On : 17/04/2026 11:11P



Investigation

Observed Value

Unit

Biological Reference Interval

PT (Prothrombin Time)

(Citratd Plasma)

Prothrombin Time**38.1****C** sec

9.4-12.5

Kindly note change in reference ranges

Control (MNPT)**11.2**

sec

= 11.0

Ratio

3.40

--

Index

29.40

%

--

PT(INR) Value**3.44**

Normal Population : 0.8 - 1.2

Standard Therapy: 2.0-3.0

High Dose Therapy: 3.0-4.5

Test done on Fully Automated Coagulometer (Clotting)

Note:

Kindly correlate with clinical and therapeutic history.

Interpretation:

- The Prothrombin time (PT) measures the extrinsic (Factor VII) & common coagulation pathways (Factors II, V, X & Fibrinogen).
- This test used is used for monitoring oral anticoagulation therapy, which lowers the levels of multiple vitamin K dependent coagulation factors in blood (Factor II, VII, IX, X).
- The results of PT is expressed as International Normalized Ratio (INR) to neutralize the influence of variable sensitivity of the reagents (thromboplastin) used in the assay by different laboratories.
- INR is the most commonly used parameters for monitoring oral anticoagulant treatment. Its therapeutic range varies with disease and treatment dosage.
- The INR is used only for patients on stable oral anticoagulant therapy. It makes no significant contribution to the diagnosis or treatment of patients whose PT is prolonged for other reasons.

Increased PT time may be due to:

1. Congenital or acquired factor deficiencies of (Factor II, V, VII, X, Fibrinogen), Coumarin (Warfarin – Vitamin K antagonists), therapy.

- Liver Diseases (Bile duct obstruction, Cirrhosis, Hepatitis).
- Hemorrhagic Disease of the newborn.
- DIC.
- Malabsorption.
- Fibrinolysis.
- Vitamin K deficiency

2. Combined Abnormal PT & APTT is found under two circumstances:



Dr. SANJAY GOHIL
 MD (Pathology)
 HOD Hematology & Transplant Immunology
 Reg No.2009093391

MEDICAL LABORATORY REPORT

** Referred Test "C" represents critical results



Name **Mr. MAHESH KAVITISHETTY**
Age / Gender : 29.2 Year(s)/ Male
Contact No. : +919967833785
Address JJ H Mumbai..
Pin code : 400008

VID No. : 250138509972839
PID No. : P14525547224709
Referred by : DR.JJ HOSPITAL
Registered On :17/04/2026 4:03PM
Collected On :17/04/2026 4:15PM
Reported On :17/04/2026 11:PM

Investigation**Observed Value****Unit****Biological Reference Interval**

- Medical: Administration of oral anticoagulants, DIC, Liver disease, Vitamin K deficiency, Massive transfusions.
- coagulation factor abnormalities: dysfibrinogenemia's; Factor V, X, and II defects.

Interference in PT/INR

Alcohol, Antibiotics, Aspirin, Cimetidine, Thrombin Inhibitors (Increase PT), Barbiturates, Oral contraceptives, Hormone-replacement therapy (HRT), and Vitamin K (Decrease PT).

-- End of Report --



Test Marked with NABL symbol are in the scope of accreditation



Dr. SANJAY GOHIL
MD (Pathology)
HOD Hematology & Transplant Immunology
Reg No.2009093391

MEDICAL LABORATORY REPORT

** Referred Test