


# Distribution




# Bernoulli Distribution



# Bernouli


- There are two possibilities (pass or fail) with probability  $p$  of success and  $q = 1-p$  of failure..



Expectation :  $p$   
Variance :  $pq$



# Bernouli



$$\text{Expectation}, E(x) = \sum x_i P(x_i)$$

$$= 1 * p + 0 * q$$

$$= p$$

$$\text{Variance}, Var = \sum (x_i - \mu)^2 P(x_i)$$

$$= (1 - p)^2 * p + (0 - p)^2 * (1 - p)$$

$$= p(1 - p)$$


$$= pq$$



# Geometric Distribution


Number of independent and identical Bernoulli trials needed to get ONE success.

Eg. Number of attempts before I pass the exam



# Geometric Distribution

- You run a series of independent trial.
- There can be either a success or failure for each trial, and the probability of success is the same for each trial.
- How many trials are needed in order to get the first successful outcome.



# Geometric Distribution

- $PMF^* = P(X = r) = q^{r-1}p$


- $P(X > r) = q^r$

- $CDF^{**}, P(X \leq r) = 1 - q^r$

- $E(X) = \frac{1}{p} \quad var(x) = \frac{q}{p^2}$

PMF : Probability Mass Function

CDF : Cumulative Distribution Function





# BINOMIAL DISTRIBUTION






# Binomial Experiment

- The process consists of a sequence of  $n$  trials.
- Only two exclusive outcomes are possible in each trial. One outcome is called “Success” and other a “failure”.
- The probability of a success denotes  $p$ , does not change from trial to trial. The probability of failure is  $1-p$  and is also fixed from trial to trial.
- The trials are independent; the outcome of previous trial not influence future trial.




# Binomial Variable

- Let's consider a coin  $P(H) = 0.6$
- $P(T) = 0.4$
- $X = \#$  of heads after 10 flip of my coin
- Made up of independent trails
- Each trail can be classified as either success or failure
- Probability of success on each trail is constant




# Binomial distribution


- $X$  = # of heads after flipping coin 4 times
- Possible outcomes =  $2 * 2 * 2 * 2 = 16$
- $P(X = 0) = \frac{C_0^4}{32} = \frac{1}{32}$
- $P(X = 1) = \frac{C_1^4}{32} = \frac{5}{32}$
- $P(X = 2) = \frac{C_2^4}{32} = \frac{6}{32}$




# Binomial distribution



# Binomial distribution



# Binomial distribution





# Binomial Probability example

P(Score) = 70 % or 0.7

P(miss) = 30 % or 0.3

P(Exactly 2 scores in 6 attempts)

$$= C_2^6 (0.7)^2 (0.3)^4$$

P(Exactly k scores in n attempts)

$$= C_k^n (p)^k (1 - p)^{n-k}$$



# Question

In this area it is expected that 9 cars are passing in a hours. What is the probability that there are exactly 6 cars are passing in this hour

$$E(X) = \lambda = n * p = 9 \text{ cars/hr}$$


Since , 1 hours = 60 minutes

So,  $n = 60$

$K = 6$  (number of expected cars in hour)

$$P(x = k) = C_k^n (p)^k (1 - p)^{n-k}$$

$$p = \lambda/n = 9 / 60$$





# Question

In this area it is expected that 9 cars are passing in a hours. What is the probability that there are exactly 6 cars are passing in this hour

$$p = \lambda / n = 9 / 60$$

$$1 - p = 51 / 60$$

Therefore, probability of exactly 6 cars are passing in an hours is

$$P(x = 6) = C_6^{60} \left(\frac{9}{60}\right)^6 \left(\frac{51}{60}\right)^{54}$$

$$n = 3600 \text{ (no. of seconds in an hour)}$$



A decorative graphic in the background consisting of a central white circle containing a grey stylized flower or star shape with eight petals. This is surrounded by a red circular arc, and at the top and bottom of the arc are two solid red circles.

# Poisson Distribution



# Poisson


$$E(X) = \tilde{\lambda} = n * p$$

$$P(X = k) = \lim_{n \rightarrow \infty} C_k^n (p)^k (1 - p)^{n-k}$$

$$= \frac{\tilde{\lambda}^k}{k!} e^{-\tilde{\lambda}}$$

$\tilde{\lambda} = 9$  cars pass

$P(X = 2)$



# Question

Q. Probability that a cars will not pass in 6 min

$$= \frac{\lambda^k}{k!} e^{-\lambda}$$

For 1<sup>st</sup> second:

$$P(X = 0) = \frac{\lambda^0}{0!} e^{-\lambda} = e^{-\lambda} \quad (\text{for 1 sec})$$

For  $n$  second:

Probability that a cars will pass for  $n$  sec.

$$= e^{-n\lambda}$$



# Exponential Distribution

Q. Probability that a cars will pass in  $n$  sec

$$1 - e^{-n\lambda}$$

$$\text{CDF} = 1 - e^{-n\lambda}, n \geq 0$$

$$\text{PDF} = \lambda e^{-n\lambda}, n \geq 0$$



# Exponential Distribution


- Poisson process
- Continuous analog of Geometric distribution

$$E(X) = \frac{1}{\lambda} \quad \text{var}(X) = \frac{1}{\lambda^2}$$



# Reference

- <https://www.mathsisfun.com/data/index.html>
- **Head First: Statistics**
- **KhanAcademy**





# INNOMATICS RESEARCH LAB

