



Pathogenesis of rheumatoid arthritis and c-Fos/AP-1

Shunichi Shiozawa & Ken Tsumiyama

To cite this article: Shunichi Shiozawa & Ken Tsumiyama (2009) Pathogenesis of rheumatoid arthritis and c-Fos/AP-1, Cell Cycle, 8:10, 1539-1543, DOI: [10.4161/cc.8.10.8411](https://doi.org/10.4161/cc.8.10.8411)

To link to this article: <https://doi.org/10.4161/cc.8.10.8411>



Published online: 15 May 2009.



Submit your article to this journal [↗](#)



Article views: 282



Citing articles: 33 View citing articles [↗](#)

Review

Pathogenesis of rheumatoid arthritis and c-Fos/AP-1

Shunichi Shiozawa^{1-3,*} and Ken Tsumiyama²

¹Division of Rheumatology; Department of Medicine; Kobe University Graduate School of Medicine; and The Center for Rheumatic Diseases; Kobe University Hospital; Chuoku, Kobe Japan; ²Division of Bioregulation; Department of Biophysics; Kobe University Graduate School of Health Science; Sumaku, Kobe Japan; ³Investigator of the Center of Excellence (COE); Japan

Key words: rheumatoid arthritis, c-Fos/AP-1, inflammatory cytokine, matrix metalloproteinase, joint destruction

c-Fos/AP-1 controls the expression of inflammatory cytokines and matrix-degrading matrix metalloproteinases (MMPs) important in arthritis via promoter AP-1 binding motif. Among inflammatory cytokines, IL-1 β is the most important inducer of a variety of MMPs, and mainly responsible for cartilage breakdown and osteoclastogenesis. IL-1 β and c-Fos/AP-1 influence each other's gene expression and activity, resulting in an orchestrated cross-talk that is crucial to arthritic joint destruction, where TNF α can act synergistically with them. While how to stop the degradation of bone and cartilage, i.e., to control MMP, has long been the central issue in the research of rheumatoid arthritis (RA), selective inhibition of c-Fos/AP-1 does resolve arthritic joint destruction. Thus, the blockade of IL-1 β and/or c-Fos/AP-1 can be promising as an effective therapy for rheumatoid joint destruction in addition to the currently available TNF α blocking agents that act mainly on arthritis.

Introduction

Immediate early gene *c-fos* contributes significantly to arthritis. This somewhat unexpected role of c-Fos/AP-1 is by no means exceptional when considering the roles of c-Fos/AP-1 in the transactivation of the genes responsible for arthritis such as inflammatory cytokines and matrix metalloproteinases (MMPs).¹⁻⁴ Most of the genes encoding inflammatory cytokines and matrix-degrading MMPs are under the control of c-Fos/AP-1, and c-Fos/AP-1 transactivates such genes by binding directly their promoter AP-1 motifs.¹⁻¹⁴ How to stop the degradation of bone and cartilage, i.e., to control MMP, has long been the central issue in the research of rheumatoid arthritis (RA), since cartilaginous matrix can only be degraded after the first attack of collagen fibrils by MMP.^{15,16} Here I review the pathogenesis of rheumatoid joint destruction in relation transcription factor including c-Fos/AP-1.

Cell Cycle and Arthritis

Cell cycle is organized by a family of cyclin-dependent protein kinases (cdks). There are several lines of evidence suggesting that cell cycle progression is distorted in RA. First, the manner of the growth of rheumatoid synovial cell is 'tumor-like': cells with 'transforming' appearance containing abundant cytoplasm, large pale nuclei and prominent nucleoli with karyotypic alteration are typically found adjacent to the affected joints in human RA and experimental animal model of arthritis.¹⁷⁻¹⁹ Second, the transcription factors acting at the G₁ phase of cell cycle including c-Fos/AP-1,^{20,21} Erg-1,²² CREB²³ and C/EBP²⁴ are increased, whereas the proportion of PCNA⁺ or Ki-67⁺ cells is decreased^{25,26} in rheumatoid synovial. Third, Wee1 kinase which inhibits mitotic cell division by phosphorylating cdc2 is under the control of c-Fos/AP-1, and Wee1 is increased simultaneously with c-Fos/AP-1 in RA.^{10,11} The findings suggest that mitotic cell division is arrested in face of heightened cell proliferation in RA due to upregulated c-Fos/AP-1 and Wee1, thereby causing a 'tumor-like' synovial overgrowth, an important, classical attribute of rheumatoid synovia.²⁷

Factors that Initiate Arthritis

Arthritic joint destruction is, needless to say, heralded by arthritis. The initial phase of arthritis is characterized by (1) 'tumor-like' synovial cellular overgrowth not necessarily accompanied by lymphocytes,¹⁷⁻¹⁹ (2) peri-articular osteoporosis seen as early as a few month after the onset of RA,²⁸ and (3) loss of cartilage typically seen within 1 year after disease onset. Matrix-degrading MMPs and inflammatory cytokine IL-1 β released from synovial cell and/or osteoclast, appear to be responsible for the initiation of arthritis (Fig. 1A). IL-1 β is the most important inducer of a variety of MMPs including MMP1, 3, 8, 13 and 14,²⁹ and IL-1 β can activate osteoclasts³⁰ and degrade cartilaginous matrix most strongly.^{29,31,32} and vice versa, MMPs cleave IL-1 β into its active forms.³³ Time course studies in animal experiment show that IL-1 β , IL-6 and MMP-3 are initially increased in arthritis,^{34,35} and IL-6, and possibly TNF α , appears activated subsequent to IL-1 β in collagen-induced arthritis (CIA) and human synovial cells (Fig. 1B).^{31,36-38} IL-1 β is increased in sera, synovia and joints of patients with RA,³⁹⁻⁴¹ and serum levels of IL-1 β correlate with its disease severity.^{42,43}

*Correspondence to: Shunichi Shiozawa; The Center for Rheumatic Diseases; Kobe University Hospital; 7-5 Kusunokicho; Chuoku, Kobe 650-0017 Japan; Email: shioz@med.kobe-u.ac.jp

Submitted: 03/10/09; Accepted: 03/13/09

Previously published online as a *Cell Cycle* Epublication:
<http://www.landesbioscience.com/journals/cc/article/8411>




Figure 1. Relationship between inflammatory cytokines, c-Fos/AP-1 and matrix degrading MMPs in arthritis and arthritic joint destruction. (A) On initiation of arthritis, c-Fos/AP-1 is upregulated in the cell and IL-1 β and matrix-degrading MMPs are released. IL-1 β and the increase of c-Fos/AP-1 in the cell stimulate osteoclast to degrade joints. AP, antigen presentation. (B) In full-blown arthritis, IL-1 β stimulates Th to release IL-6 and TNF α . IL-1 β , IL-6 and TNF α then stimulate Th17 cell to release IL-17. Inflammatory cytokines IL-6, TNF α and IL-17 play essential roles in arthritis. Besides, IL-6 also acts on hepatocyte to release C-reactive protein (CRP). (C) In the advanced phase of arthritis, IL-6 and TNF α stimulate synovial cell to release IL-1 β and then mutual positive interaction between IL-1 β , IL-6 and TNF α is established. (D) At later stage, IL-1 β , IL-6 and TNF α stimulate osteoclast and synovial cells to release MMPs to enhance farther destruction of joints.

Factors that Aggravate Arthritis

TNF α , IL-6 and IL-17 are the main players in full-blown arthritis (Fig. 1B). TNF-blocking agents, infliximab, etanercept and adalimumab, not only inhibit arthritis but also halt radiologic joint destruction.⁴⁴⁻⁴⁹ TNF α is produced from lymphoid cells quickly in response to infection or inflammation via NF κ B-dependent and NF κ B non-dependent pathways, the latter including ATF2, c-Jun, Erg 1 and Sp1.^{50,51} TNF α upregulates NF κ B directly and c-Fos/AP-1 indirectly via MAPK/JNK pathway.^{1,52} TNF α phosphorylates ATF2, thereby upregulating IL-6,⁵² IL-17,^{53,54} and TNF α itself⁵⁰ (Fig. 1B). TNF α increases MMP1, 3 and 9 and inhibits bone formation via Dickkopf-1 (Dkk-1).⁵⁵⁻⁵⁸ However, TNF α cannot stimulate osteoclasts directly, just helping M-CSF- or RANKL-induced osteoclast formation.²⁹ The direct effect of TNF α on MMPs is far weaker than that of IL-1 β .^{29,31,32}

IL-1 β and TNF α promote Th17 cell commitment and IL-17 production, a process principally driven by IL-6 and TGF β

(Fig. 1B).⁵⁹ Here, the presence of IL-6 is of particular importance, because stimulation of CD4 T cell with IL-6 plus TGF β potently induces Th17 differentiation whereas stimulation with TGF β triggers the development of regulatory T cells.⁶⁰⁻⁶² IL-17 is increased in the synovial tissue and synovial fluid of patients with RA.⁶³⁻⁶⁷ IL-17 induces IL-1 β , TNF α , IL-6, IL-8 and macrophage inflammatory protein (MIP)-1 α , MMPs and receptor activator of NF κ B ligand (RANKL) (Fig. 1C).^{65,68-73} IL-17 acts in synergy with IL-1 β and TNF α to induce cytokine or chemokine⁷⁴ and cartilage destruction^{73,75} in vitro. While inhibition of IL-17 substantially improves arthritis,⁷⁶⁻⁷⁹ the effect of IL-17 blockade appears clinically modest as compared with a dramatic effect of IL-1 β blockade.^{51,80}

IL-6 receptor (IL-6R) blockade also inhibits arthritis significantly. IL-6 is a major inducer of C-reactive protein from liver and thus a key mediator of inflammation.²⁹ During inflammation, IL-1 β acts on bone marrow to release neutrophils⁸¹ and on endothelial cells to release IL-6, and IL-6 thus-released increases

C-reactive protein and thrombocytosis.^{81,82} IL-6 vice versa, induces IL-1 β , TNF α and c-Fos/AP-1.^{50,84} Although C-reactive protein is an independent risk factor for bone loss and fracture and IL-6 can differentiate Th17 cells in cooperation with TGF β thereby increasing RANKL on osteoclasts (Fig. 1D),^{29,85} IL-6R blockade is less protective on radiologic joint destruction.⁸⁶⁻⁸⁸

Factors for Arthritic Joint Destruction

A recombinant form of the drug of naturally occurring IL-1 receptor antagonist (IL-1ra) is with lower clinical efficacy for RA.⁸⁹⁻⁹¹ This could be due to the fact that a 50% inhibition of IL-1 activity requires approximately 100-fold excess of IL-1ra, however, IL-1ra is on the other hand highly effective in the treatment of IL-1 β -centered autoinflammatory syndrome.⁹² This would mean that cascades of inflammatory cytokine are responsible for the pathogenesis of RA, and thus IL-1 β blockade alone appears less effective. Nevertheless, IL-1 β is the most important inducer of a variety of MMPs,^{29,93} even compared in molar basis.^{30,94} It induces the release of latent form of MMPs such as MMP3 or MMP13 crucial for collagen breakdown⁹⁵ and of active aggrecanase which is the dominant enzyme responsible for proteoglycan loss.^{30,96,97} There are additive effects on the induction of MMPs between IL-1 β and other cytokines or growth factors including TNF α , oncostatin M, FGF and PDGF.²⁹ Cartilage damage is fully dependent, even in TNF-triggered arthritis, on IL-1.²⁹ IL-1 β is 100-fold potent than TNF α in inducing cartilage destruction in vivo, where IL-1 β can act synergistically with TNF α ,^{97,98} suggesting that IL-1 is a key link between synovitis and cartilage breakdown (Fig. 1D). IL-1 β also facilitates osteoclast formation and significantly enhance RANKL-induced osteoclastogenesis.^{29,90,91} A total lack of chronic erosive arthritis is noted in IL-1 β -deficient mice,⁹⁶ and arthritic joint destruction in TNF α transgenic mice is totally inhibited by anti-IL-1R antibody in face of still increased serum TNF α .⁹⁹ Arthritis in IL-1ra-deficient mice are highly progressive.³⁸ Thus, IL-1 is crucially important in the progression of joint destruction in RA and all animal models.^{37,95}

c-Fos/AP-1 and Joint Destruction

The invasive front of rheumatoid pannus, a granulation tissue degrading cartilage and bone, is composed of synovial mesenchymal cells, but not lymphocytes.¹⁷ The synovial mesenchymal cells, which can present antigen to T cells¹⁰⁰⁻¹⁰² and release IL-1 β and other inflammatory mediators, are driven by upregulated c-Fos/AP-1 in RA.^{6,20,21} Upregulated c-Fos/AP-1 in RA causes synovial overgrowth^{103,104} and osteoporosis^{8,28} typically seen in arthritis. We found that selective inhibition of c-Fos/AP-1 resolved arthritis by the use of a specific small-molecule inhibitor of c-Fos/AP-1 designed and synthesized using 3D pharmacophore modeling based on the crystal structure of the AP-1-DNA complex.³⁵ c-Fos/AP-1 controls the expression of the inflammatory cytokines and matrix-degrading MMPs important in arthritis via promoter AP-1 binding motif.^{1-5,12-14} This inhibitor³⁵ inhibits most of the matrix-degrading MMPs, including MMP3, 9 and 13, as well as inflammatory cytokines including IL-1 β . While specific inhibition of individual MMPs, such as MMP3, cannot ameliorate

CIA possibly owing to compensation by another MMP,¹⁰⁵ broad-spectrum MMP inhibition successfully reduces the clinical severity and cartilage destruction of arthritis.³⁵ Interestingly, c-Fos/AP-1 and IL-1 β influence each other's gene expression and activity, resulting in an orchestrated cross-talk that is crucial to arthritic joint destruction,^{2,12-14,30,106,107} both c-Fos/AP-1 and IL-1 β are indispensable for arthritic joint destruction.^{30,36-38,99,108-110}

Concluding Remarks

A controversy exists regarding the importance of mesenchymal synovial cell and/or lymphocyte in the pathogenesis of RA. It is clear that synovial mesenchymal cells are responsible for joint destruction, however, T cell-centered immunity is also important with regards to triggering antigen. A representative mediators released from mesenchymal synovial cells would be IL-1 β , c-Fos/AP-1, MMP and/or TNF α and those of lymphocytes would be TNF α and/or IL-6. There are substantial cross-talk between them, and the change in one mediator would surely influence the others. However, as discussed, TNF α , IL-6 and IL-17 are important for arthritis, and IL-1 β and c-Fos/AP-1 for joint destruction. Thus, c-Fos/AP-1 inhibitor appears to be a promising drug for rheumatoid joint destruction. However, the fact that c-Fos/AP-1 acts mainly on mesenchymal cells and also upregulates Wee1 kinase opens a new application of c-Fos/AP-1 inhibitor for sleep disturbance. We recently showed that there is a mutual interaction between biological clock gene *Cry* and Wee1 kinase. Arthritis is extremely aggravated in the absence of circadian rhythmicity and vice versa, arthritis disturbs the circadian rhythm of the host.¹¹¹

References

- Angel P, Karin M. The role of Jun, Fos and the AP-1 complex in cell-proliferation and transformation. *Biochim Biophys Acta* 1991; 1072:129-57.
- Karin M, Liu Z-G, Zandi E. AP-1 function and regulation. *Curr Opin Cell Biol* 1997; 9:240-6.
- Angel P, Imagawa M, Chiu R, Stein B, Imbra RJ, Rahmsdorf HJ, et al. Phorbol ester-inducible genes contain a common cis element recognized by a TPA-modulated trans-acting factor. *Cell* 1987; 49:729-39.
- Schonthal A, Herrlich P, Rahmsdorf HJ, Ponta H. Requirement for *fos* gene expression in the transcriptional activation of collagenase by other oncogenes and phorbol esters. *Cell* 1988; 54:325-34.
- Gutman A, Waslyk B. The collagenase gene promoter contains a TPA and oncogene-responsive unit encompassing the PEA3 and AP-1 binding sites. *EMBO J* 1990; 9:2241-6.
- Shiozawa S, Tanaka Y, Fujita T, Tokuhisa T. Destructive arthritis without lymphocyte infiltration in H2-*c-fos* transgenic mice. *J Immunol* 1992; 148:3100-4.
- Kuroki Y, Shiozawa S, Sugimoto T, Fujita T. Constitutive expression of *c-fos* gene inhibits type 1 collagen synthesis in transfected osteoblasts. *Biochem Biophys Res Commun* 1992; 182:1389-94.
- Miyauchi A, Kuroki Y, Fukase M, Fujita T, Chihara K, Shiozawa S. Persistent expression of proto-oncogene *c-fos* stimulates osteoclast differentiation. *Biochem Biophys Res Commun* 1994; 205:1547-55.
- Shiozawa S, Shimizu S, Tanaka K, Hino K. Studies on the contribution of *c-fos*/AP-1 to arthritic joint destruction. *J Clin Invest* 1997; 99:1210-6.
- Kawasaki H, Komai K, Ouyang Z, Murata M, Hikasa M, Ohgiri M, et al. c-Fos/activator protein-1 transactivates wee1 kinase at G₁/S to inhibit premature mitosis in antigen-specific Th1 cells. *EMBO J* 2001; 20:4618-27.
- Kawasaki H, Komai K, Nakamura M, Yamamoto E, Ouyang Z, Nakashima T, et al. Human wee1 kinase is directly transactivated by and increased in association with c-Fos/AP-1: rheumatoid synovial cells overexpressing these genes go into aberrant mitosis. *Oncogene* 2003; 22:6839-44.
- Sirum-Connolly K, Brinckerhoff C. Interleukin-1 or phorbol induction of the stromelysin promoter requires an element that cooperates with AP-1. *Nucleic Acids Res* 1991; 19:335-41.

13. Hess J, Porte D, Munz C, Angel P. AP-1 and Cbfa/Runt physically interact and regulate parathyroid hormone-dependent MMP13 expression in osteoblasts through a new osteoblast-specific element 2/AP-1 composite element. *J Biol Chem* 2001; 276:20029-38.
14. Sun Y, Wenger L, Brinckerhoff CE, Misra RR, Cheung HS. Basic calcium phosphate crystals induce matrix metalloproteinase-1 through the Ras/mitogen-activated protein kinase/c-Fos/AP-1 metalloproteinase 1 pathway. *J Biol Chem* 2002; 277:1544-52.
15. Milner JM, Patel A, Rowan AD. Emerging roles of serine proteinases in tissue turnover in arthritis. *Arthritis Rheum* 2008; 58:3644-56.
16. Burrage PS, Mix KS, Brinckerhoff CE. Matrix metalloproteinase: role in arthritis. *Front Biosci* 2006; 11:529-43.
17. Shiozawa S, Shiozawa K, Fujita T. Morphologic observations in the early phase of the cartilage-pannus junction. *Arthritis Rheum* 1983; 26:472-8.
18. Fassbender HG. Is pannus a residue of inflammation? *Arthritis Rheum* 1984; 27:956-7.
19. Shiozawa S, Tanaka Y, Fujita T, Tokuhisa T. Destructive arthritis without lymphocyte infiltration in H2-c-fos transgenic mice. *J Immunol* 1992; 148:3100-4.
20. Trabandt A, Aicher WK, Gay RE, Sukhatme VP, Fassbender HG, Gay S. Spontaneous expression of immediately-early response genes c-fos and cgr-1 in collagenase-producing rheumatoid synovial fibroblasts. *Rheumatol Int* 1992; 12:53-9.
21. Asahara H, Fujisawa K, Kobata T, Hasunuma T, Maeda T, Asanuma M, et al. Direct evidence of high DNA binding activity of transcription factor AP-1 in rheumatoid arthritis synovium. *Arthritis Rheum* 1997; 40:912-8.
22. Aicher WK, Heer AH, Trabandt A, Bridges SL Jr, Schroeder HW Jr, Stransky G, et al. Overexpression of zinc-finger transcription factor Z-225/Egr-1 in synoviocytes from rheumatoid arthritis patients. *J Immunol* 1994; 152:5940-8.
23. Wakisaka S, Suzuki N, Takeno M, Takeba Y, Nagafuchi H, Saito N, et al. Involvement of simultaneous multiple transcription factor expression, including cAMP responsive element binding protein and OCT-1, for synovial cell outgrowth in patients with rheumatoid arthritis. *Ann Rheum Dis* 1998; 57:487-97.
24. Pope RM, Lovis R, Mungre S, Perlman H, Koch AE, Haines III GK. C/EBP beta in rheumatoid arthritis: correlation with inflammation, not disease specificity. *Clin Immunol* 1999; 91:271-82.
25. Jacobs RA, Perrett D, Axon JM, Herbert KE, Scott DL. Rheumatoid synovial cell proliferation, transformation and fibronectin secretion in culture. *Clin Exp Rheumatol* 1995; 13:717-23.
26. Schaser K, Kinne RW, Beil H, Kladny B, Stoss H. Proliferation of T-cells, macrophages, neutrophilic granulocytes and fibroblast-like cells in the synovial membrane of patients with rheumatoid arthritis. *Verh Dtsch Ges Pathol* 1996; 80:276-80.
27. Fassbender HG. Joint destruction in various arthritic diseases. In: Kuettner K, et al. eds. *Articular cartilage biochemistry*. New York: Raven Press 1986; 371-89.
28. Shimizu S, Shiozawa S, Shiozawa K, Imura S, Fujita T. Quantitative histologic studies on the pathogenesis of periarticular osteoporosis in rheumatoid arthritis. *Arthritis Rheum* 1985; 28:25-31.
29. Schett G, Stach C, Zwerina J, Voll R, Manger B. How antirheumatic drugs protect joints from damage in rheumatoid arthritis. *Arthritis Rheum* 2008; 58:2936-48.
30. Dayer J-M. The pivotal role of interleukin-1 in the clinical manifestations of rheumatoid arthritis. *Rheumatology* 2003; 42:3-10.
31. Van de Loo AAJ, van den Berg WB. Effects of murine recombinant IL-1 on synovial joints in mice: measurement of patellar cartilage metabolism and joint inflammation. *Ann Rheum Dis* 1990; 49:238-45.
32. Henderson B, Pettipher ER. Arthritogenic actions of recombinant IL-1 and TNF in the rabbit: evidence for synergistic interactions between cytokines in vivo. *Clin Exp Immunol* 1989; 75:306-10.
33. Kawasaki Y, Xu ZZ, Wang X, Park JY, Zhuang ZY, Tan PH, et al. Distinct roles of matrix metalloproteinases in the early- and late-phase development of neuropathic pain. *Nat Med* 2008; 14:331-6.
34. Thornton S, Duwel LE, Boivin GP, Ma Y, Hirsch R. Association of the course of collagen-induced arthritis with distinct patterns of cytokine and chemokine messenger RNA expression. *Arthritis Rheum* 1999; 42:1109-18.
35. Aikawa Y, Morimoto K, Yamamoto T, Chaki H, Hashimoto A, Narita H, et al. Treatment of arthritis with a selective inhibitor of c-Fos/activator protein-1. *Nat Biotechnol* 2008; 26:817-23.
36. Van den Berg WB. Lessons from animal models of arthritis. *Curr Rheumatol Rep* 2002; 4:232-9.
37. Joosten LAB, Helsen MMA, Saxne T, van De Loo FAJ, Heinegard D, van Den Berg D. IL-1 α blockade prevents cartilage and bone destruction in murine type II collagen-induced arthritis, whereas TNF α blockade only ameliorates joint destruction. *J Immunol* 1999; 163:5049-55.
38. Horai R, Saijo S, Tanioka H, Nakae S, Sudo K, Okahara A, et al. Development of chronic inflammatory arthropathy resembling rheumatoid arthritis in IL-1 receptor antagonist-deficient mice. *J Exp Med* 2000; 191:313-20.
39. Chikanza IC, Roux-Lombard P, Dayer JM, Panayi GS. Dysregulation of the in vivo production of interleukin-1 receptor antagonist in patients with rheumatoid arthritis. Pathogenetic implications. *Arthritis Rheum* 1995; 38:64-8.
40. Kahle P, Saal JG, Schaudt K, Zacher J, Fritz P, Pawelec G. Determination of cytokines in synovial fluids: correlation with diagnosis and histomorphological characteristics of synovial fluid. *Ann Rheum Dis* 1992; 51:731-4.
41. Koch AE, Kunkel SL, Chensue SW, Haines GK, Streiter RM. Expression of interleukin-1 and interleukin-1 receptor antagonist by human rheumatoid synovial tissue macrophages. *Clin Immunol Immunopathol* 1992; 65:23-9.
42. Eastgate JA, Symons JA, Wood NC, Grinlinton FM, di Giovine FS, Duff GW. Correlation of plasma interleukin-1 levels with disease activity in rheumatoid arthritis. *Lancet* 1988; 2:706-8.
43. Rooney M, Symons JA, Duff GW. Interleukin-1 β in synovial fluid is related to local disease activity in rheumatoid arthritis. *Rheumatol Int* 1990; 10:217-9.
44. Smolen JS, Van Der Heijde DM, St. Clair EW, Emery P, Bathon JM, Keystone E, et al. Predictors of joint damage in patients with early rheumatoid arthritis treated with high-dose methotrexate with or without concomitant infliximab: results from ASPIRE trial. *Arthritis Rheum* 2006; 54:702-10.
45. Klareskog L, van der Heijde D, de Jager JP, Gough A, Kalden J, Malaise M, et al. Therapeutic effect of the combination of etanercept and methotrexate compared with each treatment alone in patients with rheumatoid arthritis: double-blind randomized controlled trial. *Lancet* 2004; 363:675-81.
46. Keystone EC, Kavanaugh AF, Sharp JT, Tannenbaum H, Hua Y, Teoh LS, et al. Radiographic, clinical and functional outcomes of treatment with adalimumab (a human anti-tumor necrosis factor monoclonal antibody) in patients with active rheumatoid arthritis receiving concomitant methotrexate therapy: a randomized, placebo-controlled, 52-week trial. *Arthritis Rheum* 2004; 50:1400-11.
47. Lipsky PE, van der Heijde DM, St. Clair EW, Furst DE, Breedveld FC, Kalden JR, et al. Infliximab and methotrexate in the treatment of rheumatoid arthritis. *N Engl J Med* 2000; 343:1594-602.
48. Weinblatt ME, Keystone EC, Furst DE, Moreland LW, Weisman MH, Birbara CA, et al. Adalimumab, a fully human anti-tumor necrosis factor α monoclonal antibody, for the treatment of rheumatoid arthritis in patients taking concomitant methotrexate: the ARMADA trial. *Arthritis Rheum* 2003; 48:35-45.
49. Maini R, St. Clair EW, Breedveld F, Furst D, Kalden J, Weisman M, et al. Infliximab (chimeric anti-tumor necrosis factor α monoclonal antibody) versus placebo in rheumatoid arthritis patients receiving concomitant methotrexate: a randomized phase III trial. *Lancet* 1999; 354:1932-9.
50. Tsai EY, Falvo JV, Tsytsykova AV, Barczak AK, Reimold AM, Glimcher LH, et al. A lipopolysaccharide-specific enhancer complex involving Ets, Elk-1, Sp1 and CREB binding protein and p300 is recruited to the tumor necrosis factor alpha promoter in vivo. *Mol Cell Biol* 2000; 20:6084-94.
51. Finnegan A, Doodles PD. Pathways for interleukin-1-driven arthritis. *Arthritis Rheum* 2008; 58:3282-5.
52. Ventura J-J, Kennedy NJ, Lamb JA, Flavell RA, Davis RJ. c-Jun NH2-terminal kinase is essential for the regulation of AP-1 by tumor necrosis factor. *Mol Cell Biol* 2003; 23:2871-82.
53. Koenders MI, Devesa I, Marijnissen RJ, Abdollahi-Roodsaz S, Boots AM, Walgreen B, et al. Interleukin-1 drives pathogenic Th17 cells during spontaneous arthritis in interleukin-1 receptor antagonist-deficient mice. *Arthritis Rheum* 2008; 58:3461-70.
54. Stockinger B, Veldhoen M. Differentiation and function of Th17 T cells. *Curr Opin Immunol* 2007; 19:281-6.
55. Reunanan N, Li S-P, Ahonen M, Foschi M, Han J, Kahari V-M. Activation of p38 α MAPK enhances collagenase-1 (matrix metalloproteinase (MMP)-1) and stromelysin-1 (MMP-3) expression by mRNA stabilization. *J Biol Chem* 2002; 277:32360-8.
56. Yan C, Boyd DD. Regulation of matrix metalloproteinase gene expression. *J Cell Physiol* 2007; 211:19-26.
57. Ray A, Bal BS, Ray BK. Transcriptional induction of matrix metalloproteinase-9 in the chondrocyte and synovioyte cells is regulated via a novel mechanism: evidence for functional cooperation between serum amyloid A-activating factor-1 and AP-1. *J Immunol* 2005; 175:4039-48.
58. Diarra D, Stolina M, Polzer K, Zwerina J, Ominsky MS, Dwyer D, et al. Dickkopf-1 is a master regulator of joint remodelling. *Nat Med* 2007; 13:156-63.
59. Bettelli E, Carrier Y, Gao W, Korn T, Strom TB, Oukka M, et al. Reciprocal developmental pathways for the generation of pathogenic effector Th17 and regulatory T cells. *Nature* 2006; 441:235-8.
60. Mangan PR, Harrington LE, O'Quinn DB, et al. Transforming growth factor- β induces development of the Th17 lineage. *Nature* 2006; 441:231-4.
61. Veldhoen M, Hocking RJ, Atkins CJ, Locksley RM, Stockinger B. TGF β in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells. *Immunity* 2006; 24:179-89.
62. Chabaud M, Durand JM, Buchs N, Fossiez F, Page G, Frappart L, et al. Human interleukin-17: a T cell-derived proinflammatory cytokine produced by the rheumatoid synovium. *Arthritis Rheum* 1999; 42:963-70.
63. Ziolkowska M, Koc A, Luszczkiewicz G, Ksiezopolska-Pietrzak K, Klimczak E, Chwalinska-Sadowska H, et al. High levels of IL-17 in rheumatoid arthritis patients: IL-15 triggers in vitro IL-17 production via cyclosporine A-sensitive mechanism. *J Immunol* 2000; 164:2832-8.
64. Kotake S, Udagawa N, Takahashi N, Matsuzaki K, Itoh K, Ishiyama S, et al. IL-17 in synovial fluids from patients with rheumatoid arthritis is a potent stimulator of osteoclastogenesis. *J Clin Invest* 1999; 103:1345-52.

65. Hwang SY, Kim HY. Expression of IL-17 homologs and their receptors in the synovial cells of rheumatoid arthritis patients. *Mol Cell* 2005; 19:180-4.
66. Raza K, Falciani F, Curnow SJ, Ross EJ, Lee CY, Akbar AN, et al. Early rheumatoid arthritis is characterized by a distinct and transient synovial fluid cytokine profile of T cell and stromal cell origin. *Arthritis Res Ther* 2005; 7:784-95.
67. Jovanovic DV, Di Battista JA, Martel-Pelletier J, Jolicoeur FC, He Y, Zhang M, et al. IL-17 stimulates the production and expression of proinflammatory cytokines, IL-1 β and TNF α , by human macrophages. *J Immunol* 1998; 160:3513-21.
68. Fossiez F, Djossou O, Chomarat P, Flores-Romo L, Ait-Yahia S, Maat C, et al. T cell interleukin-17 induces stromal cells to produce proinflammatory and hematopoietic cytokines. *J Exp Med* 1996; 183:2593-603.
69. Yao Z, Fanslow WC, Seldin MF, Rousseau AM, Painter SL, Comeau MR, et al. Herpesvirus Saimiri encodes a new cytokine, IL-17, which binds to a novel cytokine receptor. *Immunity* 1995; 3:811-21.
70. Chabaud M, Garnerio P, Dayer JM, Guerne PA, Fossiez F, Miossec P. Contribution of interleukin 17 to synovium matrix destruction in rheumatoid arthritis. *Cytokine* 2000; 12:1092-9.
71. Benderdour M, Tardif G, Pelletier JR, Di Battista JA, Reboul P, Ranger P, et al. Interleukin 17 (IL-17) induces collagenase-3 production in human osteoarthritic chondrocytes via AP-1 dependent activation: differential activation of AP-1 members by IL-17 and IL-1 β . *J Rheumatol* 2002; 29:1262-72.
72. Koshy PJ, Henderson N, Logan C, Life PF, Cawston TE, Rowan AD. Interleukin 17 induces cartilage collagen breakdown: novel synergistic effects in combination with proinflammatory cytokines. *Ann Rheum Dis* 2002; 61:704-13.
73. Koenders MI, Joosten LAB, van der Berg WB. Potential new targets in arthritis therapy: interleukin (IL)-17 and its relation to tumour necrosis factor and IL-1 in experimental arthritis. *Ann Rheum Dis* 2006; 3:29-33.
74. Van Bezooijen RL, van der Wee-Pals L, Papapoulos SE, Lowik CW. Interleukin-17 synergises with tumour necrosis factor α to induce cartilage breakdown in vitro and in vivo. *Ann Rheum Dis* 2002; 61:870-6.
75. Nakae S, Saijo S, Horai R, Sudo K, Mori S, Iwakura Y. IL-17 production from activated T cells is required for the spontaneous development of destructive arthritis in mice deficient in IL-1 receptor antagonist. *Proc Natl Acad Sci USA* 2003; 100:5986-90.
76. Lubberts E, Koenders MI, Oppers-Walgreen B, van den Bersselaar L, Coenen-de Roo CJ, Joosten LA, et al. Treatment with a neutralizing anti-murine interleukin-17 antibody after the onset of collagen-induced arthritis reduces joint inflammation, cartilage destruction and bone erosion. *Arthritis Rheum* 2004; 50:650-9.
77. Fujimoto M, Serada S, Mihara M, Uchiyama Y, Yoshida H, Koike N, et al. Interleukin-6 blockade suppresses autoimmune arthritis in mice by the inhibition of inflammatory Th17 responses. *Arthritis Rheum* 2008; 58:3710-9.
78. Horai R, Nakajima A, Habiro K, Kotani M, Nakae S, Matsuki T, et al. TNF α is crucial for the development of autoimmune arthritis in IL-1 receptor antagonist-deficient mice. *J Clin Invest* 2004; 114:1603-11.
79. Finnegan A, Grusby MJ, Kaplan CD, O'Neill SK, Eibel H, Koreny T, et al. IL-4 and IL-12 regulate proteoglycan-induced arthritis through Stat-dependent mechanisms. *J Immunol* 2002; 169:3345-52.
80. Doodes PD, Cao Y, Hamel KM, Wang Y, Farkas B, Iwakura Y, et al. Development of proteoglycan-induced arthritis is independent of IL-17. *J Immunol* 2008; 181:329-37.
81. Dinarello CA. The many worlds of reducing interleukin-1. *Arthritis Rheum* 2005; 52:1960-7.
82. Hoffman HM, Rosengren S, Boyle DL, Cho JY, Nayar J, Mueller JL, et al. Prevention of cold-associated acute inflammation in familial cold autoinflammatory syndrome by interleukin-1 receptor antagonist. *Lancet* 2004; 364:1779-85.
83. Ogura H, Murakami M, Okuyama Y, Tsuruoka M, Kitabayashi C, Kanamoto M, et al. Interleukin-17 promotes autoimmunity by triggering a positive-feedback loop via interleukin-6 induction. *Immunity* 2008; 29:628-36.
84. Yang E, Lerner L, Besser D, Darnell E Jr. Independent and cooperative activation of chromosomal c-fos promoter by STAT3. *J Biol Chem* 2003; 278:15794-9.
85. Wong PK, Quinn JM, Sims NA, van Nieuwenhuijze A, Campbell IK, Wicks IP. Interleukin-6 modulates production of T lymphocyte-derived cytokines in antigen-induced arthritis and drives inflammation-induced osteoclastogenesis. *Arthritis Rheum* 2006; 54:158-68.
86. Maini RN, Taylor PC, Szechinski J, Pavelka K, Bröll J, Balint G, et al. Double-blind randomized controlled clinical trial of the interleukin-6 receptor antagonist, tocilizumab, in European patients with rheumatoid arthritis who had an incomplete response to methotrexate. *Arthritis Rheum* 2006; 54:2817-29.
87. Nishimoto N, Hashimoto J, Miyasaka N, Yamamoto K, Kawai S, Takeuchi T, et al. Study of active controlled monotherapy used for rheumatoid arthritis, an IL-6 inhibitor (SAMURAI): evidence of clinical and radiographic benefit from an x ray reader-blinded randomized controlled trial of tocilizumab. *Ann Rheum Dis* 2007; 66:1162-7.
88. Smolen JS, Beaulieu A, Rubbert-Roth A, Ramos-Remus C, Rovinsky J, Alecock E, et al. Effect of interleukin-6 receptor inhibition with tocilizumab in patients with rheumatoid arthritis (OPTION study): a double-blind, placebo-controlled, randomized trial. *Lancet* 2008; 371:987-97.
89. Cohen SB, Moreland LW, Cush JJ, Greenwald MW, Block S, Shergy WJ, et al. A multicentre, double blind, randomized, placebo controlled trial of anakinra (Kineret), a recombinant interleukin 1 receptor antagonist, in patients with rheumatoid arthritis treated with background methotrexate. *Ann Rheum Dis* 2004; 63:1062-8.
90. Jiang Y, Genant HK, Watt I, Cobby M, Bresnahan B, Aitchison R, et al. A multicenter, double-blind, dose-ranging, randomized, placebo-controlled study of recombinant human interleukin-1 receptor antagonist in patients with rheumatoid arthritis: radiologic progression and correlation of Genant and Larsen scores. *Arthritis Rheum* 2000; 43:1001-9.
91. Zwerina J, Redlich K, Polzer K, Joosten L, Krönke G, Distler J, et al. TNF-induced structural joint damage is mediated by interleukin-1. *Proc Natl Acad Sci USA* 2007; 104:11742-7.
92. Braddock M, Quinn A. Targeting IL-1 inflammatory disease: new opportunities for therapeutic intervention. *Nat Rev Drug Discovery* 2004; 3:1-10.
93. Pap T, Gay S, Schett G. Matrix metalloproteinases. In: Smolen JS, Lipsky PE, editors. *Biological therapies in rheumatology*. (London): Taylor & Francis Ltd 2002; 483-97.
94. Dayer JM, Beutler B, Cerami A. Cachectin/tumor necrosis factor stimulates collagenase and prostaglandin E2 production by human synovial cells and dermal fibroblasts. *J Exp Med* 1985; 162:2163-8.
95. Van den Berg WB. Anti-cytokine therapy in chronic destructive arthritis. *Arthritis Res* 2001; 3:18-26.
96. Gravallese EM, Goldring SR. Cellular mechanisms and the role of cytokines in bone erosions in rheumatoid arthritis. *Arthritis Rheum* 2000; 43:2143-51.
97. Van de Loo AAJ, van den Berg WB. Effects of murine recombinant IL-1 on synovial joints in mice: measurement of patellar cartilage metabolism and joint inflammation. *Ann Rheum Dis* 1990; 49:238-45.
98. Henderson B, Pettipher ER. Arthritogenic actions of recombinant IL-1 and TNF in the rabbit: evidence for synergistic interactions between cytokines in vivo. *Clin Exp Immunol* 1989; 75:306-10.
99. Probert T, Plows D, Kontogeorgos G, Kollias G. The type I IL-1 receptor acts in series with TNF α to induce arthritis in TNF α transgenic mice. *Eur J Immunol* 1995; 25:1794-7.
100. Shiozawa S, Shiozawa K, Fujita T. Presence of HLA-DR antigen on synovial type A and B cells: an immunoelectron microscopic study in rheumatoid arthritis, osteoarthritis and normal traumatic joints. *Immunology* 1983; 50:587-94.
101. Shimizu S, Shiozawa S, Shiozawa K, Imura S, Ishikawa H, Hirohata K, et al. The restoration of proliferation and differentiation of peripheral blood mononuclear non-adherent cells into immunoglobulin-secreting cells by autologous synovial adherent cells from patients with rheumatoid arthritis. *Virchow Arch B Cell Pathol* 1988; 54:350-6.
102. Klareskog L, Forsum U, Kabelitz D, et al. Immune functions of human synovial cells: phenotypic and T cell regulatory properties of macrophage-like cells that express HLA-DR. *Arthritis Rheum* 1982; 25:488-501.
103. Kuroki Y, Shiozawa S, Yoshihara R, Hotta T. The contribution of human c-fos DNA to cultured synovial cells: a transfection study. *J Rheumatol* 1993; 20:422-8.
104. Shimizu K, Kawasaki H, Morisawa T, et al. Spontaneous and cytokine regulated c-fos gene expression in rheumatoid synovial cells: resistance to cytokine stimulation when the c-fos gene is overexpressed. *Ann Rheum Dis* 2000; 59:636-40.
105. Mudgett JS, et al. Susceptibility of stromelysin 1-deficient mice to collagen-induced arthritis and cartilage destruction. *Arthritis Rheum* 1998; 41:110-21.
106. Whitmarsh AJ, Shore P, Sharrocks AD, Davis RJ. Integration of MAP kinase signal transduction pathways at the serum response element. *Science* 1995; 269:403-7.
107. Scott DL, Kingsley GH. Tumor necrosis factor inhibitors for rheumatoid arthritis. *N Engl J Med* 2006; 355:704-12.
108. Teitelbaum SL. RANKing c-Jun in osteoclast development. *J Clin Invest* 2004; 114:463-5.
109. Redlich K, Hayer S, Ricci R, David JP, Tohidast-Akrad M, Kollias G, et al. Osteoclasts are essential for TNF-mediated joint destruction. *J Clin Invest* 2002; 110:1419-27.
110. Kobayashi K, Takahashi N, Jimi E, Udagawa N, Takami M, Kotake S, et al. Tumor necrosis factor stimulates osteoclast differentiation by a mechanism independent of the ODF/RANKL-RANK interaction. *J Exp Med* 2000; 191:275-86.
111. Hashiramoto A, Yamane T, Tsumiyama K, et al. Biological clock modulates arthritis. Submitted.