

DAVE3700 - Matte 3000

Kristian Sørdal

June 22, 2022

Contents

1	Introduction	3
2	Domains, Graphs and Level Sets	4
2.1	Domain of definition	4
2.2	Graphs of functions	4
2.3	Level Sets	5
3	Derivatives	7
3.1	Partial Derivatives	7
3.2	Higher order derivatives	7
3.3	Chain Rule	7
3.4	The Gradient	8
4	Directional Derivatives and Critical Points	9
4.1	Directional Derivatives	9
4.2	Critical Points	10
5	Lecture 5	11
5.1	Hessian Matrix	11
5.2	Global extremal values	11
5.3	Constrained optimization	13
6	Lecture 6	14
6.1	Substitution Method	14
6.2	Lagrange's Method	14
6.3	Method for Langrange	15
7	Lecture 7	18
7.1	Parametrized Curve	18
7.2	Kinematics	22
8	Lecture 8	24
8.1	Determining Motion	24
8.2	Arc Length	25
8.3	Tangent Vectors	28
8.4	Normal Vectors	29
9	Lecture 9	31
9.1	Normal Vectors - Continued	31
9.2	Curvature	33
10	Lecture 10	37
10.1	Curves in polar form	37
10.2	Areas In Polar Form	39
11	Scalar and Vector Fields	41
11.1	Gradient, Divergence and Curl	43

12 Lecture 11	45
12.1 Operations on fields	45
12.2 Intepretation of Divergence	45
12.3 Interpretation of Curl	46
12.4 Scalar Field from Gradient	47
13 Lecture 12	49
13.1 Identities Between Operations	49
13.2 Line Integrals	49
13.3 Parametrization and Orientation	51
13.4 Case of Vector Fields	52
14 Line Integrals, Parametrization and Vector Fields	55
14.1 Identities between operations	55
14.2 Line Integrals	55
14.3 Parametrization and orientation	56
14.4 Case of Vector Fields	57

1 Introduction

2 Domains, Graphs and Level Sets

2.1 Domain of definition

A function may not be defined for all real numbers.

Example. $f(x) = \frac{1}{x}$ is not defined for $x = 0$ ◇

Definition 1. The domain of a function f is the set of numbers for which it is defined. We write the domain of f as D_f .

For instance, for $f(x) = \frac{1}{x}$ we have that

$$D_f = \{x \in \mathbb{R} : x \neq 0\}$$

This is the largest possible domain, we can also consider smaller domains. We have the interval from 1 to 2.

$$[1, 2] = \{x \in \mathbb{R} : 1 \leq x \leq 2\}$$

Example. Find the largest domain of $f(x, y) = \frac{1}{y-x}$. The denominator should be non-zero, we get

$$D_f = \{(x, y) \in \mathbb{R}^2 : y - x \neq 0\}$$
◇

Example. Same exercise with $f(x, y) = \sqrt{y - x^2}$.

Argument: $y - x^2 \geq 0$ (because square root). We will then have $y \geq x^2$ ◇

2.2 Graphs of functions

The plot of a function f describes its behaviour visually. Mathematically, a plot corresponds to the notion of a graph.

Definition 2. The graph of a function $f(x, y)$ with domain D_f is the set of points (x, y, z) such that:

$$(x, y) \in D_f \text{ and } z = f(x, y)$$

We write G_f for the set

$$G_f = \{(x, y, z) \in \mathbb{R}^3 : (x, y) \in D_f \text{ and } z = f(x, y)\}$$

The graph of a function of two variables will, in general, be a surface.

Example. Lets consider $f(x, y) = 1$, with domain \mathbb{R}^2 . The graph of f is:

$$G_f = \{(x, y, 1) : (x, y) \in \mathbb{R}^2\}$$

All points have $z = 1$, this is a plane. More generally, the graph of $f(x, y) = ax + by + c$ is a plane with linear dependence on x and y . \diamond

Example. Consider the graph of:

$$f(x, y) = x^2 + y^2, \quad D_F = \mathbb{R}^2$$

This surface is called a paraboloids. \diamond

Example. A sphere of radius r is defined by

$$x^2 + y^2 + z^2 = r^2$$

All points x, y, z satisfy the equation. \diamond

Is this the graph of a function? No!

There is no unique value of z , associated with (x, y) because:

$$z = \pm \sqrt{r^2 - x^2 - y^2}$$

Both satisfy the sphere equation. Lets consider the graph of

$$f(x, y) = \sqrt{r^2 - x^2 - y^2}$$

With the domain $x^2 + y^2 \leq r^2$. The grap is a half sphere.

2.3 Level Sets

Another way to visualize functions.

Definition 3. A level set of a function $f(x, y)$ is constant. Essentially, this is a topographic map.

Example. Consider the function

$$f(x, y) = x^2 + y^2$$

The level sets for $c > 0$ are circles.

$$f(x, y) = x^2 + y^2 = c = (\sqrt{c})^2$$

This is a circle with radius \sqrt{c} .

Now consider the case $c < 0$, then:

$$f(x, y) = x^2 + y^2 = c$$

which doesn't work, because the level sets are empty.

For $c = 0$, we only have the point $(x, y) = (0, 0)$. Generally, level sets of $f(x, y)$ is a curve.

◇

3 Derivatives

3.1 Partial Derivatives

In the case of one variable, we have

$$\frac{df}{dt} = \lim_{n \rightarrow 0} \frac{f(x+n) - f(x)}{n}$$

Similarly, for two or more variables, we have the following definition

Definition 4. The partial derivative of $f(x, y)$ with respect to x :

$$\frac{\partial f}{\partial x} = \lim_{n \rightarrow 0} \frac{f(x+n, y) - f(x, y)}{n}$$

Also written as f_x , for $\frac{\partial f}{\partial y}$, we have f_y

Note, the expression above is $\frac{\partial f}{\partial x}(x, y)$, which is the value at the point (x, y)

3.2 Higher order derivatives

Given $\frac{\partial f}{\partial x}$, we can take further derivatives. We have

$$\frac{\partial^2 f}{\partial^2 x} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right), \quad \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right)$$

Also written as $f_{xx}, f_{yy}, f_{xy}, f_{yx}$. In most cases, f_{xy} and f_{yx} coincide.

Theorem 5. Schwartz theorem: Suppose f_{xy} and f_{yx} exist, and are continuous, then

$$f_{xy} = f_{yx}$$

Similar definitions and results for the case of more variables: x_1, \dots, x_n , with n variables.

3.3 Chain Rule

Suppose $f(x) = g(h(x))$, for instance

$$f(x) = (\cos x)^2 \text{ with } g(x) = x^2, h(x) = \cos x$$

then the chain rule is

$$\frac{df}{dt}(x_0) = \frac{dg}{dh}(h(x_0)) \cdot \frac{dh}{dt}(x_0)$$

Generalization to more variables.

Theorem 6. Chain rule: consider $f(x, y)$ x and y depending on a variable t . Then:

$$\frac{df}{dt}t_0 = \frac{\partial f}{\partial x}(x(t_0), y(t_0)) \frac{dx}{dt}t_0 + \frac{\partial f}{\partial y}(x(t_0), y(t_0)) \frac{dy}{dt}t_0$$

The "short form" of this result is

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$$

Example. Consider $f(x, y) = xy$, where

$$x(t) = \cos t, y(t) = \sin t$$

This cannot be computed directly with $\frac{df}{dt}$.

$$f(t) = f(x(t), y(t)) = f(\cos t, \sin t) = \cos t \cdot \sin t$$

We can compute

$$\frac{df}{dt} = (\cos t)' \sin t + \cos t (\sin t)' = -(\sin t)^2 + (\cos t)^2$$

Using the chain rule, we get

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$$

◇

3.4 The Gradient

Define an operation that takes a scalar function, and returns a vector function.

Definition 7. The gradient of $f(x, y)$ at (x_0, y_0) is

$$\nabla f(x_0, y_0) = (f_x(x_0, y_0), f_y(x_0, y_0))$$

4 Directional Derivatives and Critical Points

4.1 Directional Derivatives

We have seen the following

- f_x = the rate of change along the x -direction.
- f_y = the rate of change along the y -direction.

What about general directions?

Definition 8. Let $\vec{u} = (a, b)$, the directional derivative along \vec{u} at (x, y) is

$$D_{\vec{u}}f(x, y) = \lim_{h \rightarrow 0} \frac{f(x + ha, y + hb) - f(x, y)}{h}$$

Note that

$$\vec{u} = (1, 0) \rightarrow D_{\vec{u}} = f_x$$

$$\vec{u} = (0, 1) \rightarrow D_{\vec{u}} = f_y$$

To compare directions, we take $|\vec{u}| = 1$. Here $|\vec{u}|$ is the length of \vec{u} , that is

$$|\vec{u}| = \sqrt{\vec{u} \cdot \vec{u}}$$

Proposition 1. We have the following result

$$D_{\vec{u}} \cdot f = \nabla f \cdot \vec{u}$$

Proof. Consider the following function

$$g(t) = f(x + ta, y + tb)$$

Its derivative at $t = 0$ is

$$\begin{aligned} \frac{dg}{dt}(0) &= \lim_{h \rightarrow 0} \frac{g(h) - g(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x + ha, y + hb) - f(x, y)}{h} \\ &= D_{\vec{u}} \cdot f(x, y) \end{aligned}$$

On the other hand, using the chain rule, we get

$$\begin{aligned}\frac{dg}{dt}(0) &= \frac{\partial f}{\partial x} \frac{d(x+ta)}{dt} + \frac{\partial f}{\partial y} \frac{d(y+tb)}{dt} \\ &= \frac{\partial f}{\partial x} a + \frac{\partial f}{\partial y} b = \nabla f \cdot \vec{u}\end{aligned}$$

□

We can now state another property of ∇f . The direction where f changes the most.

Proposition 2. $|D_{\vec{u}} \cdot f|$ is the largest when \vec{u} is parallel to ∇f .

Proof. Recall that, given two vectors, \vec{v} and \vec{w} , we have that

$$\vec{v} \cdot \vec{w} = |\vec{v}| \cdot |\vec{w}| \cdot \cos \alpha$$

When is $|\vec{v} \cdot \vec{w}|$ the largest? We have

$$|\vec{v} \cdot \vec{w}| = |\vec{v}| \cdot |\vec{w}| \cdot |\cos \alpha|, \quad |\cos \alpha| \leq 1$$

It is the largest when the following condition is true

$$|\cos \alpha| = 1, \quad \alpha = 0 \vee \pi$$

Which means that the vectors are pointing in the same, or opposite direction. Applying this to $D_{\vec{u}} \cdot f$, we get

$$\begin{aligned}|D_{\vec{u}} \cdot f| &= |\nabla f \cdot \vec{u}| \\ &= |\nabla f| \cdot |\vec{u}| \cdot |\cos \alpha|\end{aligned}$$

For fixed values of $|\vec{u}|$, this is the largest when $\alpha = 0 \vee \pi$. That is ∇f and \vec{u} are parallel. □

4.2 Critical Points

How do we find the maxima and minima of $f(x)$? Lets take a look at $f'(x_0) = 0$.

Definition 9. We say that x_0, y_0 is a critical point of f if:

$$\frac{\partial f}{\partial x}(x_0, y_0) = 0 \quad \wedge \quad \frac{\partial f}{\partial y}(x_0, y_0) = 0$$

Similarly for

5 Lecutre 5

5.1 Hessian Matrix

Example. Consider again the function $f(x, y) = x^2 - y^2$.

$$f_{xx} = 2, \quad f_{yy} = -2, \quad f_{yx} = 0$$

$$H = \begin{bmatrix} 2 & 0 \\ 0 & -2 \end{bmatrix} = \det H = -4 < 0$$

Hence (x_0, y_0) is a saddle point. ◇

Example. Consider the function $f(x, y) = x^2 + y^2$, we have

$$(f_x, f_y) = (2x, 2y)$$

The only critical point is $(x_0, y_0) = (0, 0)$.

$$H = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \det H = 4 > 0$$

Since $f_{xx} = 2 > 0$, we conclude that $(0, 0)$ is a local minima. In this case, it is actually a global minimum, because $f(x, y) = x^2 + y^2 \geq 0$. ◇

5.2 Global extremal values

A function can have many maxima and minimas. Usually, we are intrested in the largest and smallest values.

Definition 10. Let $f(x, y)$ be with domain D_f . Then we have

- (x_0, y_0) is a global maxima if $f(x_0, y_0) \geq f(x, y)$ for all $(x, y) \in D_f$.
- (x_0, y_0) is a global minima if $f(x_0, y_0) \leq f(x, y)$ for all $(x, y) \in D_f$.

Trivial example: for $f(x, y) = 1$, all points are global maxima and minima. Note that global maxima and minima need not be critical points.

Example. We have $f(x) = x$ with $D_f = [-1, 1]$.

- Global maxima at $x = 1, f(1) = 1$.
- Global minima at $x = -1, f(-1) = -1$

We have no critical points because $f'(x) = 1 \neq 0$. Also not that maxima and minima depend on the chosen domain.

If we take $D_f = [-2, 3]$, then

Max: $x = 3$, Min: $x = -2$

◇

Theorem 11. Let f be continuous with domain D_f . Suppose D_f is closed and bounded, then there is at least one global maxima, and one global minima.




Figure 1: The circle is the boundary.

Some terminology:

$$\text{open} = \{x^2 + y^2 < 1\}$$

$$\text{closed} = \{x^2 + y^2 \leq 1\}$$

The method for finding maxima and minima is as follows:

1. Find critical points of f in D_f , and characterize them.
2. Study the points that are on the boundary.
3. Compare them.

Example. Consider $f(x, y) = x^2 + y^2$ with domain

$$D = \{(x, y) \in \mathbb{R} : x^2 + y^2 \leq 1\}$$

The domain in this case is a disc. The red circle is the boundary.

We compute $f_x = 2x, f_y = 2y$. The only critical point is the origin at $(x_0, y_0) = (0, 0)$. This is a global minimum since $f(0, 0) = 0$ and $f(x, y) \geq 0$.

Now, let's consider the boundary

$$C = \{x^2 + y^2 = 1\}$$

For any point (x_0, y_0) on the circle C , we have

$$f(x_0, y_0) = x_0^2 + y_0^2 = 1$$

We claim that this point is a global maximum. For any (x, y) in domain D_f , we have

$$f(x, y) = x^2 + y^2 \leq 1$$

The value $f(x, y) = 1$ is obtain only at the boundary C . Any point on the circle is a global maxima. \diamond

5.3 Constrained optimization

In this section, we will discuss how to find maxima and minima of $f(x, y)$ with constraint $g(x, y) = 0$. Think of $g = 0$ as a budget, or a geometrical constraint.

Example. We want to minimize $f(x, y) = x^2 + y^2$ with the constraint $g(x, y) = xy - 1 = 0$.




Figure 2: $y = \frac{1}{x}$

We have the following strategy

- Solve $g = 0$ for one variable. For instance $y = \frac{1}{x}$.
- Consider: $h(x) = f\left(x, \frac{1}{x}\right) = x^2 + x^{-2}$

Now we can study this function of one variable with no constraints. We can proceed as usual.

$$\frac{dh}{dx} = 2x - 2x^{-3} = 0$$

This is equivalent to $x^4 = 1$. The real solutions are $x = \pm 1$. Since $y = \frac{1}{x}$, we get the critical points:

$$(x, y) = (1, 1), \quad (x, y) = (-1, -1)$$

\diamond

6 Lecture 6

6.1 Substitution Method

We want to maximize / minimize $f(x, y)$ with constraint $g(x, y) = 0$. We can solve $g(x, y) = 0$ for one variable $y = y(x)$.

Example. Consider $f(x, y) = x^2 + y^2$ and $g(x, y) = xy - 1 = 0$. In this case, we have $f = \frac{1}{x}$ from $g = 0$, we get

$$h(x) = f(x, x^{-1}) = x^2 + x^{-2}$$

We have found the minima at $(1, 1)$ and $(-1, -1)$. ◇

This method isn't always feasible, so let's look at some alternatives.

6.2 Lagrange's Method

Example. Let's look at the level curves, which are circles. We have that $f(x, y) = x^2 + y^2 = c$.





Figure 3: $f(x, y)$

Smaller circles correspond to smaller values of $f(x, y)$, but we must also satisfy $g(x, y) = 0$. In the best case, $f(x, y) = c$ is just touching $g(x, y) = 0$. If this is worked out geometrically, we get $(1, 1)$ and $(-1, -1)$. ◇

This idea is used in Lagrange's method. We want $f(x, y) = c$ to be parallel to $g(x, y) = 0$. More precisely: Their vectors should be parallel.



Equivalently, their normal vectors are also parallel. Recall that a normal vector to $g = 0$ is given by ∇g . Similarly, ∇f is normal to $f = c$.

6.3 Method for Lagrange

Suppose we want to find a local maxima and minima of $f(x, y)$ with constraint $g(x, y) = 0$. We proceed as follows

1. Find all possible solutions to the equations

$$\nabla f(x, y) = \lambda \nabla g(x, y), \quad g(x, y) = 0$$

2. Plug in all solutions from 1. into $f(x, y)$ and identify the largest and smallest.

- The number λ is called the *Lagrange Multiplier*.
- Easy extension to n variables.
- Can also be generalized to multiple constraints, g_1, \dots, g_n

Example. We have the following functions

$$f(x, y) = x^2 + y^2, \quad g(x, y) = xy - 1 = 0$$

We compute the gradients

$$\nabla f(x, y) = (2x, 2y), \quad \nabla g(x, y) = (y, x)$$

The Lagrange equations are

$$2x = \lambda y, \quad 2y = \lambda x, \quad xy - 1 = 0$$

Observe that $(x, y, \lambda) = (0, 0, 0)$ is not a solution. For $x \neq 0$, we get $y = \frac{1}{x}$, from the third equation.

$$\begin{aligned}
 2x &= \lambda y \Rightarrow 2x = \lambda \frac{1}{x} \Rightarrow \lambda = 2x^2 \\
 2y &= \lambda x \Rightarrow \frac{2}{x} = 2x^2 \cdot x \Rightarrow x^4 = 1 \\
 x^4 &= 1 \Rightarrow x = \pm 1, \quad y = \frac{1}{x}
 \end{aligned}$$

$$(x, y) = (1, 1) \quad \wedge \quad (x, y) = (-1, -1)$$

Are these points the minima? We have

$$f(1, 1) = f(-1, -1) = 2$$


Lets Compare this to some other points, such that $g(x, y) = 0$. For instance, $(x, y) = (2, \frac{1}{2})$

$$g\left(2, \frac{1}{2}\right) = 2 \cdot \frac{1}{2} - 1 = 0$$

$$f\left(2, \frac{1}{2}\right) = 2^2 + \left(\frac{1}{2}\right)^2 > 2$$

This tells us that $(x, y) = (\pm 1, \pm 1)$ are local minima. ◇

Example. Consider a box of surface area 24 cm^2 .



Determine the dimensions (x, y, z) such that the volume is max.
We have the surface area $2xy + 2xz + 2yz$. Our constraint is

$$g(x, y, z) = 2xy + 2xz + 2yz - 24 = 0$$

The goal is to maximize $f(x, y, z) = 0$, with the constraint $g(x, y, z) = 0$.
The equation $\nabla f = \lambda \nabla g$ gives

$$yz = 2\lambda(y + 2), \quad xz = 2\lambda(x + 2), \quad xy = 2\lambda(x + y)$$

Observe that $\lambda \neq 0$, since $x, y, z > 0$. To solve the equations, we can multiply by x, y and z , respectively, then we get

$$x(y + z) = y(x + z) = x(y + x)$$

Consider $x(y + z) = y(x + z)$.

$$x(y + z) = y(x + z) \Rightarrow (x - y)z = 0$$

$$z \neq 0, \quad x = y$$

It's the same for the other equations, so $x = y = z$. We also need to use $g = 0$. Setting $x = y = z$, we get

$$g(x, x, x) = 2x^2 + 2x^2 + 2x^2 - 24 = 0$$

$$x^2 = 4$$

$$x = 2$$

Lagrange's method gives:

$$(x, y, z) = (2, 2, 2) \Rightarrow V = 8$$


◇

7 Lecture 7

7.1 Parametrized Curve

A curve is described as a sset of points in \mathbb{R}^2 and \mathbb{R}^3 . For instance a line is described by

$$f = \{(x, y) \in \mathbb{R}^2 : x = y\}$$



This is a static picture. But how do we give a dynamical picture? We'll use parametrized curves.


Definition 12. A parametrization of a curve c in \mathbb{R}^2 , is given by

$$\vec{r}(t) = (x(t), y(t)), \quad t_0 \leq t \leq t_1$$

Such that $\vec{r}(t) \in c$ for all time t .

- A parametrization describes motion. Think of t as the time.
- A parametrization is not unique.
- Various natural assumptions, such as continuity and differentiability.

Example. Consider the funtion $\vec{r}(t)$ with $0 \leq t \leq 1$.



$$\vec{r}(0) = (0, 0), \quad \vec{r}(1) = (1, 1)$$

We have the portion of the line where $y = x$. Notice that here, $x(t) = t$, $y(t) = t$ and $y(t) = x(t) = t$ for all t .

Lets consider a different function,

$$\vec{r}(t) = (2t, 2t), \quad 0 \leq t \leq \frac{1}{2}$$

When we parametrize, we get

$$\vec{r}(0) = (0, 0), \quad \vec{r}\left(\frac{1}{2}\right) = (1, 1)$$

We are moving along the curve at twice the speed.

◇


Example. Given $f(x)$, we consider

$$\vec{r}(t) = (t, f(t)), \quad t_0 \leq t \leq t_1$$

This describes a portion of the graph f , with

$$\text{Start: } (t_0, f(t_0)), \quad \text{End: } (t_1, f(t_1))$$

For instance, consider the line $y = mx + c$, we have



We have that

$$\vec{r}(t) = (t, mt + c), \quad t_0 \leq t \leq t_1$$

◇

Example. We want to describe a line with

$$\text{Start: } A = (x_0, y_0), \quad \text{End: } B = (x_1, y_1)$$

Then we take the parametrization

$$\vec{r}(t) = (1 - t)A + tB, \quad 0 \leq t \leq 1$$


More explicitly, we have

$$\vec{r}(t) = ((1 - t)x_0 + tx_1, (1 - t)y_0 + ty_1)$$

Note that $\vec{r}(0) = A$ and $\vec{r}(1) = B$.

◇

Example. Consider $\vec{r}(t) = (\cos t, \sin t), 0 \leq t \leq 2\pi$. What curve does this describe?



It describes a circle.

$$x(t)^2 + y(t)^2 = (\cos(t)^2 + \sin(t)^2) = 1$$

We start at $(1, 0)$ and move counter-clockwise.

◇


Example. Here is an example from physics. Consider

$$x(t) = v_x t, \quad y(t) = v_y t - \frac{1}{2} g t^2, \quad 0 \leq t \leq \frac{2v_y}{g}$$

This describes the motion of an object with initial velocity $\vec{v} = (v_x, v_y)$, under gravity. We write $t_0 = 0$ and $t_1 = \frac{2v_y}{g}$. Note that

$$\vec{r}(t_0) = (0, 0), \quad \vec{r}(t_1) = \left(\frac{2v_x v_y}{g}, 0 \right)$$

The object falls back to the ground at time t_1 .



Well known fact: This motion is parabolic, we will rederive this.

From $x(t) = v_x t$, we get $t = \frac{x(t)}{v_x}$. Then

$$y(t) = v_y t - \frac{1}{2} g t^2 \Rightarrow \frac{v_x}{v_y} x(t) - \frac{1}{2} \frac{g}{v_x^2} x(t)^2$$

This is the expression of a parabola

$$y = ax^2 + bx + c, \quad a \neq 0$$

It can also be written as

$$y(t) = -\frac{1}{2} \frac{g}{v_x^2} \left(x(t) - \frac{v_y}{g} \right)^2 + \frac{1}{2} \frac{v_y}{g} \frac{v_y}{v_x}$$

◇

7.2 Kinematics

Kinematics describes position, velocity and acceleration of an object.

Definition 13. The position vector is $\vec{r}(t)$. The velocity vector is $\vec{v}(t) \frac{d\vec{r}}{dt}$. The acceleration vector is $\vec{a}(t) = \frac{d^2\vec{r}}{dt^2}$.

If we write $\vec{r}(t) = (x(t), y(t))$, then

$$\vec{v}(t) = \left(\frac{dx}{dt}, \frac{dy}{dt} \right) = (x'(t), y'(t))$$

Similarly

$$\vec{a}(t) = \left(\frac{d^2x}{dt^2}, \frac{d^2y}{dt^2} \right) = (x''(t), y''(t))$$

Example. Consider again the gravity example. Here we have

$$\vec{r}(t) = \left(v_x t, v_y t - \frac{1}{2} g t^2 \right)$$

The velocity is

$$\vec{v}(t) = (v_x, v_y - g t)$$

Note that $v(0) = (v_x, v_y)$ is the initial velocity of the object. For acceleration, we get

$$\vec{a}(t) = \frac{d\vec{v}}{dt} = (0, -g)$$

◇

8 Lecture 8

8.1 Determining Motion

Given acceleration $\vec{a}(t)$, can we find $\vec{v}(t)$ and $\vec{s}(t)$? Yes, with some initial conditions given, we can. This is done by integration, consider

$$\vec{v}(t) = \frac{d\vec{s}}{dt}$$

This is a differential equation for $\vec{s}(t)$. To solve it, we integrate both sides in t , from t_1 , to t_2 . We get

$$\int_{t_0}^{t_1} \vec{v}(t) dt = \int_{t_0}^{t_1} \frac{d\vec{s}}{dt} dt$$

The fundamental theorem of calculus gives

$$\vec{s}(t_1) - \vec{s}(t_0) = \int_{t_0}^{t_1} \vec{v}(t) dt$$

We can determine $\vec{s}(t)$ for any t if we know $\vec{v}(t)$ and the initial condition $\vec{s}(t_0)$.

Example. Consider an object with acceleration

$$\vec{a}(t) = (1, t) = \vec{i} - t\vec{j}$$

We have the following initial conditions

$$\vec{s}(0) = (2, 0) = 2\vec{i} \quad \wedge \quad \vec{v}(0) = 0$$

We want to determine $\vec{s}(t)$. First, to determine $\vec{v}(t)$, we compute

$$\begin{aligned} \int_0^t \vec{a}(t) dt &= \vec{i} \int_0^t 1 dt + \vec{j} \int_0^t t dt \\ &= t\vec{i} + \frac{1}{2}t^2\vec{j} \end{aligned}$$

Here $t_0 = 0$, since $\vec{v}(0) = 0$, then

$$\vec{v}(t) = \vec{v}_0 + \int_0^t \vec{a}(t) dt = t\vec{i} + \frac{1}{2}t^2\vec{j}$$

To determine \vec{s} , we compute

$$\begin{aligned} \int_0^t \vec{v}(t) dt &= \vec{i} \int_0^t t dt + \vec{j} \int_0^t \frac{1}{2}t^2 dt \\ &= \frac{1}{2}t^2\vec{i} + \frac{1}{6}t^3\vec{j} \end{aligned}$$

Since $\vec{s}(0) = (2, 0) = 2\vec{i}$, we get

$$\vec{s}(t) = \vec{s}(0) + \int_0^t \vec{v}(t) dt = \left(\frac{t^2}{2} + 2\right)\vec{i} + \frac{1}{6}t^3\vec{j}$$

◇

8.2 Arc Length

A formula that we can concretely use to compute the length of a curve (using a parametrization). Consider a curve c , with parametrization


$$\vec{s}(t) = (x(t), y(t)), \quad t_0 \leq t \leq t_1$$

Definition 14. The arc length of c is given by

$$S = \int_{t_0}^{t_1} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$


Idea: summing segments of length

$$\sqrt{(\Delta x)^2 + (\Delta y)^2}$$



To compute S , we need a parametrization of c . Does S depend on this choice? No!

Example. Consider the line segment below



From elementary geometry, its length is $\sqrt{1^2 + 1^2} = \sqrt{2}$. Consider the parametrization

$$\vec{s}(t) = (t, t), \quad 0 \leq t \leq 1$$

We have $(x'(t), y'(t)) = (1, 1)$. Then

$$S = \int_0^1 \sqrt{1^2 + 1^2} dt = \sqrt{2} \int_0^1 1 dt = \sqrt{2}$$

Instead we choose

$$\vec{s}(t) = (2t, 2t), \quad 0 \leq t \leq \frac{1}{2}$$

We have $(x'(t), y'(t)) = (2, 2)$. Then

$$S = \int_0^{\frac{1}{2}} \sqrt{2^2 + 2^2} dt = \sqrt{8} \int_0^{\frac{1}{2}} 1 dt = \sqrt{8} \cdot \frac{1}{2} = \sqrt{2}$$

◇

Question: What is the distance crossed up to time t ?

Definition 15. The arc length parameter is

$$S(t) = \int_{t_0}^t \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

The difference is that we integrate up to t , not t_1 . Special case when $S(t_1) = S$. There is an important relation between $S(t)$ and $\vec{v}(t)$.

Proposition 3. We have

$$|\vec{v}(t)| = \frac{dS}{dt}$$

Proof. The fundamental theorem of calculus states that if

$$F(x) = \int_a^x f(x)dt \rightarrow F'(x) = f(x)$$

Applying this to $S(t)$, then

$$S'(t) = \frac{ds}{dt} = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$$


On the other hand, we have that

$$\vec{v}(t) = (x'(t), y'(t)) \quad \wedge \quad |\vec{v}(t)| = (t) = \sqrt{x'(t)^2 + y'(t)^2}$$

The two expressions coincide. □

Example. Consider a circle of radius R , with

$$\vec{S}(t) = (R \cos t, R \sin t), \quad 0 \leq t \leq 2\pi$$



We want to compute $S(t)$, we have

$$\sqrt{x'(t)^2 + y'(t)^2} = \sqrt{R^2 (\sin t)^2 + R^2 (\cos t)^2} = R$$

We want to check that $\frac{dS}{dt} = |\vec{v}(t)|$. We have

$$\vec{v}(t) = \frac{d\vec{S}}{dt} = (-R \sin t, R \cos t)$$


Its length is equal to $|\vec{v}(t)| = R$. Since $S(t) = R(t)$, we see that

$$\frac{dS}{dt} = |\vec{v}(t)|$$

◇

8.3 Tangent Vectors

Geometrically, the velocity $\vec{v}(t)$ is tangent to a curve. It is useful to define a tangent vector of length 1.



Definition 16. The unit tangent vector is


$$\vec{T}(t) = \frac{\vec{v}(t)}{|\vec{v}(t)|}$$

Note that \vec{T} has length 1 since

$$\vec{T}(t) \cdot \vec{T}(t) = \frac{\vec{v}(t) \cdot \vec{v}(t)}{|\vec{v}(t)|^2} = 1$$

8.4 Normal Vectors

Normal vectors are normal to the curve, or in other words, they are orthogonal. Recall that for implicit curves $f(x, y) = 0$, a normal vector is given by ∇f .



Lets now consider parametrised curves. We have

$$S(t) = (x(t), y(t)), \quad t_0 \leq t \leq t_1$$

Definition 17. A unit normal vector to the curve is defined by

$$\vec{N}(t) = \frac{\frac{d\vec{T}}{dt}}{\left| \frac{d\vec{T}}{dt} \right|}$$

We need to check that \vec{N} is orthogonal to \vec{T} , that is: $\vec{N}(t) \cdot \vec{T}(t) = 0$ for all t .

Proposition 4. We have that

$$\vec{N}(t) \cdot \vec{T}(t) = 0$$

Proof. Since \vec{T} is a unit vector, we have that, for all t

$$\vec{T}(t) \cdot \vec{T}(t) = 1$$

Take the time derivative, the $\left(\vec{T} \cdot \vec{T} \right) = 0$. We also have

$$\frac{d}{dt} \left(\vec{T} \cdot \vec{T} \right) = \frac{d\vec{T}}{dt} \cdot \vec{T} + \vec{T} \cdot \frac{d\vec{T}}{dt} = 2 \frac{d\vec{T}}{dt} \cdot \vec{T}$$


Since $\left(\vec{T} \cdot \vec{T} \right) = 0$, we get $\frac{d\vec{T}}{dt} \cdot \vec{T} = 0$. Dividing by $\left| \frac{d\vec{T}}{dt} \right|$, we get $\vec{N}(t) \cdot \vec{T}(t) = 0$ □

9 Lecture 9

9.1 Normal Vectors - Continued

Yesterday we saw that

$$\vec{N}(t) = \frac{\frac{d\vec{T}}{dt}}{\left| \frac{d\vec{T}}{dt} \right|}$$



Example. Consider the circle

$$\vec{r}(t) = (\cos t, \sin t), 0 \leq t \leq 2\pi$$

The velocity is

$$\vec{v}(t) = \vec{r}' = (-\sin t, \cos t)$$

We have $|\vec{v}(t)| = 1$, since

$$\vec{v}(t) \cdot \vec{v}(t) = (\sin^2 t + \cos^2 t) = 1$$

We find that $\vec{T}(t) = \vec{v}(t)$. To find \vec{N} , we need first


$$\frac{d\vec{T}}{dt} = \frac{d\vec{v}}{dt} = (-\cos t, -\sin t)$$

We check that $\left| \frac{d\vec{T}}{dt} \right| = 1$, then

$$\vec{N}(t) = (-\cos t, -\sin t) = -\vec{r}'(t)$$

We compute more explicitly:

$$\vec{N}(t) \cdot \vec{T}(t) = (-\cos t, -\sin t) \cdot (-\sin t, \cos t) = \cos t \cdot \sin t - \sin t \cdot \cos t = 0$$



◇

We revisit the implicit case $f(x, y) = 0$.

Proposition 5. Let C be defined by $f(x, y) = 0$. A unit normal to C is given by $\vec{n} = \frac{\nabla f}{|\nabla f|}$

Proof. Suppose C is parametrized by

$$\vec{r}(t) = (x(t), y(t)), \quad t_0 \leq t \leq t_1$$

By definition of C , we have

$$f(\vec{r}(t)) = f(x(t), y(t)), \quad t_0 \leq t \leq t_1$$

We have that $\frac{d}{dt}f(\vec{r}(t)) = 0$, but using the chain rule, we get

$$\frac{df(\vec{r}(t))}{dt} = \nabla f(\vec{r}(t)) \cdot \frac{d\vec{r}}{dt}$$

Hence, we get

$$\nabla f(\vec{r}(t)) \cdot \vec{v}(t) = 0$$

Therefore $\nabla f(\vec{r}(t))$ is normal to the curve C , or orthogonal to the tangent $\vec{v}(t)$ □

Example. The circle of radius 1 can be described implicitly by $f(x, y) = 0$ with

$$f(x, y) = x^2 + y^2 - 1$$

The gradient is $\nabla f(x, y) = (2x, 2y)$, not of length 1, since

$$|\nabla f(x, y)| = \sqrt{4x^2 + 4y^2} = 2\sqrt{x^2 + y^2}$$


We are only interested in (x, y) such that $f(x, y) = 0$, that is $x^2 + y^2 = 1$, then

$$|\nabla f| = 2$$

Hence we get

$$\vec{n} \frac{\nabla f}{|\nabla f|} = \frac{(2x, 2y)}{2} = (x, y)$$


The normal vectors point outside of the circle, note: $\vec{n} = -\vec{N}$, compare with the parameter.



◇

9.2 Curvature

We want to compute how much a curve "curves".



How do we quantify this?

Definition 18. Let C be a parametrized curve, and \vec{T} its unit tangent vector. The curvature is then defined by

$$K = \left| \frac{d\vec{T}}{dS} \right|$$

Where S is the arc length parameter of C . We also define the radius of curvature as

$$\rho = \frac{1}{K}$$

Note: We consider S , not t . Reason: K does not depend on parametrization.

Example. Consider the following

$$\vec{r}(t) = (0, 0), \quad 0 \leq t \leq 1$$

We expect K to be zero. We can easily compute

$$S(t) = \int_0^t \sqrt{1^2 - 0^2} dt = t$$

Here we have $S(t) = t$. Now we compute \vec{T} . We have $\vec{v}(t) = (1, 0)$, and $|\vec{v}(t)| = 1$. Then $\vec{T}(t) = \vec{v}(t) = (1, 0)$. Furthermore, we get that

$$\frac{d\vec{T}}{dS} = 0 \quad \wedge \quad K = 0$$

◇

Example. Consider a circle of radius R with

$$\vec{r}(t) = (R \cos t, R \sin t), \quad 0 \leq t \leq 2\pi$$

The velocity is

$$\vec{v}(t) = (-R \sin t, R \cos t)$$

We get that $|\vec{v}(t)| = R$, then

$$\vec{T}(t) = \frac{\vec{v}(t)}{|\vec{v}(t)|} = (-\sin t, \cos t)$$

To compute K , we need to express \vec{T} in terms of $\vec{S}(t)$. We saw that

$$S(t) = R \cdot t \Rightarrow t = \frac{S}{R}$$


Now we compute

$$\frac{d\vec{T}}{dS} = \left(-\frac{1}{R} \cos \frac{S}{R}, -\frac{1}{R} \sin \frac{S}{R} \right)$$

Finally, we have

$$K = \left| \frac{d\vec{T}}{dS} \right| = \sqrt{\frac{1}{R^2} \left(\cos \frac{S}{R} \right)^2 + \frac{1}{R^2} \left(\sin \frac{S}{R} \right)^2} = \frac{1}{R}$$

We have non-zero, constant curvature, we also have that $\rho = R$. Note that $K \rightarrow 0$, as $r \rightarrow \infty$



Large Curvature



Small Curvature

◇

Proposition 6. We have that

$$K = \frac{1}{|\vec{v}|} = \left| \frac{d\vec{T}}{dt} \right|$$

Example. Consider again

$$\vec{r}(t) = (R \cos t, R \sin t)$$

We have seen that

$$|\vec{v}(t)|, \vec{T}(t) = (-\sin t, \cos t)$$

To get K , we compute

$$\frac{d\vec{T}}{dt} = (-\cos t, -\sin t)$$

◇

10 Lecture 10

10.1 Curves in polar form

Polar coordinates are an alternative description to cartesian coordinates (x, y) .


Definition 19. The polar coordinates r, θ are defined by

$$x = r \cos \theta, \quad y = r \sin \theta$$

Their range is respectively

$$0 \leq r \leq \infty, \quad 0 \leq \theta \leq 2\pi$$


Geometrical meaning is that r is the distance from the origin, and θ is the angle.



From (x, y) to (r, θ) , we can use

$$r^2 = x^2 + y^2$$

We can describe curves using (r, θ) . The idea is to give r as a function of θ . The curve will be "traced" as we vary θ . It is an analogue of $y = f(x)$.



Example. Consider the curve

$$r(\theta) = 1, \quad 0 \leq \theta \leq 2\pi$$

What curve is it? All points have distance 1 from origin ($r = 1$)

Using $r^2 = x^2 + y^2$, we find that $x^2 + y^2 = 1$. We have a circle of radius 1. \diamond

Example. The next curve is called the cardioid, it is defined by

$$r(\theta) = 1 - \cos \theta, \quad 0 \leq \theta \leq 2\pi$$

It is described in cartesian coordinates by

$$(x^2 + y^2 + x)^2 = x^2 + y^2$$

Polar coordinates work best in the presence of spherical symmetry. The length of C can be computed using polar coordinates. \diamond

Proposition 7. Let C be given in polar form by

$$r = r(\theta), \quad \alpha \leq \theta \leq \beta$$

Then its arc length can be computed by

$$S = \int_{\alpha}^{\beta} \sqrt{\left(\frac{dr}{d\theta}\right)^2 + r^2} d\theta$$

Proof. Parametrize C by

$$\vec{r}(\theta) = (x(\theta), y(\theta)), \quad \alpha \leq \theta \leq \beta$$

where we set

$$x(\theta) = r(\theta) \cos \theta, \quad y(\theta) = r(\theta) \sin \theta$$

The derivatives are, with $r' = \frac{dr}{d\theta}$

$$\frac{dx}{d\theta} = r' \cos \theta - r \sin \theta, \quad \frac{dy}{d\theta} = r' \sin \theta + r \cos \theta$$

After some computation we get

$$(x')^2 + (y')^2 = r^2 + \left(\frac{dr}{d\theta}\right)^2$$

Therefore we get

$$\begin{aligned} S &= \int_{\alpha}^{\beta} \sqrt{(x')^2 + (y')^2} d\theta \\ &= \int_{\alpha}^{\beta} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta \end{aligned}$$

□

Example. We have a circle given by

$$r(\theta) = R, \quad 0 \leq \theta \leq 2\pi$$

We have $\frac{dr}{d\theta} = 0$, then


$$S = \int_0^{2\pi} \sqrt{R^2 + 0^2} d\theta = R \int_0^{2\pi} d\theta = 2\pi R$$

This gives us the circumference of the circle.


◇

10.2 Areas In Polar Form


We want to compute the area inside a closed curve in polar form.



Basic observation:



Has area πr^2



Has area $\frac{\theta}{2\pi} \cdot \pi r^2 = \frac{1}{2} r^2 \theta$

Add small regions with angle $\Delta\theta$ and area $\frac{1}{2} r^2 \Delta\theta$.

Proposition 8. Consider a closed curve described by

$$r = r(\theta), \quad \alpha \leq \theta \leq \beta$$

Then its area is

$$A = \frac{1}{2} \int_{\alpha}^{\beta} r(\theta)^2 d\theta$$

Example. We have a circle of radius R

$$r(\theta) = R, \quad 0 \leq \theta \leq 2\pi$$

we get

$$A = \frac{1}{2} \int_0^{2\pi} R^2 d\theta = \frac{1}{2} R^2 \int_0^{2\pi} d\theta = \pi R^2$$

◇

Example. Consider the cardioid

$$r(\theta) = 1 - \cos \theta, \quad 0 \leq \theta \leq 2\pi$$

The area is given by:

$$\begin{aligned} A &= \int_0^{2\pi} (1 - \cos \theta)^2 d\theta \\ &= \frac{1}{2} \int_0^{2\pi} (1 - 2\cos \theta + (\cos \theta)^2) d\theta \end{aligned}$$

To compute this, we use

$$\int \cos \theta d\theta = \sin \theta + C, \quad \int (\cos \theta)^2 d\theta = \frac{1}{2}\theta + \frac{1}{2}\sin \theta \cos \theta + C$$


Finally, we obtain

$$A = \frac{1}{2} \cdot 2\pi + 0 + \frac{1}{2} \cdot \frac{1}{2} 2\pi = \frac{3}{2}\pi$$

◇

11 Scalar and Vector Fields

Idea 1. A field describes a property of a region R



Mathematically described by

$$f : \mathbb{R}^m \rightarrow \mathbb{R}^n, \quad m \text{ inputs, } n \text{ outputs}$$

For scalar fields the output is a scalar. For vector fields the output is a vector.

Example. The temperature is a scalar field.

$$T : (x, y, z) \rightarrow T(x, y, z)$$

The wind velocity is a vector field

$$\vec{W} : (x, y, z) \rightarrow \vec{W}(x, y, z)$$

◇


Notation. The notation for vector fields:

$$\begin{aligned} \vec{F}(x, y, z) &= (P(x, y, z), Q(x, y, z), R(x, y, z)) \\ &= P(x, y, z)\vec{i} + Q(x, y, z)\vec{j} + R(x, y, z)\vec{k} \end{aligned}$$

P, Q, R are component functions.

In 2D we visualize vector fields by vector plots. For instance, take

$$\vec{F}(x, y) = (0, y)$$



A vector field \vec{F} and a scalar field f can be related as follows

Definition 20. If $\vec{F} = \Delta f$, we say that \vec{F} is a gradient field, and f is a potential.

11.1 Gradient, Divergence and Curl

These are operation defined in terms of the formal vector

$$\nabla = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right) = \frac{\partial}{\partial x} \vec{i} + \frac{\partial}{\partial y} \vec{j} + \frac{\partial}{\partial z} \vec{k}$$

Definition 21. The gradient of a scalar field f is

$$\text{grad } f = \nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right)$$

The output is a vector.

The divergence of a vector field $\vec{F}(P, Q, R)$ is

$$\text{div } \vec{F} = \nabla \cdot \vec{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

The output is a scalar field.

The curl of a vector field \vec{F} is

$$\text{curl } \vec{F} = \nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix}$$

These operations can all be obtained from ∇ .

Operation	Input	Output	Symbol
Gradient	Scalar	Vector	∇f
Divergence	Vector	Scalar	$\nabla \cdot f$
Curl	Vector	Vector	$\nabla \times f$

12 Lecture 11

12.1 Operations on fields

Example. Consider $\vec{F}(x, y, z) = (x^2, y^2, z^2)$, that is

$$P = x^2, \quad Q = y^2, \quad R = z^2$$

Its divergence is

$$\nabla \cdot \vec{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} = 2x + 2y + 2z$$

We can check that \vec{F} is a gradient field. We have that $\vec{F} = \nabla f$, with the potential

$$f(x, y, z) = \frac{1}{3}(x^2 + y^2 + z^2)$$

◇

$\nabla \times \vec{F}$ is computed as a determinant. We can use the cofactor, or the Laplace expansion.

$$\begin{aligned} \nabla \times \vec{F} &= \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \partial_x & \partial_y & \partial_z \\ P & Q & R \end{vmatrix} = \begin{bmatrix} \partial_y & \partial_z \\ Q & R \end{bmatrix} \vec{i} - \begin{bmatrix} \partial_x & \partial_z \\ P & R \end{bmatrix} \vec{j} + \begin{bmatrix} \partial_x & \partial_y \\ P & Q \end{bmatrix} \vec{k} \\ &= \vec{i}(\partial_y R - \partial_z Q) - \vec{j}(\partial_x R - \partial_z P) + \vec{k}(\partial_x Q - \partial_y P) \end{aligned}$$

This is a concrete formula for $\nabla \times \vec{F}$.

Example. Consider $\vec{F} = xy\vec{i} + (x+z)\vec{j} + yz\vec{k}$. We compute

$$\nabla \times \vec{F} = \vec{i}(\partial_y R - \partial_z Q) - \vec{j}(\partial_x R - \partial_z P) + \vec{k}(\partial_x Q - \partial_y P)$$

where

$$P = xy, \quad Q = x + z, \quad R = yz$$


$$\nabla \times \vec{F} = (2-1)\vec{i} + (1-x)\vec{k}$$

◇

Note. Gradient and divergence can be defined in any dimension. The curl is only defined in up to three dimensions.

12.2 Intepretation of Divergence

Think of \vec{F} as the velocity of a fluid. $\nabla \cdot \vec{F}$ at a point P is the amount of fluid entering / leaving a small region around P .



Example. Consider $\vec{F}_1(x, y) = (x, y)$, then

$$\nabla \cdot \vec{F}_1 = \frac{\partial x}{\partial x} + \frac{\partial y}{\partial y} = 2$$

This corresponds to fluid leaving the region. Similarly for $\vec{F}_2 = (-x, -y)$, then

$$\nabla \cdot \vec{F}_2 = \frac{\partial(-x)}{\partial x} + \frac{\partial(-y)}{\partial y} = -2$$

This corresponds to fluid entering the region. Finally consider $\vec{F}_3 = (0, 1)$, then

$$\nabla \cdot \vec{F}_3 = 0$$


This is an equilibrium situation. ◇

12.3 Interpretation of Curl

$\nabla \times \vec{F}$ measures the "rotation" of \vec{F} .

Example. Consider $\vec{F}(x, y, z) = (x^2, 0, 0)$. We expect no rotation, we compute

$$\nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \partial_x & \partial_y & \partial_z \\ x^2 & 0 & 0 \end{vmatrix} = 0$$



◇

Example. Consider $\vec{F}(x, y, z) = (-\omega y, \omega x, 0)$, where ω is a non-zero constant. We compute

$$\nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \partial_x & \partial_y & \partial_z \\ -\omega y & \omega x & 0 \end{vmatrix} = \begin{bmatrix} \partial_x & \partial_y \\ -\omega y & -\omega x \end{bmatrix} \vec{k} = 2\omega \vec{k}$$

$\nabla \times \vec{F} \neq 0$ gives non-zero rotation. Also $\nabla \times \vec{F} = \text{const}$ gives that there is some rotation everywhere.

For a physical interpretation of this, we can write

$$\vec{v} = \vec{F}, \quad \vec{\omega} = (0, 0, \omega), \quad \vec{r} = (x, y, z)$$

Then we can check that $\vec{v} = \vec{\omega} \cdot \vec{r}$. This is the velocity corresponding to the angular velocity $\vec{\omega}$

◇

12.4 Scalar Field from Gradient

Suppose that we know ∇f . Can we recover f ? Yes, up to the initial conditions. The strategy is the following, first we write

$$\vec{F} = \nabla f = (P, Q, R)$$

By the definition of the gradient field, we have

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right)$$

Comparing, we get

$$\frac{\partial f}{\partial x} = P, \quad \frac{\partial f}{\partial y} = Q, \quad , \quad \frac{\partial f}{\partial z} = R$$

Example. Suppose we are given the gradient field

$$\vec{F} = \nabla f = \left(2x + 2y, \frac{1}{2}x^2 + 3y \right)$$

We want to find f , we must have

$$\frac{\partial f}{\partial x} = 2x + 2y, \quad \frac{\partial f}{\partial y} = \frac{1}{2}x^2 + 3y$$

Integrating the first equation in x gives us

$$\begin{aligned} f(x, y) &= \int (2x + xy) dx \\ &= x^2 + \frac{1}{2}x^2y + g(y) \end{aligned}$$

Where $g(y)$ is the integration constant. It can depend on y .
Now we compute $\frac{\partial f}{\partial y}$. We get

$$\frac{\partial f}{\partial y} = \frac{1}{2}x^2 + \frac{dg}{dy}$$

But we also have that $\frac{\partial f}{\partial y} = \frac{1}{2}x^2 + 3y$.
Comparing them, we get

$$\cancel{\frac{1}{2}x^2} + \frac{dg}{dy} = \cancel{\frac{1}{2}x^2} + 3y$$

Then $\frac{dg}{dy} = 3y$. Integrating in y we get

$$g(y) = \int 3y dy = \frac{3}{2}y^2 + C$$

Here, C is a constant. Finally, inserting it, we get

$$\begin{aligned} f(x, y) &= x^2 + \frac{1}{2}x^2y + g(y) \\ &= x^2 + \frac{1}{2}x^2y + \frac{3}{2}y^2 + C \end{aligned}$$

The constant C is usually not important. It can be fixed by an initial condition. For instance $f(0, 0)$ implies that $C = 0$

◇

13 Lecture 12

13.1 Identities Between Operations

We have seen three operations defined by ∇ .

$$\text{Gradient: } \nabla f, \quad \text{Divergence: } \nabla \cdot \vec{F}, \quad \text{Curl: } \nabla \times \vec{F}$$

There are many identities, we'll now look at one.

Proposition 9. For any scalar field f , we have

$$\nabla \times (\nabla f) = 0$$

Proof. We have $\nabla f = (f_x, f_y, f_z)$, then

$$\nabla \times (\nabla f) = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \partial_x & \partial_y & \partial_z \\ f_x & f_y & f_z \end{vmatrix} = (f_{zy} - f_{yz}, -f_{zx} + f_{xz}, f_{yx} - f_{xy})$$


But partial derivatives can be exchanged. Then we find that $\nabla \times (\nabla f) = 0$ □

We are going to use this when we discuss conservative fields.

13.2 Line Integrals

Consider the curve C with

$$\vec{r}(t) = (x(t), y(t)), t_0 \leq t \leq t_1$$



$$\Delta x = x(t + \Delta t) - x(t), \quad \Delta y = y(t + \Delta t) - y(t)$$

The distance between the two points is

$$\Delta S = \sqrt{(\Delta x)^2 + (\Delta y)^2}$$

When Δt is small, we can use the following approximation

$$\Delta x \approx \frac{dx}{dt} \Delta t, \quad \Delta y \approx \frac{dy}{dt} \Delta t$$

Then, for the distance, we get

$$S = \sqrt{\left(\frac{dx}{dt} \Delta t\right)^2 + \left(\frac{dy}{dt} \Delta t\right)^2} = \Delta t \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$$

Letting $\Delta t \rightarrow 0$ leads to the following


Definition 22. The line integral of $f(x, y)$ along a curve C is defined by

$$\int_C f ds = \int_{t_0}^{t_1} f(x(t), y(t)) \sqrt{x'(t)^2 + y'(t)^2} dt$$

Observe. Note that f is restricted to $\vec{r}(t) = (x(t), y(t))$. When $f = 1$, we recover the arc length.

Example. Consider C defined by

$$x(t) = t, \quad y(t) = 0, \quad 0 \leq t \leq 1$$



First, we compute

$$\sqrt{x'(t)^2 + y'(t)^2} dt = \sqrt{1^2 + 0^2} = 1$$

Now, consider $f(x, y) = x^2 + y$. We want to compute $\int_C f ds$. Restricting f to C gives


$$f(x(t), y(t)) = x(t)^2 + y(t)^2 = t^2 + 0 = t^2$$

Then we obtain

$$\int_C f ds = \int_0^1 t^2 \cdot 1 dt = \frac{1}{3}$$

◇

Note. Line integrals can be used to compute the mass of a 1-dimensional object. The curve C describes the object, and the function $\int_C f ds$ is the mass.



13.3 Parametrization and Orientation

The next result is as for the arc length.

Proposition 10. The integral $\int_C f ds$ does not depend on the parametrization of C .

We will consider a special case


Example. Consider C with

$$\vec{r}(t) = (x(t), y(t)), t_0 \leq t \leq t_1$$

We have

$$\text{Start: } A = (x(t_0), y(t_0)), \quad \text{End: } B = (x(t_1), y(t_1))$$

We want to go from B to A .



We can do this in the following manner

$$\vec{r}_{\text{opp}} = (x(-t), y(-t)), \quad -t_1 \leq t \leq -t_0$$

Note that

$$\vec{r}_{\text{opp}}(-t_1) = (x(t_1), y(t_1)) = B$$

$$\vec{r}_{\text{opp}}(-t_0) = (x(t_0), y(t_0)) = A$$

◇

Example. Consider the segment C from $(0, 0)$ to $(1, 0)$. Take $f(x, y) = x$. Show that $\int_C f ds = \frac{1}{2}$ using $\vec{r}(t)$ and $\vec{r}_{\text{opp}}(t)$.

If C parametrized by $\vec{r}(t)$, we use $-C$ when considering $\vec{r}_{\text{opp}}(t)$. We have

$$\int_C f ds = \int_{-C} f ds$$

The situation will be different for vector fields.

◇

13.4 Case of Vector Fields

Consider the curve with

$$\vec{r}(t) = (x(t), y(t)), \quad t_0 \leq t \leq t_1$$

$$\vec{r}'(t) = (x'(t), y'(t)) \quad (\text{Velocity vector})$$

Definition 23. The line integral of \vec{F} along C is

$$\int_C \vec{F} \cdot d\vec{r} = \int_{t_0}^{t_1} \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt$$

Here $\vec{F}(\vec{r}(t)) = \vec{F}(x(t), y(t))$

In physics, we have that \vec{F} is the force, and $\int_C \vec{F} d\vec{r}$ is the work done by \vec{F} along C . The elementary case is given by $W = FS$, or work = force \cdot displacement. More explicitly, we consider

$$\vec{F}(x, y) = (P(x, y), Q(x, y))$$

Then we have

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = P(x(t), y(t)) \cdot x'(t) + Q(x(t), y(t)) \cdot y'(t)$$

The line integral is then


$$\int_C \vec{F} \cdot d\vec{r} = \int_{t_0}^{t_1} P(x(t), y(t)) \cdot x'(t) dt + \int_{t_0}^{t_1} Q(x(t), y(t)) \cdot y'(t) dt$$

We will write the expression as

$$\int_C \vec{F} \cdot d\vec{r} = \int_C P dx + \int_C Q dy$$

Example. Consider the curve C with

$$\vec{r}(t) = (t, t^2), \quad 0 \leq t \leq 1$$



We have $x(t) = t$ and $y(t) = t^2$, its derivative is

$$\vec{r}(t) = (x'(t), y'(t)) = (1, 2t)$$

Now consider the vector field

$$\vec{F}(x, y) = (x + y, x)$$

That is $P = x + y$ and $Q = x$, when this is restricted to C , we get

$$\vec{F}(\vec{r}(t)) = (x(t) + y(t), y(t)) = (t + t^2, t)$$

We want to compute

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}' = (t + t^2, t) \cdot (1, 2t) = (3t^2 + t)$$

We finally obtain

$$\int_C \vec{F} \cdot d\vec{r} = \int_0^1 (3t^2 + t) dt = \frac{3}{2}$$

◇

14 Line Integrals, Parametrization and Vector Fields

14.1 Identities between operations

We have seen three operations defined by ∇ .

Gradient: ∇f

Divergence: $\nabla \cdot \vec{F}$

Curl: $\nabla \times \vec{F}$

There are many of them, we look at only one.

Proposition 11. For any scalar field f we have

$$\nabla \times (\nabla f) = 0$$

Proof. We have $\nabla f = (f_x, f_y, f_z)$. Then

$$\nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \partial_x & \partial_y & \partial_z \\ f_x & f_y & f_z \end{vmatrix} = \vec{i}(f_{zy} - f_{yz}) - f_{zx} + f_{xz}, f_{yx} - f_{cy})$$

But partial derivatives can be exchanged, $f_{xy} = f_{yx}$. Then we find that

$$\nabla \times (\nabla f) = 0$$

□

We are going to use this when we discuss conservative fields.

14.2 Line Integrals

Consider the curve C with

$$\vec{r}(t) = (x(t), y(t)), \quad t_0 \leq t \leq t_1$$

Consider the points $\vec{r}(t)$ and $\vec{r}(t + \Delta t)$ where Δt is small.

$$\Delta x = x(t + \Delta t) - x(t), \quad \Delta y = y(t + \Delta t) - y(t)$$

The distance between the two points is

$$\Delta S = \sqrt{(\Delta x)^2 + (\Delta y)^2}$$

When Δt is small, we can use the following approximation

$$\Delta x \approx \frac{dx}{dt} \Delta t, \quad \Delta y \approx \frac{dy}{dt} \Delta t$$

Then, for the distance, we get

$$S = \sqrt{\left(\frac{dx}{dt} \Delta t\right)^2 + \left(\frac{dy}{dt} \Delta t\right)^2} = \Delta t \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$$

Definition 24. The line integral of $f(x, y)$ along a curve C is defined by

$$\int_C f dS = \int_{t_0}^{t_1} f(x(t), y(t)) \sqrt{x'(t)^2 + y'(t)^2} dt$$

Some observations:

- Note that f is restricted to $\vec{r}(t) = (x(t), y(t))$.
- When $f = 1$, we recover the arc length.

Example. Consider C defined by

$$x(t), y(t) = 0, \quad 0 \leq t \leq 1$$

First, we compute

$$\sqrt{x'(t)^2 + y'(t)^2} dt = \sqrt{1^2 + 0^2} = 1$$

◇

Now, consider $f(x, y) = x^2 + y$. Compute $\int_C f dS$. Restricting f to C gives

$$f(x(t), y(t)) = x(t)^2 + y(t) = t^2 + 0 = t^2$$

Then we obtain

$$\int_C f dS = \int_0^1 t^2 \cdot 1 dt = \frac{1}{3}$$

- Line integrals can be used to compute the mass of an objects (1 - dimensional)
- The curve C describes the object
- The function $\int_C f dS$ is the mass.

14.3 Parametrization and orientation

Next results as for the arc length.

Proposition 12. The integral $\int_C f dS$ does not depend on the parametrization of C .

We will consider a special case.

Example. Consider C with

$$\vec{r}(t) = (x(t), y(t)), \quad t_0 \leq t \leq t_1$$

$$\text{Start: } A = (x(t_0), y(t_0)), \quad \text{End: } B = (x(t_1), y(t_1))$$

We want to go from B to A, which can be done in the following manner

$$\vec{r}_{\text{opp}} = (x(-t), y(-t)), \quad -t_1 \leq t \leq -t_0$$

Note that

$$\vec{r}_{\text{opp}}(-t_1) = (x(t_1), y(t_1)) = B$$

$$\vec{r}_{\text{opp}}(-t_0) = (x(t_0), y(t_0)) = A$$

◇

Example. Consider C = segment from $(0,0)$, to $(1,0)$, take $f(x, y) = x$. Show that $\int_C f dS = \frac{1}{2}$ using $\vec{r}(t)$ and $\vec{r}_{\text{opp}}(t)$. ◇

If C parametrized by $\vec{r}(t)$, we will use $-C$ when considering $\vec{r}_{\text{opp}}(t)$. We have that

$$\int_C f dS = \int_{-C} f dS$$

The situation will be different for vector fields.

14.4 Case of Vector Fields

Consider the curve with

$$\vec{r}(t) = (x(t), y(t)), \quad t_0 \leq t \leq t_1$$

$$\vec{r}'(t) = (x'(t), y'(t)) \quad \text{Velocity Vector}$$

Definition 25. The line integral of \vec{F} along C is

$$\int_C \vec{F} \cdot d\vec{r} = \int_{t_0}^{t_1} \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt$$

Here $\vec{F}(\vec{r}(t)) = \vec{F}(x(t), y(t))$

- In physics, we have that \vec{F} = Force and $\int_C \vec{F} d\vec{r}$ = Work done by \vec{F} along C .
- The elementary case $W = FS$, where W is work, F is Force and S is the displacement.

More explicitly, we consider

$$\vec{F}(x, y) = (P(x, y), Q(x, y))$$

Then we have that

$$\vec{F}(\vec{r}(t) \cdot \vec{r}'(t)) = P(x(t), y(t)) \cdot x'(t) + Q(x(t), y(t)) \cdot y'(t)$$

The line integral is then

$$\int_C \vec{F} \cdot d\vec{r} = \int_{t_0}^{t_1} P(x(t), y(t)) \cdot x'(t) dt + \int_{t_0}^{t_1} Q(x(t), y(t)) \cdot y'(t) dt$$

We will write the expression as

$$\int_C \vec{F} d\vec{r} = \int_C P dx + \int_C Q dy$$

Example. Consider a curve C with

$$\vec{r}(t) = (t, t^2), \quad 0 \leq t \leq 1$$

We have $x(t) = t$ and $y(t) = t^2$, its derivative is

$$\vec{r}'(t) = (x'(t), y'(t)) = (1, 2t)$$

Now consider the vector field

$$\vec{F}(x, y) = (x + y, x)$$

That is $P = x + y$ and $Q = x$. When restricted to C , we get

$$\vec{F}(\vec{r}(t)) = (x(t) + y(t), y(t)) = (t + t^2, t)$$

We want to compute

$$\vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) = (t + t^2, t) \cdot (1, 2t) = (3t^2 + t)$$

We then obtain the following

$$\int_C \vec{F} \cdot d\vec{r} = \int_0^1 (3t^2 + t) dt = \frac{3}{2}$$

◇