Proceedings of Spring 2026 MECE E3311
Heat Transfer
Mechanical Engineering
Columbia University

Cryotherapy and Its Significance In Medical Treatment

Team #1
Patrick Adah, Hudson Aibel, Kristina Dee, Enrique Lopez Sanchez, Prutha Patil, Yuxi Rui

***Abstract*—Cryotherapy is the application of low temperatures to biological tissues, and encompasses a wide range of clinical and wellness techniques such as whole-body cryotherapy, cold-water immersion, and cryosurgery. Despite this range, they all share the same fundamental mechanisms of heat transfer —conduction at the instrument-body interface, and convection between the body and its surroundings. This paper reviews the heat transfer mechanisms that underly cryotherapy including the Pennes Bioheat Equation and latent heat phase change in freezing tissue. The paper also discusses types and applications of cryotherapy treatment that include cryoablation, whole-body cryotherapy, local cryotherapy, and cold-water immersion, and extends this discussion to examine some possible improvements in cryo-therapeutic treatments.**

I. Introduction

To many, the term cryotherapy often refers to the application of cold substances, such as ice packs, to help reduce pain and inflammation resulting from tissue discomfort. This definition, while accurate, lacks the coverage of what ”cold substances” really encompasses.

In medical terms, cryotherapy is defined as the use of very low temperatures to treat tissue, reduce pain, or control inflammation. In medicine, it ranges from simple ice-pack treatment to liquid-nitrogen destruction of abnormal skin tissue.

While modern cryotherapy encompasses a wide range of clinical and wellness techniques, its underlying principle is consistent: controlled removal of heat from the body to achieve a desired physiological or therapeutic outcome. Cryotherapy is fundamentally a heat transfer problem. The body acts as a thermal system, and cold application drives heat flux away from tissue, inducing responses from the body [1].

The use of cryotherapy is not a new phenomenon. As early as 2500 BCE, Egyptian physicians documented the use of cold compresses to reduce inflammation and manage injury. Hippocrates and later Roman physicians employed snow and ice to slow bleeding and numb pain before surgical procedures [2]. However, cryotherapy in its modern form emerged in the late 19th century when Dr. James Arnott used a mixture of ice and salt (capable of reaching approximately -20°C) to treat tumors of the cervix and breast. The result was both analgesic and tumor-reducing effects. This represented the first documented use of extreme cold for targeted tissue destruction, laying the groundwork for what would become cryosurgery [1].

The mid-20th century brought significant technical advancement. The development of liquid nitrogen as a practical cryogen (with a boiling point of -196°C) gave doctors a powerful and relatively accessible tool for delivering intense localized cold. Irving Cooper’s introduction of the cryoprobe in the 1960s enabled precise application of these temperatures to internal tissues, expanding cryotherapy into neurosurgery and oncology [3] From a heat transfer perspective, the cryoprobe operates by circulating liquid nitrogen or another refrigerant through a probe tip, establishing a steep thermal gradient between the probe surface and the surrounding biological tissue. This drives rapid conductive heat transfer, forming an iceball whose growth and geometry can be predicted and controlled [4].

Cryotherapy expanded beyond just the surgical field. Localized surface cryotherapy, such as ice packs, cold water immersion, and coolant sprays, became standard practice in sports medicine and physical rehabilitation throughout the 20th century. These methods rely primarily on conduction and convection to extract heat from superficial tissue, reducing local metabolic rate, nerve conduction velocity, and inflammatory activity [5]. Whole-body cryotherapy, known as WBC, was introduced in Japan by Dr. Toshima Yamauchi in the late 1970s for rheumatoid arthritis treatment. It extended the concept of cryotherapy to systemic exposure. Patients enter a chamber cooled by liquid nitrogen vapor to temperatures between -110°C and -160°C for short durations of two to four minutes. Unlike other localized methods, WBC relies heavily on convective heat transfer from the skin surface to the surrounding cold gas, triggering a systemic response from the body [6].

Understanding cryotherapy through the lens of heat transfer requires accounting for the complex thermal behavior of biological tissue. Unlike simple homogeneous materials, tissue conducts heat anisotropically and is continuously heated from within by blood perfusion and cellular metabolism [7]. Blood flow acts as a convective heat source, warm arterial blood delivers thermal energy to cooled regions, opposing the cooling effect and playing a critical role in determining how quickly and deeply temperature drops penetrate tissue. This is formally captured by the Pennes Bioheat Equation, which models the transient temperature distribution in tissue as a function of conduction, blood perfusion, and metabolic heat generation.

Phase change adds an additional layer of complexity. When tissue water freezes, it undergoes a liquid-to-solid transition that releases latent heat, temporarily slowing the rate of temperature decrease at the freezing front [8]. This phenomenon is central to cryosurgery, where the controlled propagation of an ice front through tissue is the primary mechanism of cell destruction. Accurate prediction of iceball growth and the extent of lethal tissue damage depends on proper modeling of this phase change.

Together, all of the different principles of heat transfer: conduction, convection, radiation, blood perfusion, metabolic heating, and latent heat of phase change form the backbone of

all of cryotherapy, whether that be treating a sprained ankle or attacking a metastatic cancer.

II. Heat Transfer Processes Involved In Cryotherapy

A. Conduction

Conduction is a method of heat transfer that relies on contact between two surfaces, often being the primary source to localized treatments such as ice packs and cryoprobes. It is governed by Fourier's Law, which shows how heat flux (q) is proportional to the temperature gradient (∇T) and the thermal conductivity of the material (k):

$$q = -k\nabla T \tag{1}$$

For cryotherapy, the temperature gradient is established between internal body temperatures and the external cryogenic agent at the skin's surface. However, the efficiency of the heat transfer process is highly limited due to tissue layers [17].

The human body is made up of a non-homogeneous, multi-layered system that consists of epidermis, dermis, adipose tissue, and skeletal muscle. Each layer has a different thermal conductivity that affects the body's ability to remove heat. The most significant issue is the thickness of the adipose layer (fat). This tissue has a thermal conductivity of $0.23\frac{W}{mK}$ which is significantly lower compared to muscle tissue ($k = 0.46\frac{W}{mK}$). Its material properties cause the tissue to act as an insulator [18]. As a result, people with higher adipose concentrations require longer exposure time to therapeutic temperatures. Research suggests an inversely proportional relationship between the heat loss rate in intramuscular tissue and the thickness of fat [17], [19].

It is important to consider the non-linearity of these thermal properties within organic tissue. As cryotherapy progresses and temperatures decline, thermal conductivity tends to increase. Frozen tissue is shown to have higher thermal conductivity than non-frozen tissue, creating a freezing front within the body [20]. To accurately model this, k must be treated as a temperature dependent variable:

$$k(T) = \begin{cases} k_s & T < T_s \\ k_s + (k_l - k_s)\frac{T-T_s}{T_l-T_s} & T_s \leq T \leq T_l \\ k_l & T > T_l \end{cases} \tag{2}$$

where k_s is thermal conductivity of the solid state, k_l is the thermal conductivity of the liquid state, and T_l and T_s represent the temperatures at the liquid and solid states, respectively [22].

Modeling becomes increasingly complex as penetration depth requires solving the heat equation for a non-stationary temperature field. This shift in analytical procedure accounts for the change in k-values as intracellular water begins to shift phases (Equation 2) [21].

B. Convection

To further expand the process of removing heat from the body, the energy must transfer away from the skin to the environment. This is where convection comes in, allowing heat to move into the surrounding fluid medium, typically through cold air, liquid nitrogen, or water. The process of convection is governed by Newton's Law of Cooling:

$$q = h(T - T_\infty) \tag{3}$$

where h is the heat transfer coefficient of the fluid, T is the temperature of the skin, and T_∞ is the temperature of the bulk fluid. Equation 3 represents the heat flux from the skin to the ambient.

Unlike conduction, convection is reliant on the fluid's flow properties rather than material properties. In Whole-Body Cryotherapy (WBC), fans are used to create forced convection to significantly increase the value of h compared to natural convection. From fluid mechanics, h is inversely proportional to the thickness of the thermal boundary layer at the skin-fluid interface. Therefore, the addition of the fan reduces the thickness of the thermal boundary layer and rapidly cools the epidermis [23].

The selection of the cooling medium is also very important due to the difference in h. The convective heat transfer coefficient for air in a typical cryochamber can range from 15 to $40\frac{W}{m^2K}$, depending on the flow of air [24]. Water, on the other hand, has a coefficient above $250\frac{W}{m^2K}$ in ice baths [7]. The difference in coefficient of heat transfer is why a $10°$C ice bath results in significantly higher heat extraction than -$110°$C air in a cryogenic chamber [6].

C. Latent Heat

Latent heat is a form of energy that accounts for a phase change. In medicinal cryotherapy, tissue will freeze, causing intracellular and extracellular water to transition from liquid to solid. This process releases a significant amount of energy, at approximately $334\frac{kJ}{kg}$ and must occur at a constant temperature. Therefore, a thermal plateau is created and the cooling rate of the tissue stalls until the phase change is complete [25].

In cryoablations, the phase change accounts for the morphology of ice balls. However, accurate models must be completed to ensure that lethal temperatures are focused on the target volume to spare surrounding healthy tissue. The lethal isotherm is typically below -$40°$C [26]. To mathematically handle this process, the Enthalpy Method is incorporated by reformulating the heat equation to include latent heat as a function of total energy instead of temperature alone.

$$\rho\frac{\partial H}{\partial t} = \nabla \cdot (k\nabla T) + \omega_b\rho_b c_b(T_a - T) + Q_m \tag{4}$$

Equation 4 allows for the tracking of a non-linear moving boundary between two zones of different phases. This is critical for predicting the depth of necrosis [22].

Alternatively, the process can be modeled by the Effective Heat Capacity (C_{eff}) which is typically used for Finite Element Analysis (FEA):

$$\rho C_{eff}(T)\frac{\partial T}{\partial t} = \nabla \cdot (k\nabla T) \tag{5}$$

where

$$C_{eff} = c_p + L\frac{df}{dT} \tag{6}$$

In Equation 6, c_p is the sensible specific heat of the tissue, L is the latent heat of fusion, and $\frac{df}{dT}$ is the liquid fraction (f) with respect to temperature [27]. This treats the latent heat as a temporary spike in the material's specific heat during the phase change.

D. Pennes Bioheat

The Pennes Bioheat Equation was developed by Harry H. Pennes in 1948 to model the temperature distribution in human tissue, modifying the heat conduction equation (Equation 1). The method includes the circulatory system, which serves as an "internal radiator".

$$\rho c \frac{\partial T}{\partial t} = \nabla \cdot (k \nabla T) + \omega_b \rho_b c_b (T_a - T) + Q_m \qquad (7)$$

Equation 7 incorporates biological factors where ω_b is the blood perfusion rate (s^{-1}), ρ_b is the density of blood, and c_b is the specific heat of blood. The factors of heat include T_a which represents the arterial blood and typically represented by core body temperature, and Q_m, the volumetric metabolic heat generation [7].

1) Blood Perfusion: The term $\omega_b \rho_b c_b (T_a - T)$ represents a convective heat exchange between blood and tissue. This acts as a volumetric heat source and as the cooling agent extracts heat, the body will pump arterial blood to affected area to conserve the core body temperature.

Blood perfusion has a heat sink effect that limits the growth of ice balls during cryoablation. This is especially true near major blood vessels, where blood flow is likely to extract heat at a much more significant rate than the probe can freeze tissue. This effect has the potential to keep malignant cells alive [28].

2) Metabolic Heat Generation (Q_m): Q_m accounts for the energy that is produced by cellular respiration. However, the effects are often negligible because it is often several orders of magnitude smaller than the other components. Therefore, Q_m is typically treated as a constant that can be omitted for simplification but is still included for energy balance [22], [27].

III. Types And Applications of Cryotherapy Treatment

Cryotherapy may be categorized according to scale of tissue exposure and method of bodily heat extraction. Major forms include local cryotherapy, cryoablation (cryosurgery), whole-body cryotherapy, and cold-water immersion. Although such modalities rely on a reduction of tissue temperature, they differ heavily in their dominant heat-transfer mechanisms, achievable cooling depth, and clinical purpose.

A. Local Cryotherapy

Local cryotherapy is the application of cold to a specific body region, using cooling agents such as ice packs, cooling pads, or liquid nitrogen (vapocoolant) sprays [9]. Such is commonly used in sports medicine, physical therapy, and acute injury management to reduce pain, edema, inflammation, and muscle spasms.

From a heat transfer perspective, conduction is the dominant mechanism in most local cryotherapy techniques, requiring direct contact between the cold source and skin. Heat flows from the warmer body tissue to the colder external medium according to Fourier's Law. In spray-based applications, convection also contributes by transporting heat away from the tissue surface via cold gas or liquid carriers [10]. These processes were proven to reduce tissue temperature, blood flow, inflammation, and nerve conduction velocity, leading to pain relief and swelling reduction. The rate and depth of tissue cooling depend strongly on the temperature difference, duration of exposure, contact quality, and thermal resistance of intervening tissue layers (most especially subcutaneous fat).

Clinically, local cryotherapy is widely used for acute musculoskeletal injuries including sprains, strains, and contusions, as well as for postoperative pain control and rehabilitation support. Local metabolic demand and nerve conduction velocity both decrease with tissue temperature, thus helping limit secondary hypoxic tissue injury after trauma.

B. Cryoablation (Cryosurgery)

Cryosurgery, also referred to as cryoablation, makes the use of extremely cold substances such as liquid nitrogen, nitrous oxide, and argon-based systems (dependent on whether treatment is applied superficially or via probe) to intentionally destroy pathological tissue including warts, actinic keratoses, and superficial skin cancers as well as in interventional medicine for treatment of tumors in the prostate, liver, kidney, lung, and bone. [11].

During treatment, heat rapidly transfers from tissue to the cryogenic applicator through conduction. In spray cryotherapy, forced convection further enhances surface cooling. As tissue temperature drops below freezing, phase change occurs causing intracellular and extracellular water to freeze, forming ice crystals that damage cellular structures. This introduces latent heat effects that heavily influence the thermal history of the tissue and the growth of the freezing front. Repeated freeze-thaw cycles are typically employed to increase cellular injury through such crystal formation, osmotic stress, membrane rupture, and vascular stasis [12].

Application-wise, cryoablation proves useful for its highly localized destruction of diseased tissue while minimizing damage to surrounding structures, with its effectiveness dependent on accurate prediction of iceball geometry, lethal isotherm location, and the thermal effects of blood perfusion, which may function as a heat sink and reduce freezing near large vessels.

C. Whole-Body Cryotherapy

Whole-body cryotherapy (WBC) exposes nearly the entire body to extremely cold air, typically between $-110°$C and $-160°$C for short durations of approximately 2-4 minutes inside a cryochamber. It is commonly marketed for exercise recovery, pain relief, soreness reduction, and temporary inflammatory relief, and has been more recently explored for conditions such as rheumatoid arthritis and chronic pain.

Unlike local cryotherapy and cryoablation, WBC relies primarily on convective heat transfer as circulating cold gas removes heat from the skin surface. Since air has a relatively low heat-transfer coefficient (in comparison to water), WBC can expose the body to extremely low ambient temperatures while limiting deeper tissue cooling, rapidly cooling the skin surface rather than in deep muscle layers. Limited radiative heat loss also occurs because of the large temperature difference between the body and surroundings[6]. The rapid reduction in skin temperature may decrease inflammation and temporarily reduce muscle soreness and pain perception [13]. However, scientific evidence supporting long-term therapeutic benefits remains limited [14]. WBC is particularly interesting in its demonstration that very low air temperatures do not necessarily correspond to greater total heat remove than colder liquid-based methods, due to the relatively small convective heat-transfer coefficient of air.

D. Cold-Water Immersion

Lastly, cold-water (CWI) which is commonly known as ice baths or cold plunges is widely used in sports medicine and rehabilitation to reduce soreness and inflammation after exercise or injury [15]. Heat is removed from the body mainly through convection (particularly in a liquid medium), with circulation of cold water continuously absorbing heat from the skin more effectively than cold air, with additional conduction occuring at the skin–water interface [16]. This process leads to a reduction in tissue temperature, decreasing metabolic activity, inflammation, and nerve conduction velocity, contributing to pain reduction and recovery support. Since water has a much larger heat transfer coefficient and volumetric heat capacity than air, it extracts body heat far more efficiently even at much higher temperatures than those used in cryochambers. This makes CWI one of the most thermally effective noninvasive cryotherapy methods for lowering skin and superficial muscle temperature. However, the stronger cooling effect necessitates careful control of exposure duration and water temperature to avoid adverse physiological responses.

IV. Possible Improvements in Cryotherapy

In cryotherapeutic cold chambers, liquid nitrogen remains the most popular cryogen due to its low cost and wide availability [29]. However, there is a recent shift to electric cryotherapy chambers instead of nitrogen- and argon-based chambers for improved safety, such as reduced risk of cold burns and anoxia [30]. Studies have shown that mechanical refrigeration systems based on the Joule-Thomson effect can achieve comparable therapeutic temperatures to traditional nitrogen systems, while maintaining safer oxygenated environments [31]. Unlike nitrogen chambers in which the head must remain outside of the cooling enclosure due to risk of asphyxiation, electric systems allow the full body to be cooled through circulating oxygenated air. Using multiple fans placed at various heights to create forced convection, electric cryotherapy chambers cool the body more uniformly than nitrogen chambers which directly pump in liquid nitrogen, leading the lower half of the chamber to be cooler than the upper [32]. Whole body cooling is shown to stimulate the autonomous nervous system [33] and specifically increase parasympathetic activity [34], which can improve immune response and reduce stress.

Due to associated risks such as frostbite, skin burns, and nerve damage, improving temperature regulation and real-time monitoring are important areas for further cryotherapy research. Recent studies have explored measuring bioimpedance [35] to detect abnormal freezing events during treatment, and incorporating artificial intelligence into thermal imaging for continuous feedback and automated temperature adjustments [36].

Another area of improvement is decreasing the size and weight of cryotherapeutic devices for increased portability. New designs of thermoelectric cooling (TEC) units bring the refrigeration source directly to the tissue to be treated, improving upon current devices that circulate ice water from a remote insulated container to a cooling bladder placed on the skin [37]. By reducing the device size and energy requirements, these devices are especially useful on-site for athletes or in outdoor spaces where rapid treatment is needed.

In the field of cryoablation, nanomaterials are increasingly being investigated to improve the uniformity and precision of cancer treatments [38]. Nanomaterials can be used as imaging contrast agents to guide cryosurgeries, and also as vehicles for drug delivery. Additionally, specific nanomaterials can generate reactive oxygen species (ROS) from radiation or heat from light, increasing oxidative stress and inducing tumor cell death. Current cryoablation limitations include incomplete destruction of tumors especially on the periphery of ablation zones, restrictions on tumor size, inadequate ice formation, and possibility of ice extending into healthy tissues. Nanomaterials can help address this by improving control over ice formation time and shape. Metal nanoparticles, such as gold, iron oxide, and gallium-based liquid alloys, have a higher thermal conductivity and can exchange heat more efficiently, permeating throughout the tumor and creating a more uniform thermal gradient [36]. In contrast, non-metal nanoparticles have a lower thermal conductivity and can be used to protect healthy cells from cryoinjury, such as adding phase change materials (PCMs) that can buffer against temperature fluctuations.

V. Conclusion

This paper looks at past developments of cryotherapy, especially early works in the mid 20th century, and how different heat transfer processes play a critical role. The common heat transfer processes include Conduction (Fourier's Law), Convection (Newton's Law of Cooling), Latent Heat (Phase Change), and Pennes Bioheat (physiological application of conduction). Future areas of research include transitioning to electric cryochambers from liquid nitrogen-based chambers, advancements in temperature regulation and monitoring, increased compactness of devices for portability and everyday use, and the use of nanomaterials for more targeted cryoablation applications.

Cryotherapy has evolved into a highly accessible treatment method because its various forms range from simple household

applications, such as ice packs and cold-water immersion, to advanced clinical procedures like cryosurgery and whole-body cryotherapy. Its ability to provide rapid pain relief, reduce inflammation, and assist tissue recovery makes it valuable for individuals experiencing physical distress or injury.

REFERENCES

[1] Gage AA. History of cryosurgery. *Semin Surg Oncol*. 1998 Mar;14(2):99-109. doi: 10.1002/(sici)1098-2388(199803)14:2¡99::aid-ssu2¿3.0.co;2-1. PMID: 9492880.

[2] Freiman A, Bouganim N. *History of cryotherapy*. Dermatol Online J. 2005;11(2):9. PMID: 16150211.

[3] Cooper IS. *Cryogenic surgery*: a new method of destruction or extirpation of benign or malignant tissues. N Engl J Med. 1963;268:743-749. doi: 10.1056/NEJM196304042681401.

[4] Baust JG, Gage AA. *The molecular basis of cryosurgery*. BJU Int. 2005;95(9):1187-1191. doi: 10.1111/j.1464-410X.2005.05502.x. PMID: 15892793.

[5] Bleakley C, McDonough S, MacAuley D. *The use of ice in the treatment of acute soft-tissue injury: a systematic review of randomized controlled trials*. Am J Sports Med. 2004;32(1):251-261. doi: 10.1177/0363546503260757. PMID: 14754753.

[6] Bleakley CM, Bieuzen F, Davison GW, Costello JT. *Whole-body cryotherapy: empirical evidence and theoretical perspectives*. Open Access J Sports Med. 2014;5:25-36. doi: 10.2147/OAJSM.S41655. PMID: 24648781.

[7] Pennes HH. *Analysis of tissue and arterial blood temperatures in the resting human forearm*. J Appl Physiol. 1948;1(2):93-122. doi: 10.1152/jappl.1948.1.2.93. PMID: 18887578.

[8] Diller KR. *Heat transfer in health and healing*. J Heat Transfer. 2015;137(9):090801. doi: 10.1115/1.4030424.

[9] Knight, K. L. *Cryotherapy in Sport Injury Management*. Human Kinetics, 1995. ISBN: 9780873227711

[10] Jutte, L. S., Merrick, M. A., Ingersoll, C. D., and Edwards, J. E. "The Relationship Between Intramuscular Temperature, Skin Temperature, and Adipose Thickness During Cryotherapy and Rewarming," *Archives of Physical Medicine and Rehabilitation*, Vol. 82, No. 6, pp. 845–850, 2001. doi: 10.1053/apmr.2001.23195

[11] Gage, A. A., and Baust, J. "Mechanisms of Tissue Injury in Cryosurgery," *Cryobiology*, Vol. 37, No. 3, pp. 171–186, 1998. doi: 10.1006/cryo.1998.2115

[12] Rubinsky, B. "Cryosurgery," *Annual Review of Biomedical Engineering*, Vol. 2, pp. 157–187, 2000. doi: 10.1146/annurev.bioeng.2.1.157

[13] Banfi, G., Lombardi, G., Colombini, A., and Melegati, G. "Whole-Body Cryotherapy in Athletes," *Sports Medicine*, Vol. 40, No. 6, pp. 509–517, 2010. doi: 10.2165/11531940-000000000-00000

[14] Mayo Clinic Press. "The Chilling Truth: Exploring the Health Benefits and Risks of Cryotherapy," 2024.

[15] Bleakley, C. M., and Davison, G. W. "What is the Biochemical and Physiological Rationale for Using Cold-Water Immersion in Sports Recovery?" *British Journal of Sports Medicine*, Vol. 44, No. 3, pp. 179–187, 2010. doi: 10.1136/bjsm.2009.065565

[16] Wilcock, I. M., Cronin, J. B., and Hing, W. A. "Physiological Response to Water Immersion: A Method for Sport Recovery?" *Sports Medicine*, Vol. 36, No. 9, pp. 747–765, 2006. doi: 10.2165/00007256-200636090-00003

[17] Merrick, M. A., Jutte, L. S., & Smith, M. E. (2003). *Cold modalities with different thermodynamic properties produce different surface and intramuscular temperatures*. Journal of Athletic Training, 38(2), 149.

[18] Otte, J. W., et al. (2002). *Subcutaneous Adipose Tissue Thickness Alters the Effect of Ice Pack Application on Intramuscular Temperature*. Journal of Athletic Training, 37(2), 126–132.

[19] *Cooling Efficiency of 4 Common Cryotherapeutic Agents*. (2007). Journal of Athletic Training, 42(3), 343–348.

[20] *Mathematical modeling of the frozen zone dynamics: Towards using thermal imagers in cryotherapy*. (2026). PMC. Available at: https://pmc.ncbi.nlm.nih.gov/articles/PMC13086359/

[21] *Analysis and Simulation of Dynamic Heat Transfer and Thermal Distribution in Burns with Multilayer Models Using Finite Volumes*. (2025). MDPI. https://www.mdpi.com/2673-8716/5/4/41

[22] Rabin, Y., & Shitzer, A. (1998). *Numerical solution of the multidimensional freezing problem during cryosurgery*. Journal of Biomechanical Engineering, 120(1), 32-37.

[23] Kelly, J. S., & Bird, E. (2021). *Impact of Whole-Body Cryotherapy on Physiological and Cognitive Responses*. Frontiers in Physiology.

[24] Westerlund, T., et al. (2003). *Thermal responses during and after whole-body cryotherapy (-110 °C)*. Journal of Thermal Biology, 28(8), 601-608.

[25] He, X. (2011). *Thermostability of Biological Systems: Fundamentals, Challenges, and Quantification*. The Open Biomedical Engineering Journal, 5, 47–73.

[26] Kotova, T. G., et al. (2016). *Calculation of Effective Freezing Time in Lung Cancer Cryosurgery Based on Godunov Simulation*. Sovremennye tehnologii v medicine, 8(1), 48–54.

[27] Diller, K. R. (2015). *Heat Transfer in Health and Healing*. Journal of Heat Transfer, 137(1).

[28] Shafirstein, G., & Feng, Y. (2013). *The role of mathematical modelling in thermal medicine*. International Journal of Hyperthermia, 29(4), 259-261.

[29] Cooper, S. M. and Dawber, R. P. *The history of cryosurgery*, Journal of the Royal Society of Medicine, **94**(4), 196–201 (2001).

[30] Legrand, F. D., et al. *Evaluating safety risks of whole-body cryotherapy/cryostimulation (WBC): a scoping review from an international consortium*, European Journal of Medical Research, **28**(1), 387 (2023). doi: 10.1186/s40001-023-01385-z.

[31] Wang, X., et al. *Development and performance test of a whole-body cryotherapy cabin cooled by mixed-refrigerant*, Applied Thermal Engineering, **263**, 125369 (2025). doi: 10.1016/j.applthermaleng.2024.125369.

[32] De Nardi, M. et al. *Thermal responses induced by nitrogen and forced convection based partial-body cryostimulation*, Journal of Thermal Biology, **115**, 103620 (2023). doi: 10.1016/j.jtherbio.2023.103620.

[33] Louis, J., et al. *The use of whole-body cryotherapy: time- and dose-response investigation on circulating blood catecholamines and heart rate variability*, European Journal of Applied Physiology, **120**(8), 1733–1743 (2020). doi: 10.1007/s00421-020-04406-5.

[34] Ikäheimo, T., et al. *Autonomic nervous function during whole-body cold exposure before and after cold acclimation*, Aviation, Space, and Environmental Medicine, **79**, 875–882 (2008). doi: 10.3357/ASEM.2235.2008.

[35] N. Vaquero-Gallardo, O. Millán-Blasco, and H. Martínez-García, *Real-time detection system for preventing tissue damage in cryotherapy through temperature and bioimpedance monitoring*, Measurement, **254**, 117898 (2025). doi: 10.1016/j.measurement.2025.117898.

[36] Yao, Y., et al. *Cryotherapy and thermotherapy in the management of osteoarthritis and rheumatoid arthritis: A comprehensive review*, Fundamental Research, **5**(6), 2409–2431 (2025). doi: 10.1016/j.fmre.2024.07.008.

[37] Mejia, N., et al. *An On-Site Thermoelectric Cooling Device for Cryotherapy and Control of Skin Blood Flow*, Journal of Medical Devices, **9**(4), 0445021–0445026 (2015). doi: 10.1115/1.4029508.

[38] Kwak, K., et al. *Recent progress in cryoablation cancer therapy and nanoparticles mediated cryoablation*, Theranostics, **12**(5), 2175–2204 (2022). doi: 10.7150/thno.67530.