


# First Principles of Math



Sindre Sogge Heggen

*"Wahrlich es ist nicht das Wissen, sondern das Lernen,  
nicht das Besitzen, sondern das Erwerben,  
nicht das Da-Seyn, sondern das Hinkommen,  
was den grössten Genuss gewährt"*

*"It is not knowing, rather learning,  
not possessing, rather obtaining,  
not being present, rather reaching there,  
which serves the greatest joy."*

— Carl Friedrich Gauss

This document is created by Sindre Sogge Heggen. The text is written in L<sup>A</sup>T<sub>E</sub>X and the figures are made using Asymptote.

This book is part of the OpenMathBooks project. OpenMathBooks © 2022 by Sindre Sogge Heggen is licensed under CC BY-NC-SA 4.0. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>

15.02.2023

# Foreword

The extent and applications of mathematics are enormous, but a fair share of it is founded on a manageable amount of principles: I wish to present these in this book. I have chosen to call a principle in summarized form a *rule*. You will find the rules in blue text boxes, usually followed by an example of its usage. One of the main targets of this book is presenting the logical justification for the rules. In Chapter 1 - 5 you will find explanations<sup>1</sup> preceding every rule, while in chapter 6 some explanations are found directly after stating the rules (and eventual examples). As of chapter 7, some explanations are found in a concluding section named *Explanations*. This indicates that they are rather intricate or are so intuitively true that many will find the explanation superfluous.

## The structure of the book

The book consists of a *Part I* and a *Part II*. *Part I* focuses on the basic understanding of the numbers and operations of calculation. *Part II* introduces the concept of algebra and the closely related topics of powers, equations, and functions. In addition, both *Part I* and *Part II* end with a chapter on geometry.

*Notice! You will not find practice problems and applications of mathematics in real life in this book. These are two very important elements to come, either integrated in this book or as an independent document.*

## A note on convention

Although I am very much aware of the convention of writing commas and dots in center-aligned equations, I opted against this<sup>2</sup>. In this way, a center-aligned equation is a grammatical hybrid; it can end with both an invisible comma or dot, or nothing at all.

---

<sup>1</sup>To *explain* the rules rather than *proving* them is a deliberate decision. A proof demands mathematical rigor that often forces a lot of assumptions and definitions along the way. This can make the main insight disappear in the crowd of details. However, some of the explanations are valid as proofs.

<sup>2</sup>I've never liked the looks of it.

**I want to thank:**

Anne Jordal Myrset

Charlotte Merete Dahl

Brenden Callahan

For brilliant suggestions and comments.

## Symbols

$=$	"equals"
$<$	"less than"
$>$	"greater than"
$\leq$	"less than or equal to"
$\geq$	"greater than or equal to"
$\in$	"included in"
$\vee$	"or"
$\wedge$	"and"
$[a, b]$	"closed interval from $a$ to $b$ "
$ a $	"length/absolute value of $a$ "
$\perp$	"perpendicular to"
$\parallel$	"parallel with"
$\triangle$	"triangle"
$\square$	"quadrilateral"

# Contents

<b>I</b>	<b>Numbers, calculations, and geometry</b>	<b>8</b>
<hr/>		
<b>1</b>	<b>The numbers</b>	<b>9</b>
1.1	The equal sign, amounts, and number lines . . . . .	10
1.2	Numbers, digits and value . . . . .	12
1.3	Coordinate systems . . . . .	16
	Exercises . . . . .	17
<b>2</b>	<b>The four elementary operations</b>	<b>20</b>
2.1	Addition . . . . .	21
2.2	Subtraction . . . . .	23
2.3	Multiplication . . . . .	25
2.4	Division . . . . .	28
	Exercises . . . . .	31
<b>3</b>	<b>Factorization and order of operations</b>	<b>35</b>
3.1	Order of operations . . . . .	36
3.2	Factorization . . . . .	42
	Exercises . . . . .	44
<b>4</b>	<b>Fractions</b>	<b>47</b>
4.1	Introduction . . . . .	48
4.2	Values, expanding and simplifying . . . . .	51
4.3	Addition and subtraction . . . . .	54
4.4	Fractions multiplied by integers . . . . .	58
4.5	Fractions divided by integers . . . . .	60
4.6	Fractions multiplied by fractions . . . . .	63
4.7	Cancellation of fractions . . . . .	64
4.8	Division by fractions . . . . .	68
4.9	Rational numbers . . . . .	71
	Exercises . . . . .	73
<b>5</b>	<b>Negative numbers</b>	<b>81</b>
5.1	Introduction . . . . .	82
5.2	The elementary operations . . . . .	84
5.3	Negative numbers as amounts . . . . .	90
	Exercises . . . . .	91
<b>6</b>	<b>Calculation methods</b>	<b>93</b>
6.1	Addition . . . . .	94
6.2	Subtraction . . . . .	96
6.3	Multiplication . . . . .	99
6.4	Division . . . . .	104
6.4.1	Division using decimal numbers . . . . .	112
6.5	Calculation with time . . . . .	113
6.6	Rounding and estimates . . . . .	114
6.6.1	Estimate . . . . .	115

6.7	Standard form . . . . .	118
	Exercises . . . . .	121
<b>7</b>	<b>Geometry</b>	<b>124</b>
7.1	Terms . . . . .	125
7.2	Attributes of triangles and quadrilaterals . . . . .	134
7.3	Perimeter . . . . .	138
7.4	Area . . . . .	139
7.5	Tredimensjonal geometri . . . . .	146
7.6	Volum . . . . .	149
	Exercises . . . . .	151

## II Algebra and geometry 157

---

<b>8</b>	<b>Algebra</b>	<b>158</b>
8.1	Introduction . . . . .	159
8.2	Powers . . . . .	164
8.3	Irrational numbers . . . . .	172
	Exercises . . . . .	173
<b>9</b>	<b>Equations</b>	<b>179</b>
9.1	Introduction . . . . .	180
9.2	Solving with the elementary operations . . . . .	181
9.3	Solving with elementary operations summarized . . . . .	188
9.4	Power equations . . . . .	191
	Exercises . . . . .	193
<b>10</b>	<b>Functions</b>	<b>196</b>
10.1	Introduction . . . . .	197
10.2	Linear functions and graphs . . . . .	200
	Exercises . . . . .	208
<b>11</b>	<b>Geometry</b>	<b>214</b>
11.1	Formulas of area and perimeter . . . . .	215
11.2	Congruent and similar triangles . . . . .	225
11.3	Explanations . . . . .	230
	Exercises . . . . .	242
	<b>Exercise answers</b>	<b>252</b>
	<b>Index</b>	<b>260</b>




## Part I

# Numbers, calculations, and geometry

# Chapter 1

## The numbers



Bilde fra [pixabay.com](https://pixabay.com)

## 1.1 The equal sign, amounts, and number lines

### The equal sign

As the name implies, the *equal sign*  $=$  refers to things that are the same. In what sense some things are the same is a philosophical question and initially we are bound to this: What equality  $=$  points to must be understood by the context in which the sign is used. With this understanding of  $=$  we can study some basic properties of our numbers and then later return to more precise meanings of the sign.

#### The language box

Common ways of expressing  $=$  is

- "equals"
- "is the same as"

### Amounts and number lines

There are many ways a number can be defined, however, in this book we shall stick to two ways of interpreting a number; a number as an *amount* and a number as a *placement on a line*. All representations of numbers rely on the understanding of 0 and 1.

#### Numbers as amounts

Talking about an amount, the number 0 is<sup>1</sup> connected to "nothing". A figure showing nothing will therefore equal 0:

$$= 0$$

1 we'll draw like a box:

$$\square = 1$$

In this way, other numbers are defined by how many one-boxes (ones/units) we have:

$$\begin{array}{c} \square \\ \square \end{array} = 2$$

$$\begin{array}{c} \square \\ \square \\ \square \end{array} = 3$$

$$\begin{array}{c} \square \\ \square \\ \square \\ \square \end{array} = 4$$

---


<sup>1</sup>In [Chapter 6](#) we'll see that there are other interpretations of 0.

## Numbers as placements on a line


When placing numbers on a line, 0 is our starting point:



Now we place 1 a set length to the right of 0:



Other numbers are now defined by how many one-lengths (ones/units) we are away from 0:



## Positive integers

We'll soon see that numbers do not necessarily have to be a *whole* amount of ones, but those which *are* have their own name:

### 1.1 Positive integers

Numbers which are a whole amount of ones are called *positive<sup>1</sup> integers*. The positive integers are

1, 2, 3, 4, 5 and so on.

Positive integers are also called *natural numbers*.

### What about 0?

Some authors also include 0 in the definition of natural numbers. This is in some cases beneficial, in others not.

---


<sup>1</sup>We'll see what the word *positive* refers to in chapter [chapter 5](#).

# 1.2 Numbers, digits and value

Our numbers consist of the *digits* 0, 1, 2, 3, 4, 5, 6, 7, 8 and 9 along with their *positions*. The digits and their positions defines<sup>1</sup> the *value* of numbers.


## Integers larger then 10

Let's, as an example, write the number *fourteen* by our digits.



We can now make a group of 10 ones, then we also have 4 ones. By this, we write fourteen as


fourteen = 14




<sup>1</sup>Later on, we'll also see that *signs* have an impact on a numbers value (see [Chapter 5](#)).

# Decimal numbers


Sometimes we don't have a whole amount of ones, and this brings about the need to divide "ones" into smaller pieces. Let's start off by drawing a one:




Now we divide our one into 10 smaller pieces:




Since we have divided 1 into 10 pieces, we name one such piece *a tenth*:



We indicate tenths by using the *decimal mark*: .



## Example



## The language box

In a lot of countries, a comma is used in place of the period for the decimal mark.

3,5 (*other*)

3.5 (*English*)

# Base-10 positional notation


So far, we have seen how we can express the value of a number by placing digits according to the amount of tens, ones and tenths. The pattern continues:

## 1.2 Base-10 positional notation

The value of a number is given by the digits 0, 1, 2, 3, 4, 5, 6, 7, 8 and 9 and their position. In respect to the digit indicating ones,

- digits to the left indicate amounts of tens, hundreds, thousands etc.
- digits to the left indicate amounts of tenths, hundredths, thousandths etc.

### Example 1



### Example 2

thousands  
hundreds  
tens  
ones  
tenths  
hundredths  
3805.72

### 1.3 Even and odd numbers

Integers with 0, 2, 4 or 6 on the ones place are called *even numbers* .

Integers with 1, 3, 5, 7 or 9 on the ones place are called *odd numbers* .

#### Example

The first ten (positive) even numbers are

0, 2, 4, 6, 8, 10, 12, 14, 16, og 18

The first ten (positive) odd numbers are

1, 3, 5, 7, 9, 11, 13, 15, 17, og 19



## 1.3 Coordinate systems


Two number lines can be put together to form a *coordinate system*. In that case we place one number line *horizontally* and one *vertically*. A position in a coordinate system is called a *point*.

In fact, there are many types of coordinate systems, but we'll use the *cartesian coordinate system*. It is named after the French mathematician and philosopher, René Descartes.

A point is written as two numbers inside a bracket. We shall call these two numbers the *first coordinate* and the *second coordinate*.

- The first coordinate tells how many units to move along the horizontal axis.
- The second coordinate tells how many units to move along the vertical axis.




In the figure, the points  $(2, 3)$ ,  $(5, 1)$  and  $(0, 0)$  are shown. The point where the axes intersect,  $(0, 0)$ , is called *origo*.



# Exercises for Chapter 1




## 1.1.1

Write the value of the number.





## 1.1.2

Write the value of the number marked red.



## 1.1.3


Write the value of the number marked red.



1.1.4




Write the value of the number.




1.1.5

Choose which one of a), b) and c) have the value 0.1.




1.1.6




Choose which one of a), b) and c) have the value 0.1.



1.1.7


Choose which one of a), b) and c) have the value 0.1.






- a)  b)  c) 

1.1.8


Choose which one of a), b) and c) have the value 0.01.






- a)  b)  c) 

1.1.9

Choose which one of a), b) and c) have the value 0.01.



- a)  b)  c) 

## Chapter 2

# The four elementary operations




Bilde fra [pxhere.com](https://pxhere.com)

## 2.1 Addition

### Addition with amounts


When we have an amount and wish to add more, we use the symbol  $+$ . If we have 2 and want to add 3, we write

$$2 + 3 = 5$$



The order in which we add have no impact on the results; starting off with 2 and adding 3 is the same as starting off with 3 and adding 2:

$$3 + 2 = 5$$



#### The language box

A calculation involving addition includes two or more *terms* and one *sum*. In the calculation

$$2 + 3 = 5$$

both 2 and 3 are terms while 5 is the sum.

Common ways of saying  $2 + 3$  include

- "2 plus 3"
- "2 added to 3"
- "2 and 3 added"

## 2.1 Addition is commutative

The order of the terms has no impact on the sum.

### Example

$$2 + 5 = 7 = 5 + 2$$


$$6 + 3 = 9 = 3 + 6$$

## Addition on the number line: moving to the right

On a number line, addition with positive numbers involves moving *to the right*:


### Example 1

$$2 + 7 = 9$$



### Example 2

$$4 + 11 = 15$$




### Interpretation of =

+ brings the possibility of expressing numbers in different ways, for example is  $5 = 2 + 3$  and  $5 = 1 + 4$ . In this context, = means "has the same value as". This is also the case regarding subtraction, multiplication and division which we'll look at in the next three sections.

# 2.2 Subtraction

## Subtraction with amounts

When removing a part of an amount, we use the symbol :

$$5 - 3 = 2$$

  $-$    $=$  

### The language box

A calculation involving subtraction includes one or more *terms* and one *difference*. In the calculation

$$5 - 3 = 2$$

both 5 and 3 are terms while 2 is the difference.

Common ways of saying  $5 - 3$  include

- "5 minus 3"
- "3 subtracted from 5"

### A new interpretation of 0

As mentioned earlier in this book, 0 can be interpreted as "nothing". However, subtraction brings the possibility of expressing 0 by other numbers, for example  $7 - 7 = 0$  and  $19 - 19 = 0$ . In many practical situations, 0 indicates some form of equilibrium, like two equal but opposite forces.




## Subtraction on the number line: Moving to the left

In [Section 2.1](#), we have seen that  $+$  (with positive numbers) involves moving *to the right* on the number line. With  $-$  it's the opposite, we move *to the left*<sup>1</sup>:

### Example 1

$$6 - 4 = 2$$



### Example 2

$$12 - 7 = 5$$



### Notice

At first it may seem a bit odd moving in the opposite direction of the way in which the arrows point, as in *Example 1* and *2*. However, in [Chapter 5](#) this will turn out to be useful.

---

<sup>1</sup>In figures with number lines, the red colored arrows indicates that you shall start at the arrowhead and move to the other end.

# 2.3 Multiplication

## Multiplication by integers: initial definition

When adding equal numbers, we can use the multiplication symbol  $\cdot$  to write our calculations more compact:

### Example

$$4 + 4 + 4 = 4 \cdot 3$$

$$8 + 8 = 8 \cdot 2$$

$$1 + 1 + 1 + 1 + 1 = 1 \cdot 5$$

### The language box

A calculation involving multiplication includes several *factors* and one *product*. In the calculation

$$4 \cdot 3 = 12$$

both 4 and 3 are factors, while 12 is the product.

Common ways of saying  $4 \cdot 3$  include

- "4 times 3"
- "4 multiplied by 3"
- "4 and 3 multiplied together"

A lot of texts use  $\times$  instead of  $\cdot$ . In computer programming,  $*$  is the most common symbol for multiplication.

## Multiplication involving amounts

Let us illustrate  $2 \cdot 3$ :

$$2 \cdot 3 = \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} + \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} + \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}$$

Now notice the product of  $3 \cdot 2$ :

$$3 \cdot 2 = \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} + \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}$$

## 2.2 Multiplication is commutative

The order of the factors has no impact on the product.

### Example

$$3 \cdot 4 = 12 = 4 \cdot 3$$

$$6 \cdot 7 = 42 = 7 \cdot 6$$


$$8 \cdot 9 = 72 = 9 \cdot 8$$

### Multiplication on the number line

We can also use the number line to calculate multiplications. In the case of  $2 \cdot 4$  we can think like this:

" $2 \cdot 4$  means moving 2 places to the right, 4 times."


$$2 \cdot 4 = 8$$



We can also use the number line to prove to ourselves that multiplication is commutative:

" $4 \cdot 2$  means moving 4 places to the right, 2 times."

$$4 \cdot 2 = 8$$



### Final definition of multiplication by positive integers

It may be the most intuitive to interpret " $2$  times  $3$ " as " $3$ ,  $2$  times". Then it follows:

$$"/>2 \text{ times } 3" = 3 + 3$$

In this section we introduced  $2 \cdot 3$ , that is " $2$  times  $3$ ", as  $2 + 2 + 2$ . With this interpretation,  $3 + 3$  corresponds to  $3 \cdot 2$ , but the fact that multiplication is a commutative operation ([Rule 2.2](#)) ensures that the one interpretation does not exclude the other;  $2 \cdot 3 = 2 + 2 + 2$  and  $2 \cdot 3 = 3 + 3$  are two expressions of same value.

## 2.3 Multiplication as repeated addition

Multiplication involving a positive integer can be expressed as repeated addition.

### Example 1

$$4 + 4 + 4 = 4 \cdot 3 = 3 + 3 + 3 + 3$$

$$8 + 8 = 8 \cdot 2 = 2 + 2 + 2 + 2 + 2 + 2 + 2$$

$$1 + 1 + 1 + 1 + 1 = 1 \cdot 5 = 5$$

### Notice

The fact that multiplication with positive integers can be expressed as repeated addition does not exclude other expressions. There's nothing wrong with writing  $2 \cdot 3 = 1 + 5$ .

## 2.4 Division

$:$  is the symbol for division. Division has three different interpretations:

### 2.4 The three interpretations of division

- **Distribution of amounts**

$12 : 3 =$  "The number in each group when evenly distributing 12 into 3 groups"

- **Number of equal terms**

$12 : 3 =$  "The number of 3's added to make 12"

- **The inverse operation of multiplication**

$12 : 3 =$  "The number which yields 12 when multiplied by 3"

### The language box

A calculation involving division includes a *dividend*, a *divisor* and a *quotient*. In the calculation

$$12 : 3 = 4$$

12 is the dividend, 3 is the divisor and 4 is the quotient.

Common ways of saying  $12 : 3$  include


- "12 divided by 3"
- "12 to 3"

In a lot of contexts,  $/$  is used instead of  $:$ , especially in computer programming.

Sometimes  $12 : 3$  is called "the *ratio* of 12 to 3".

### Distribution of amounts


The calculation  $12 : 3$  tells that we shall distribute 12 into 3 equal groups:



We observe that each group contains 4 boxes, which means that

$$12 : 3 = 4$$


### Number of equal terms




12 equals the sum of 4 instances of 3, that is  $12 : 3 = 4$ .

### The inverse operation of multiplication

We have just seen that if we divide 12 into 3 equal groups, we get 4 in each group. Hence  $12 : 3 = 4$ . The sum of these groups makes 12:



However, this is the same as multiplying 4 by 3, in other words: If we know that  $4 \cdot 3 = 12$ , we also know that  $12 : 3 = 4$ . As well we know that  $12 : 4 = 3$ .



**Example 1**

Since  $6 \cdot 3 = 18$ ,

$$18 : 6 = 3$$

$$18 : 3 = 6$$

**Example 2**

Since  $5 \cdot 7 = 35$ ,

$$35 : 5 = 7$$

$$35 : 7 = 5$$

## Exercises for Chapter 2

### 2.1.1

Write the numbers as the sum of two numbers.

**Example**

We can write 3 as  $1 + 2$

- a) 4      b) 5      c) 6      d) 7      e) 8      f) 9

### 2.1.2

Write the numbers as the sum of three numbers.

**Example**

We can write 4 as  $1 + 2 + 1$

- a) 5      b) 6      c) 7      d) 8      e) 9      f) 10

### 2.1.3

When the sum of two numbers equals 10, these numbers are called **friends of ten**. For example are 1 and 9 friends of ten because  $1 + 9 = 10$ .

1) Find the ten-friend of

- a) 2      b) 3      c) 4      d) 5

2) When exercise 1) is completed, why is it "unnecessary" to find the ten-friends of 6, 7 og 8?



### 2.1.4

*Note:* You can allow yourself answering the questions by simply checking a couple of examples. For proof, see [Exercise 11](#).

Choose the correct alternative of 1), 2) and 3).

- a) The sum of two even numbers is
  - 1) an even number.
  - 2) an odd number.
  - 3) sometimes an even number and sometimes an odd number.
- b) The sum of two odd numbers is
  - 1) an even number.
  - 2) an odd number.
  - 3) sometimes an even number and sometimes an odd number.
- c) The sum of an even number and an odd number is
  - 1) an even number.
  - 2) an odd number.
  - 3) sometimes an even number and sometimes an odd number.

### 2.2.1

Write the number as the difference of two numbers.

**Example**

We can write 1 as  $8 - 7$ .

- a) 2      b) 3      c) 4      d) 5      e) 6      f) 7      g) 8

### 2.2.2

*Note:* You can allow yourself answering the questions by simply checking a couple of examples. For proof, see [Exercise 11](#).

Choose the correct alternative of 1), 2) and 3).

- a) The difference of two even numbers is
- 1) an even number.
  - 2) an odd number.
  - 3) sometimes an even number and sometimes an odd number.
- b) The difference of two odd numbers is
- 1) an even number.
  - 2) an odd number.
  - 3) sometimes an even number and sometimes an odd number.
- c) The difference of an even number and an odd number is
- 1) an even number.
  - 2) an odd number.
  - 3) sometimes an even number and sometimes an odd number.

### 2.3.1

Write as a multiplication, and write the alternative sum.

**Example**



$$3 + 3 + 3 + 3 + 3 = 3 \cdot 5 = 5 + 5 + 5$$

- a)  $2 + 2 + 2$
- b)  $3 + 3 + 3 + 3 + 3 + 3$
- c)  $4 + 4$
- d)  $5 + 5 + 5 + 5 + 5 + 5 + 5 + 5 + 5 + 5$
- e)  $6 + 6 + 6 + 6$
- f)  $7 + 7 + 7 + 7$

### 2.3.2

Draw boxes to find the answer to the multiplications.

**Example**



- a)  $4 \cdot 5$
- b)  $8 \cdot 3$
- c)  $2 \cdot 9$
- d)  $5 \cdot 6$
- e)  $7 \cdot 8$

### 2.3.3

- a) Will an integer multiplied by 2 always result in an even number or an odd number?
- b) Will an even number multiplied by 5 always result in an even number or an odd number? What digit will always be positioned at the one-place?
- c) Will an odd number multiplied by 5 always result in an even number or an odd number? What digit will always be positioned at the one-place?

## Chapter 3

# Factorization and order of operations

## 3.1 Order of operations

### Priority of the operations

Look at the following calculation:

$$2 + 3 \cdot 4$$

This *could* have been interpreted in two ways:

1. "2 plus 3 equals 5. 5 times 4 equals 20. The answer is 20."
2. "3 times 4 equals 12. 2 plus 12 equals 14. The answer is."

But the answers are not the same! This points out the need to have rules for what to calculate first. One of these rules is that multiplication and division is to be calculated *before* addition or subtraction, which means that

$$\begin{aligned} 2 + 3 \cdot 4 &= \text{"Calculate } 3 \cdot 4, \text{ then add } 2\text{"} \\ &= 2 + 12 \\ &= 14 \end{aligned}$$

But what if we wanted to calculate  $2 + 3$  first, then multiply the sum by 4? We use parentheses to tell that something is to be calculated first:

$$\begin{aligned} (2 + 3) \cdot 4 &= \text{"Calculate } 2 + 3, \text{ multiply by } 4 \text{ afterwards"} \\ &= 5 \cdot 4 \\ &= 20 \end{aligned}$$

### 3.1 Order of operations

1. Expressions with parentheses
2. Multiplication or division
3. Addition or subtraction

### Example 1

Calculate

$$23 - (3 + 9) + 4 \cdot 7$$

**Answer**

$$\begin{aligned} 23 - (3 + 9) + 4 \cdot 7 &= 23 - 12 + 4 \cdot 7 && \text{Parentheses} \\ &= 23 - 12 + 28 && \text{Multiplication} \\ &= 39 && \text{Addition and subtraction} \end{aligned}$$

### Example 2

Calculate

$$18 : (7 - 5) - 3$$

**Answer**

$$\begin{aligned} 18 : (7 - 5) - 3 &= 18 : 2 - 3 && \text{Parentheses} \\ &= 9 - 3 && \text{Division} \\ &= 6 && \text{Addition and subtraction} \end{aligned}$$

## Multiplication involving parentheses

How many boxes are present in this figure?



Two correct interpretations include:

1. It is  $2 \cdot 4 = 8$  purple boxes and  $3 \cdot 4 = 12$  green boxes. In total there are  $8 + 12 = 20$  boxes. This we can write as

$$2 \cdot 4 + 3 \cdot 4 = 20$$

2. It is  $2 + 3 = 5$  boxes horizontally and 4 boxes vertically, so there are  $5 \cdot 4 = 20$  boxes in total. This we can write as

$$(2 + 3) \cdot 4 = 20$$

From these two calculations it follows that

$$(2 + 3) \cdot 4 = 2 \cdot 4 + 3 \cdot 4$$

### 3.2 Distributive law

When an expression enclosed by parentheses is a factor, we can multiply the other factors with each term inside the parentheses.

#### Example 1

$$(4 + 7) \cdot 8 = 4 \cdot 8 + 7 \cdot 8$$

#### Example 2

$$\begin{aligned}(10 - 7) \cdot 2 &= 10 \cdot 2 - 7 \cdot 2 \\ &= 20 - 14 \\ &= 6\end{aligned}$$

*Note:* Obviously, it would be easier to calculate like this:

$$(10 - 7) \cdot 2 = 3 \cdot 2 = 6$$

#### Example 2

Calculate  $12 \cdot 3$ .

**Answer**

$$\begin{aligned}12 \cdot 3 &= (10 + 2) \cdot 3 \\ &= 10 \cdot 3 + 2 \cdot 3 \\ &= 30 + 6 \\ &= 36\end{aligned}$$

#### Note

We introduced parentheses as an indicator of what to calculate first, but [Rule 3.2](#) gives an alternative and equivalent interpretation of parentheses. The rule is especially useful when working with algebra (see [Part II](#)).

### Multiplying by 0

Earlier we have seen that 0 can be expressed as the difference between two numbers, and this can help us calculate when multiplying by 0. Let's look at the calculation

$$(2 - 2) \cdot 3$$

By [Rule 3.2](#), we get

$$\begin{aligned}(2 - 2) \cdot 3 &= 2 \cdot 3 - 2 \cdot 3 \\ &= 6 - 6 \\ &= 0\end{aligned}$$

Since  $0 = 2 - 2$ , this means that

$$0 \cdot 3 = 0$$

### 3.3 Multiplication by 0

If 0 is a factor, the product equals 0.

#### Example 1

$$7 \cdot 0 = 0$$

$$0 \cdot 219 = 0$$

### Associative laws

### 3.4 Associative law for addition

The placement of parentheses between terms has no impact on the sum.

#### Example

$$(2 + 3) + 4 = 5 + 4 = 9$$

$$2 + (3 + 4) = 2 + 7 = 9$$

$$\boxed{\phantom{00}}\boxed{\phantom{00}} + \boxed{\phantom{00}}\boxed{\phantom{00}}\boxed{\phantom{00}} + \boxed{\phantom{00}} = \boxed{\phantom{00000000}}$$




### 3.5 Associative law for multiplication

The placement of parentheses between factors has no impact on the product.

#### Example

$$(2 \cdot 3) \cdot 4 = 6 \cdot 4 = 24$$

$$2 \cdot (3 \cdot 4) = 2 \cdot 12 = 24$$



Opposite to addition and multiplication, neither subtraction nor division is associative:

$$(12 - 5) - 4 = 7 - 4 = 3$$

$$12 - (5 - 4) = 12 - 1 = 11$$

$$(80 : 10) : 2 = 8 : 2 = 4$$

$$80 : (10 : 2) = 80 : 5 = 16$$

We have seen how parentheses helps indicating the *priority* of operations, but the fact that subtraction and division are non-associative brings the need of having a rule of in which *direction* to calculate.

### 3.6 Direction of calculations

Operations which by [Rule 3.1](#) have equal priority, are to be calculated from left to right.

#### Example 1

$$\begin{aligned} 12 - 5 - 4 &= (12 - 5) - 4 \\ &= 7 - 4 \\ &= 3 \end{aligned}$$

**Example 2**

$$\begin{aligned}80 : 10 : 2 &= (80 : 10) : 2 \\&= 8 : 2 \\&= 4\end{aligned}$$

**Example 3**

$$\begin{aligned}6 : 3 \cdot 4 &= (6 : 3) \cdot 4 \\&= 2 \cdot 4 \\&= 8\end{aligned}$$

## 3.2 Factorization

If an integer dividend and an integer divisor results in an integer quotient, we say that the dividend is *divisible* by the divisor. For example is 6 divisible with 3 because  $6 : 3 = 2$ , and 40 is divisible with 10 because  $40 : 10 = 4$ . The concept of divisibility contributes to the definition of *prime numbers*:

### 3.7 Primal

A natural number larger than 1, and only divisible by itself and 1, is a prime number.

#### Example

The first five prime numbers are 2, 3, 5, 7 and 11.

### 3.8 Factorization

Factorization involves writing a number as the product of other numbers.

#### Example

Factorize 24 in three different ways.

#### Answer

$$24 = 2 \cdot 12$$

$$24 = 3 \cdot 8$$

$$24 = 2 \cdot 3 \cdot 4$$

#### The language box

Since 12 is divisible by 4, we say that 4 is a factor of 12.

### 3.9 Prime factorization

Factorization involving prime factors only is called prime factorization.

#### Example

Prime factorize 12.

**Answer**

$$12 = 2 \cdot 2 \cdot 3$$

#### The prime numbers from 1-100

91	92	93	94	95	96	97	98	99	100
81	82	83	84	85	86	87	88	89	90
71	72	73	74	75	76	77	78	79	80
61	62	63	64	65	66	67	68	69	70
51	52	53	54	55	56	57	58	59	60
41	42	43	44	45	46	47	48	49	50
31	32	33	34	35	36	37	38	39	40
21	22	23	24	25	26	27	28	29	30
11	12	13	14	15	16	17	18	19	20
1	2	3	4	5	6	7	8	9	10

## Exercises for Chapter 3

### 3.1.1

Calculate by writing the numbers as sums of ones, tens and hundreds, and applying the distributive law.

#### Example

$$15 \cdot 3 = (10 + 5) \cdot 3 = 10 \cdot 3 + 5 \cdot 3 = 30 + 15 = 45$$

$$147 \cdot 2 = (100 + 40 + 7) \cdot 2 = 100 \cdot 2 + 40 \cdot 2 + 7 \cdot 2 = 200 + 80 + 14 = 294$$

- a)  $17 \cdot 2$       b)  $59 \cdot 3$       c)  $25 \cdot 4$       d)  $582 \cdot 2$       e)  $981 \cdot 3$

### 3.2.1

Write the numbers as a multiplication between two factors.

- a) 100      b) 30      c) 40      d) 70  
e) 42      f) 32      g) 84      h) 90

### 3.2.2

Prime factorize the numbers from [Exercise 3.2.1](#).

*Note:* It is recommended that the reader find it's own ways of prime factorization, but those who insist on following a schematic method can look up [Exercise 3.2.5](#).

### 3.2.3

Factorize the numbers from [Exercise 3.2.1](#) in three different ways.

### 3.2.4

6 is called a **perfect number** because the sum of it's factors 6 (including 1, but excluding 6) equals 6:  $1 + 2 + 3 = 6$ . Find the next perfect number (it lies between 15 og 30).

3.2.5

The examples below shows a method for prime factorization of numbers. Explain the method.

Example 1

$84 = 2 \cdot 2 \cdot 3 \cdot 7$

	:	
84	2	42
42	2	21
21	3	7

Example 2

$595 = 5 \cdot 7 \cdot 17$

	:	
595	5	119
119	7	17

### **Gruble 1**

Explain why the product of two odd numbers always equals an odd number.

### **Gruble 2**

Explain a method for checking if a number is a prime number.

## Chapter 4

# Fractions



## 4.1 Introduction

### 4.1 Fractions as rewriting of division

A fraction is a different way of writing a division. In a fraction the dividend is called the *numerator* and the divisor the denominator.

$$1 : 4 = \frac{1}{4} \begin{array}{l} \longleftarrow \text{Numerator} \\ \longleftarrow \text{Denominator} \end{array}$$

### The language box

Common ways of saying  $\frac{1}{4}$  are<sup>1</sup>


- "one fourth"
- "1 of 4"
- "1 over 4"

---

<sup>1</sup>We also have the expressions from the language box on page 28.

### Fractions as amounts


Let us present  $\frac{1}{4}$  as an amount. We then think of the number 1 as a box<sup>1</sup>:


$$= 1$$


---

<sup>1</sup>For practical reasons, we choose a unit box larger than the one used in [Chapter 1](#).


Further, we divide this box into four smaller, equal-sized boxes. The sum of these boxes equals 1.



One such box equals  $\frac{1}{4}$ :



However, if you from a figure only are to see how large a fraction is, the size of 1 must be known, and to make this more apparent we'll also include the "empty" boxes:



In this way, the blue and the empty boxes tell us how many pieces 1 is divided into, while the blue boxes alone tells how many of these boxes are *actually* present. In other words,

number of blue boxes = numerator

number of blue boxes + number of empty boxes = denominator


$= \frac{2}{3}$

$= \frac{7}{10}$


$= \frac{19}{20}$

## Fractions on the number line

On the number line, we divide the length between 0 and 1 into as many pieces as the denominator indicates. In the case of a fraction with denominator 4, we divide the length between 0 and 1 into 4 equal lengths:



Moreover, fractions larger than 1 are easily presented on the number line:



## Numerator and denominator summarized

Although already mentioned, the interpretations of the numerator and the denominator are of such importance that we shortly summarize them:

- The denominator tells how many pieces 1 is divided into.
- The numerator tells how many of these pieces are present.

# 4.2 Values, expanding and simplifying

## 4.2 The value of a fraction

The value of a fraction is given by dividing the numerator by the denominator.

### Example

Find the value of  $\frac{1}{4}$ .



**Answer**

$$\frac{1}{4} = 0.25$$

## Fractions with equal value

Fractions can have the same value even though they look different. If you calculate  $1 : 2$ ,  $2 : 4$  and  $4 : 8$ , you will in every case end up with 0.5 as the answer. This means that


$$\frac{1}{2} = \frac{2}{4} = \frac{4}{8} = 0,5$$



# Expanding

The fact that fractions can look different but have the same value, implies that we can change a fraction's look without changing its value. Let's, as an example, change  $\frac{3}{5}$  into a fraction of equal value but with denominator 10:

- We can make  $\frac{3}{5}$  into a fraction with denominator 10 if we divide each fifth into 2 equal pieces. In that case, 1 is divided into  $5 \cdot 2 = 10$  pieces in total.
- The numerator of  $\frac{3}{5}$  indicates that there are 3 fifths. When these are divided by 2, they make up  $3 \cdot 2 = 6$  tenths. Hence  $\frac{3}{5}$  equals  $\frac{6}{10}$ .



# Simplifying

Notice that we can also go "the opposite way". We can change  $\frac{6}{10}$  into a fraction with denominator 5 by dividing both the numerator and the denominator by 2:

$$\frac{6}{10} = \frac{6 : 2}{10 : 2} = \frac{3}{5}$$

## 4.3 Expanding of fractions

We can either multiply or divide both the numerator and the denominator by the same number without alternating the fractions value.

Multiplying by a number larger than 1 is called *expanding* the fraction. Dividing by a number larger than 1 is called *simplifying* the fraction.

**Example 1**

Expand  $\frac{3}{5}$  into a fraction with denominator 20.

**Answer**

Since  $5 \cdot 4 = 20$ , we multiply both the numerator and the denominator by 4:

$$\begin{aligned}\frac{3}{5} &= \frac{3 \cdot 4}{5 \cdot 4} \\ &= \frac{12}{20}\end{aligned}$$

**Example 2**

Expand  $\frac{150}{50}$  into a fraction with denominator 100.

**Answer**

Since  $50 \cdot 2 = 100$ , we multiply both the numerator and the denominator by 2:

$$\begin{aligned}\frac{150}{50} &= \frac{150 \cdot 2}{50 \cdot 2} \\ &= \frac{300}{100}\end{aligned}$$

**Example 3**

Simplify  $\frac{18}{30}$  into a fraction with denominator 5.

**Answer**

Since  $30 : 6 = 5$ , we divide both the numerator and the denominator by 6:


$$\begin{aligned}\frac{18}{30} &= \frac{18 : 6}{30 : 6} \\ &= \frac{3}{5}\end{aligned}$$

# 4.3 Addition and subtraction

Addition and subtraction of fractions are in large parts focused around the denominators. Recall that the denominators indicate the partitioning of 1. If fractions have equal denominators, they represent amounts of equal-sized pieces. In this case it makes sense calculating addition or subtraction of the numerators. However, if fractions have unequal denominators, they represent amounts of different-sized pieces, and hence addition and subtraction of the numerators makes no sense directly.

## Equal denominators

If we, for example, have 2 sixths and add 3 sixths, the sum is 5 sixths:



## 4.4 Addition/subtraction of fractions with equal denominators

When adding/subtracting fractions with equal denominators, we find the sum/difference of the numerators and keep the denominator.

### Example 1

$$\begin{aligned}\frac{2}{7} + \frac{8}{7} &= \frac{2+8}{7} \\ &= \frac{10}{7}\end{aligned}$$

**Example 2**


$$\frac{7}{9} - \frac{5}{9} = \frac{7-5}{9}$$

$$= \frac{2}{9}$$

**Unequal denominators**

Let's examine the calculation<sup>1</sup>

$$\frac{3}{5} - \frac{1}{2}$$




To write the difference as a single fraction, the two terms need to have denominators of equal value. Both of the fractions can have denominator 10:

$$\frac{3}{5} = \frac{3 \cdot 2}{5 \cdot 2} = \frac{6}{10} \qquad \frac{1}{2} = \frac{1 \cdot 5}{2 \cdot 5} = \frac{5}{10}$$


Hence

$$\frac{3}{5} - \frac{1}{2} = \frac{6}{10} - \frac{5}{10}$$



<sup>1</sup>Recall that the red-colored arrow indicates that you shall start at the arrowhead and then move to the other end.





Summarized, we have expanded the fractions such that they have denominators of equal value, that is 10. When the denominators are equal, we can calculate the difference of the numerators:

$$\begin{aligned}\frac{3}{5} - \frac{1}{2} &= \frac{6}{10} - \frac{5}{10} \\ &= \frac{1}{10}\end{aligned}$$

#### 4.5 Addition/subtraction of fractions with unequal denominators

When calculating addition/subtraction of fractions, we must expand the fractions such that they have denominators of equal value, and then apply [Rule 4.4](#).

##### Example 1

Calculate

$$\frac{2}{9} + \frac{6}{7}$$

Both denominators can be transformed into 63 if multiplied by a fitting integer. Therefore, we expand the fractions as follows:

$$\begin{aligned}\frac{2 \cdot 7}{9 \cdot 7} + \frac{6 \cdot 9}{7 \cdot 9} &= \frac{14}{63} + \frac{54}{63} \\ &= \frac{68}{63}\end{aligned}$$

## Common denominator

In *Example 1* above, 63 is called a *common denominator* because there exists integers which, when multiplied by the original denominators, results in 63:

$$9 \cdot 7 = 63$$

$$7 \cdot 9 = 63$$

Multiplying together the original denominators always results in a common denominator but one can avoid large numbers by finding the *smallest* common denominator. Take, for example,

$$\frac{7}{6} + \frac{5}{3}$$

$6 \cdot 3 = 18$  is a common denominator, but it's worth noticing that  $6 \cdot 1 = 3 \cdot 2 = 6$  is too. Hence,

$$\begin{aligned}\frac{7}{6} + \frac{5}{3} &= \frac{7}{6} + \frac{5 \cdot 2}{3 \cdot 2} \\ &= \frac{7}{6} + \frac{10}{6} \\ &= \frac{17}{6}\end{aligned}$$

## Example 2

Calculate

$$\frac{3}{2} - \frac{5}{8} + \frac{10}{4}$$

### Answer


All denominators can be transformed into 8 if multiplied by a fitting integer. Therefore, we expand the fractions as follows:

$$\begin{aligned}\frac{3}{2} - \frac{5}{8} + \frac{10}{4} &= \frac{3 \cdot 4}{2 \cdot 4} - \frac{5}{8} + \frac{10 \cdot 2}{4 \cdot 2} \\ &= \frac{12}{8} - \frac{5}{8} + \frac{20}{8} \\ &= \frac{27}{8}\end{aligned}$$

## 4.4 Fractions multiplied by integers

In [Section 2.3](#) we observed that multiplying by an integer corresponds to repeated addition. Hence, if we are to calculate  $\frac{2}{5} \cdot 3$ , we can write

$$\begin{aligned}\frac{2}{5} \cdot 3 &= \frac{2}{5} + \frac{2}{5} + \frac{2}{5} \\ &= \frac{2+2+2}{5} \\ &= \frac{6}{5}\end{aligned}$$



Noticing that  $2 + 2 + 2 = 2 \cdot 3$ , we get

$$\begin{aligned}\frac{2}{5} \cdot 3 &= \frac{2 \cdot 3}{5} \\ &= \frac{6}{5}\end{aligned}$$

Multiplication of integers and fractions are also commutative<sup>1</sup>:

$$\begin{aligned}3 \cdot \frac{2}{5} &= 3 \cdot 2 : 5 \\ &= 6 : 5 \\ &= \frac{6}{5}\end{aligned}$$

### 4.6 Fractions multiplied by integers

When multiplying a fraction by an integer, we multiply the numerator by the integer.

---

<sup>1</sup>Recall that  $\frac{2}{5}$  corresponds to  $2 : 5$ .

**Example 1**

$$\begin{aligned}\frac{1}{3} \cdot 4 &= \frac{1 \cdot 4}{3} \\ &= \frac{4}{3}\end{aligned}$$

**Example 2**

$$\begin{aligned}3 \cdot \frac{2}{5} &= \frac{3 \cdot 2}{5} \\ &= \frac{6}{5}\end{aligned}$$

**An interpretation of multiplying by a fraction**

By [Rule 4.6](#) we can make an interpretation of multiplying by a fraction. For example, multiplying 3 by  $\frac{2}{5}$  can be interpreted in these two following ways:

- We multiply 3 by 2 and divide by 5:

$$(3 \cdot 2) : 5 = \frac{3 \cdot 2}{5} = \frac{6}{5}$$

- We divide 3 by 5 and multiply the quotient by 2:

$$3 : 5 = \frac{3}{5} \quad , \quad \frac{3}{5} \cdot 2 = \frac{3 \cdot 2}{5} = \frac{6}{5}$$

# 4.5 Fractions divided by integers



It is now important to recall two things:

- Division can be interpreted as an equal distribution of amounts
- In a fraction, it is the numerator which indicates the amount (the denominator indicates the partitioning of 1)



## When the numerator is divisible by the divisor

Let's calculate

$$\frac{6}{8} : 2$$



We have 6 eights which are to be equally distributed into 2 groups. This results in  $6 : 2 = 3$  eights.





Thus

$$\frac{6}{8} : 2 = \frac{3}{8}$$

**When the numerator is not divisible by the denominator**




What if we are to divide  $\frac{3}{4}$  by 2?

$$\frac{3}{4} : 2$$





Thing is, we can always expand the fraction such that the numerator becomes divisible by the divisor. Since 2 is the divisor, we expand the fraction by 2:

$$\frac{3}{4} = \frac{3 \cdot 2}{4 \cdot 2} = \frac{6}{8}$$



Now we have 6 eights. 6 eights divided by 2 equals 3 eights:



Hence

$$\frac{3}{4} : 2 = \frac{3}{8}$$

In effect, we have multiplied  $\frac{3}{4}$  by 2:

$$\begin{aligned}\frac{3}{4} : 2 &= \frac{3}{4 \cdot 2} \\ &= \frac{3}{8}\end{aligned}$$

#### 4.7 Fractions divided by integers

When dividing a fraction by an integer, we multiply the denominator by the integer.

##### Example 1

$$\begin{aligned}\frac{5}{3} : 6 &= \frac{5}{3 \cdot 6} \\ &= \frac{5}{18}\end{aligned}$$

##### Note

At the start of this section we found that

$$\frac{4}{8} : 2 = \frac{2}{8}$$

In that case, there were no need to multiply the denominator by 2, such as [Rule 4.7](#) implies. However, if we do, we have

$$\frac{4}{8} : 2 = \frac{4}{8 \cdot 2} = \frac{4}{16}$$

Now,

$$\frac{2}{8} = \frac{2 \cdot 2}{8 \cdot 2} = \frac{4}{16}$$

Hence, unsurprisingly, the two answers are of equal value.

## 4.6 Fractions multiplied by fractions

We have seen that<sup>1</sup> multiplying a number by a fraction involves multiplying the number by the numerator and then dividing the product by the denominator. Let us apply this to calculate

$$\frac{5}{4} \cdot \frac{3}{2}$$

Firstly, we multiply  $\frac{5}{4}$  by 3, then we divide the resulting product by 2. By [Rule 4.6](#), we have

$$\frac{5}{4} \cdot 3 = \frac{5 \cdot 3}{4}$$

And by [Rule 4.7](#), we get

$$\frac{5 \cdot 3}{4} : 2 = \frac{5 \cdot 3}{4 \cdot 2}$$

Hence

$$\frac{5}{4} \cdot \frac{3}{2} = \frac{5 \cdot 3}{4 \cdot 2}$$

### 4.8 Fractions multiplied by fractions

When multiplying a fraction by a fraction, we multiply numerator by numerator and denominator by denominator.

#### Example 1

$$\begin{aligned}\frac{4}{7} \cdot \frac{6}{9} &= \frac{4 \cdot 6}{7 \cdot 9} \\ &= \frac{24}{63}\end{aligned}$$

#### Example 2

$$\begin{aligned}\frac{1}{2} \cdot \frac{9}{10} &= \frac{1 \cdot 9}{2 \cdot 10} \\ &= \frac{9}{20}\end{aligned}$$

---

<sup>1</sup>Look at the text box titled "An interpretation of multiplying by a fraction" on page 59.



## 4.7 Cancellation of fractions

When the numerator and the denominator are of equal value, the fractions value always equals 1. For example,  $\frac{3}{3} = 1$ ,  $\frac{25}{25} = 1$  etc. We can exploit this fact to simplify expressions involving fractions.

Let us simplify the expression

$$\frac{8 \cdot 5}{9 \cdot 8}$$

Since  $8 \cdot 5 = 5 \cdot 8$ , we can write

$$\frac{8 \cdot 5}{9 \cdot 8} = \frac{5 \cdot 8}{9 \cdot 8}$$

And, as recently seen ([Rule 4.8](#)), we have

$$\frac{5 \cdot 8}{9 \cdot 8} = \frac{5}{9} \cdot \frac{8}{8}$$

Since  $\frac{8}{8} = 1$ ,

$$\begin{aligned} \frac{5}{9} \cdot \frac{8}{8} &= \frac{5}{9} \cdot 1 \\ &= \frac{5}{9} \end{aligned}$$

When multiplication is exclusively present in a fraction, you can always shuffle the way we did in the above expressions. However, when you have understood the outcome of the shuffling, it is better to apply *cancellation*. You then draw a line across two and two equal factors, thus indicating that they constitute a fraction which equals 1. Hence, our most recent example can be simplified to

$$\frac{\cancel{8} \cdot 5}{9 \cdot \cancel{8}} = \frac{5}{9}$$

## 4.9 Cancellation of factors

When multiplication is exclusively present in a fraction, we can cancel pair of equal factors in numerator and denominator.

### Example 1

Cancel as many factors as possible in the fraction.

$$\frac{3 \cdot 12 \cdot 7}{7 \cdot 4 \cdot 12}$$

**Answer**

$$\frac{3 \cdot \cancel{12} \cdot \cancel{7}}{\cancel{7} \cdot 4 \cdot \cancel{12}} = \frac{3}{4}$$

### Example 2

Simplify the fraction  $\frac{12}{42}$ .

**Answer**

We notice that 6 is a factor of both 12 and 42, and thus

$$\begin{aligned}\frac{12}{42} &= \frac{\cancel{6} \cdot 2}{\cancel{6} \cdot 7} \\ &= \frac{2}{7}\end{aligned}$$

### Example 3

Simplify the fraction  $\frac{48}{16}$ .

**Answer**

We notice that 16 is a factor of 48, and thus

$$\begin{aligned}\frac{48}{16} &= \frac{3 \cdot \cancel{16}}{\cancel{16}} \\ &= \frac{3}{1} \\ &= 3\end{aligned}$$

*Note:* If all factors are canceled in the numerator or the denominator, 1 takes their place.

### Simplifying by prime factorization

It's not necessarily easy to notice common factors, as we did in *Example 2* and *Example 3* above. To be sure you have not overlooked any common factors, you can always prime factorize (see page ??) both the numerator and the denominator. For example,

$$\begin{aligned}\frac{12}{42} &= \frac{2 \cdot 2 \cdot \cancel{3}}{2 \cdot \cancel{3} \cdot 7} \\ &= \frac{2}{7}\end{aligned}$$

### Fractions simplify calculations

The decimal number 0.125 can be written as the fraction  $\frac{1}{8}$ . The calculation

$$0.125 \cdot 16$$

is, for the most of us, rather strenuous to carry out. However, exploiting the nature of fractions, we have

$$\begin{aligned}0.125 \cdot 16 &= \frac{1}{8} \cdot 16 \\ &= \frac{2 \cdot \cancel{8}}{\cancel{8}} \\ &= 2\end{aligned}$$

### "Cancelling zeros"

A number such as 3000 equals  $3 \cdot 10 \cdot 10 \cdot 10$ , while 700 equals  $7 \cdot 10 \cdot 10$ . Hence, we can simplify  $\frac{3000}{700}$  like this:

$$\begin{aligned}\frac{3000}{700} &= \frac{3 \cdot \cancel{10} \cdot \cancel{10} \cdot 10}{7 \cdot \cancel{10} \cdot \cancel{10}} \\ &= \frac{3 \cdot 10}{7} \\ &= \frac{30}{7}\end{aligned}$$

In practice, this is the same as "cancelling zeros":

$$\frac{300\cancel{0}}{70\cancel{0}} = \frac{30}{7}$$


*Aware!* Zeros are the only digits we can "cancel" this way. For example,  $\frac{12\cancel{3}}{13}$  cannot be simplified in any way. Also, we can only "cancel" zeros which are right-most situated, e.g. we cannot "cancel" zeros in the fraction  $\frac{10\cancel{1}}{10}$ .

# 4.8 Division by fractions

## Division by studying the number line

Let's calculate  $4 : \frac{2}{3}$ . Since the fraction have denominator 3, it could be wise to transform also 4 into a fraction with denominator 3.

$$4 = \frac{12}{3}$$




Recall that one of the interpretations of  $4 : \frac{2}{3}$  is

”The number of  $\frac{2}{3}$ ’s added to make 4.”

By studying a number line, we find that 6 instances of  $\frac{2}{3}$  added together equals 4. Hence

$$4 : \frac{2}{3} = 6$$



## A general method

We can't study the number line every time we are to divide by a fraction, so here we shall find a general method, again with  $4 : \frac{2}{3}$  as our example. In this case, we apply the following interpretation of division:

$$4 : \frac{2}{3} = \text{"The number to multiply } \frac{2}{3} \text{ by to make 4."}$$

We begin the search of this number by multiplying  $\frac{2}{3}$  by the number which results in the product equal to 1. This number is the *inverted fraction* of  $\frac{2}{3}$ , namely  $\frac{3}{2}$ :

$$\frac{2}{3} \cdot \frac{3}{2} = 1$$

Now we only have to multiply by 4 to make 4:

$$\frac{2}{3} \cdot \frac{3}{2} \cdot 4 = 4$$

Therefore, to make 4 we must multiply  $\frac{2}{3}$  by  $\frac{3}{2} \cdot 4$ . Consequently,

$$\begin{aligned} 4 : \frac{2}{3} &= \frac{3}{2} \cdot 4 \\ &= 6 \end{aligned}$$

### 4.10 Fractions divided by fractions

When dividing a number by a fraction, we multiply the number by the inverted fraction.

#### Example 1

$$\begin{aligned} 3 : \frac{2}{9} &= 3 \cdot \frac{9}{2} \\ &= \frac{27}{2} \end{aligned}$$

#### Example 2

$$\begin{aligned} \frac{4}{3} : \frac{5}{8} &= \frac{4}{3} \cdot \frac{8}{5} \\ &= \frac{32}{15} \end{aligned}$$

**Example 3**

$$\begin{aligned}\frac{3}{5} \div \frac{3}{10} &= \frac{3}{5} \cdot \frac{10}{3} \\ &= \frac{30}{15}\end{aligned}$$

In this case we should also observe that the fraction can be simplified:

$$\begin{aligned}\frac{30}{15} &= \frac{2 \cdot \cancel{15}}{\cancel{15}} \\ &= 2\end{aligned}$$

*Note:* Canceling factors along the way saves the labor of working with large numbers:

$$\begin{aligned}\frac{3}{5} \cdot \frac{10}{3} &= \frac{\cancel{3} \cdot 2 \cdot \cancel{5}}{\cancel{5} \cdot \cancel{3}} \\ &= 2\end{aligned}$$

## 4.9 Rational numbers

### 4.11 Rational numbers

Any number which can be expressed as a fraction is a *rational number*.

#### Note

Rational numbers is a collective name of

- **Integers**

For example  $4 = \frac{4}{1}$ .

- **Decimal numbers with a finite number of digits**

For example  $0,2 = \frac{1}{5}$ .

- **Decimal numbers with infinite digits in a repeating manner**

For example <sup>1</sup>  $0.08\bar{3} = \frac{1}{12}$ .

---

<sup>1</sup>  $\bar{3}$  indicates that 3 repeats infinite. Another way of expressing this is by using  $\dots$ . That is,  $0.08\bar{3} = 0.08333333\dots$

### 4.12 Mixed numbers

If we add an integer with a fraction where the numerator has a lower value than the denominator, we have a **mixed number**

#### Example 1

Three different mixed numbers:

$$2 + \frac{5}{7} \qquad 8 + \frac{2}{7} \qquad \frac{1}{10} + 4$$

#### Note

In a lot of texts you will find numbers like those from *Example 1* written like this:

$$2\frac{5}{7} \qquad 8\frac{2}{7} \qquad 4\frac{1}{10}$$



### Example 2

Write the fraction  $\frac{17}{3}$  as a mixed number.

#### Answer

We observe that 17 is the numerator and 3 the denominator. The largest number we can multiply by 3 without getting a product larger than 17 is 5, so

$$\begin{aligned}\frac{17}{3} &= \frac{5 \cdot 3 + 2}{3} \\ &= \frac{5 \cdot \cancel{3}}{\cancel{3}} + \frac{2}{3} \\ &= 5 + \frac{2}{3}\end{aligned}$$

### Example 3

Write the mixed number  $3 + \frac{4}{5}$  as a fraction.

#### Answer

We have  $3 = \frac{3}{1}$ , so

$$3 + \frac{4}{5} = \frac{3}{1} + \frac{4}{5}$$

Moreover<sup>1</sup>,

$$\begin{aligned}\frac{3}{1} + \frac{4}{5} &= \frac{3 \cdot 5}{1 \cdot 5} + \frac{4}{5} \\ &= \frac{15}{5} + \frac{4}{5} \\ &= \frac{19}{5}\end{aligned}$$

---

<sup>1</sup>See [Rule ??](#).

## Exercises for Chapter 4

### 4.1.1

Find the value of the fraction.

- a)  $\frac{18}{3}$       b)  $\frac{20}{4}$       c)  $\frac{10}{5}$       d)  $\frac{42}{6}$       e)  $\frac{63}{7}$       f)  $\frac{32}{8}$




### 4.1.2

Find the value of the fraction. Use a calculator if necessary.

- a)  $\frac{1}{2}$       b)  $\frac{1}{4}$       c)  $\frac{1}{5}$       d)  $\frac{3}{4}$       e)  $\frac{2}{5}$       f)  $\frac{3}{5}$       g)  $\frac{4}{5}$   
f)  $\frac{3}{2}$       g)  $\frac{1}{3}$       h)  $\frac{5}{2}$       i)  $\frac{5}{6}$       j)  $\frac{7}{5}$       k)  $\frac{11}{4}$       l)  $\frac{7}{10}$




### 4.1.3

Write the fraction positioned at the red mark.



### 4.1.4

Write the fraction positioned at the red mark.



### 4.2.1

**Example**

$$\frac{9}{8} \text{ expanded by } 3 = \frac{9 \cdot 3}{8 \cdot 3} = \frac{27}{24}$$

Expand

a)  $\frac{10}{3}$  by 2.      b)  $\frac{3}{4}$  by 3.      c)  $\frac{3}{7}$  by 4.

c)  $\frac{9}{8}$  by 5.      d)  $\frac{9}{5}$  by 6.      e)  $\frac{11}{4}$  by 7.

### 4.2.2

Expand

a)  $\frac{7}{3}$  to a fraction with denominator 15.

b)  $\frac{3}{4}$  to a fraction with denominator 32.

c)  $\frac{10}{9}$  to a fraction with denominator 63.

### 4.2.3

**Example**

$$\frac{10}{8} \text{ simplified by } 2 = \frac{10 : 2}{8 : 2} = \frac{5}{4}$$

Simplify

a)  $\frac{14}{26}$  by 2.      b)  $\frac{15}{12}$  by 3.      c)  $\frac{20}{16}$  by 4.

c)  $\frac{35}{50}$  by 5.      d)  $\frac{54}{18}$  by 6.      e)  $\frac{49}{63}$  by 7.

### 4.2.4

Simplify

a)  $\frac{27}{12}$  to a fraction with denominator 4.

b)  $\frac{36}{20}$  to a fraction with denominator 5.

c)  $\frac{18}{63}$  to a fraction with denominator 7.

### 4.3.1

Calculate.

a)  $\frac{4}{3} + \frac{6}{3}$       b)  $\frac{5}{4} + \frac{9}{4}$       c)  $\frac{1}{6} + \frac{10}{6}$       d)  $\frac{8}{7} + \frac{2}{7}$       e)  $\frac{1}{2} + \frac{1}{2}$

### 4.3.2

Calculate.

a)  $\frac{10}{3} + \frac{4}{3} + \frac{8}{3}$       b)  $\frac{4}{5} + \frac{3}{5} + \frac{1}{5}$       c)  $\frac{11}{7} + \frac{2}{7} + \frac{4}{7}$

### 4.3.3

Calculate.

a)  $\frac{5}{3} - \frac{4}{3}$       b)  $\frac{9}{4} - \frac{7}{4}$       c)  $\frac{11}{6} - \frac{1}{6}$       d)  $\frac{11}{7} - \frac{4}{7}$       e)  $\frac{1}{2} - \frac{1}{2}$

### 4.3.4

Calculate.

a)  $\frac{4}{5} + \frac{3}{5} - \frac{1}{5}$       b)  $\frac{11}{7} - \frac{2}{7} - \frac{4}{7}$       c)  $\frac{10}{3} - \frac{4}{3} + \frac{8}{3}$

### 4.3.5

Calculate.

a)  $\frac{2}{5} + \frac{3}{6}$       b)  $\frac{5}{7} + \frac{4}{9}$       c)  $\frac{10}{3} + \frac{7}{8}$       d)  $\frac{7}{5} + \frac{9}{4}$       e)  $\frac{1}{3} + \frac{1}{2}$

### 4.3.6

Calculate.

a)  $\frac{2}{5} - \frac{3}{10}$       b)  $\frac{5}{4} - \frac{4}{9}$       c)  $\frac{10}{9} - \frac{1}{8}$       d)  $\frac{4}{5} - \frac{1}{4}$       e)  $\frac{5}{2} - \frac{5}{3}$

### 4.3.7

Calculate.

a)  $\frac{2}{3} + \frac{1}{2} - \frac{3}{4}$       b)  $\frac{10}{2} - \frac{1}{6} + \frac{2}{5}$       c)  $\frac{9}{2} - \frac{2}{7} - \frac{1}{8}$

#### 4.4.1

Calculate.

$$\begin{array}{lllll} \text{a)} \frac{4}{3} \cdot 5 & \text{b)} \frac{5}{7} \cdot 8 & \text{c)} \frac{9}{10} \cdot 6 & \text{d)} \frac{8}{7} \cdot 10 & \text{e)} \frac{3}{2} \cdot 7 \\ \text{f)} 7 \cdot \frac{4}{3} & \text{g)} 5 \cdot \frac{7}{3} & \text{h)} 3 \cdot \frac{10}{7} & \text{i)} 1 \cdot \frac{5}{11} & \text{j)} 8 \cdot \frac{9}{17} \end{array}$$

#### 4.5.1

Calculate.

$$\begin{array}{lllll} \text{a)} \frac{4}{3} : 5 & \text{b)} \frac{5}{7} : 8 & \text{c)} \frac{9}{10} : 6 & \text{d)} \frac{8}{7} : 10 & \text{e)} \frac{3}{2} : 7 \\ \text{f)} \frac{9}{10} : 11 & \text{g)} \frac{1}{5} : 12 & \text{h)} \frac{9}{10} : 29 & \text{i)} \frac{8}{9} : 51 & \text{j)} \frac{3}{2} : 79 \end{array}$$

#### 4.6.1

Calculate.

$$\begin{array}{lllll} \text{a)} \frac{4}{3} \cdot \frac{5}{9} & \text{b)} \frac{7}{8} \cdot \frac{1}{4} & \text{c)} \frac{2}{7} \cdot \frac{9}{3} & \text{d)} \frac{10}{3} \cdot \frac{6}{5} & \text{e)} \frac{3}{2} \cdot \frac{7}{5} \\ \text{f)} \frac{2}{3} \cdot \frac{5}{7} & \text{g)} \frac{8}{9} \cdot \frac{2}{3} & \text{h)} \frac{10}{3} \cdot \frac{8}{3} & \text{i)} \frac{4}{5} \cdot \frac{9}{7} & \text{j)} \frac{7}{2} \cdot \frac{5}{6} \end{array}$$

#### 4.7.1

Calculate.

$$\begin{array}{lllll} \text{a)} \frac{3}{10} \cdot \frac{5}{4} & \text{b)} \frac{17}{8} \cdot \frac{9}{4} & \text{c)} \frac{23}{8} \cdot \frac{2}{4} & \text{d)} \frac{7}{81} \cdot \frac{3}{8} & \text{e)} \frac{7}{8} \cdot \frac{29}{41} \end{array}$$

#### 4.8.1

Cancel as many factors as possible.

$$\begin{array}{llll} \text{a)} \frac{3 \cdot 11 \cdot 8}{4 \cdot 8 \cdot 3} & \text{b)} \frac{5 \cdot 12 \cdot 7 \cdot 2}{2 \cdot 8 \cdot 12} & \text{c)} \frac{6 \cdot 10}{6 \cdot 9 \cdot 10} & \text{d)} \frac{7 \cdot 4 \cdot 3}{7 \cdot 3} \end{array}$$

### 4.8.2

Simplify the fraction as much as possible.

- a)  $\frac{28}{16}$       b)  $\frac{18}{42}$       c)  $\frac{24}{36}$       d)  $\frac{56}{49}$       e)  $\frac{25}{50}$       f)  $\frac{21}{14}$

### 4.8.3

**Example**

$$\frac{3}{4} \cdot 20 = \frac{3}{\cancel{4}} \cdot \cancel{4} \cdot 5 = 3 \cdot 5 = 15$$

Exploit that the numerator is a factor in the other factor, and calculate.

- a)  $\frac{7}{3} \cdot 21$       b)  $\frac{9}{5} \cdot 30$       c)  $\frac{10}{7} \cdot 49$       d)  $\frac{8}{9} \cdot 18$       e)  $\frac{5}{4} \cdot 24$   
f)  $8 \cdot \frac{3}{2}$       g)  $35 \cdot \frac{5}{7}$       h)  $63 \cdot \frac{2}{9}$       i)  $48 \cdot \frac{1}{6}$       j)  $27 \cdot \frac{7}{3}$

### 4.9.1

Calculate.

- a)  $4 : \frac{9}{8}$       b)  $7 : \frac{3}{5}$       c)  $10 : \frac{7}{3}$       d)  $5 : \frac{4}{5}$       e)  $2 : \frac{5}{11}$

### 4.9.2

Calculate, and simplify the fraction as much as possible.

- a)  $4 : \frac{8}{9}$       b)  $7 : \frac{21}{5}$       c)  $10 : \frac{5}{3}$       d)  $5 : \frac{5}{4}$       e)  $2 : \frac{8}{11}$

### 4.9.3

Calculate.

- a)  $\frac{2}{3} : \frac{5}{7}$       b)  $\frac{8}{9} : \frac{5}{3}$       c)  $\frac{10}{3} : \frac{7}{3}$       d)  $\frac{1}{5} : \frac{4}{7}$       e)  $\frac{6}{5} : \frac{3}{11}$

#### 4.9.4

**Example**

$$\frac{3}{4} : \frac{15}{8} = \frac{\cancel{3}}{\cancel{2} \cdot \cancel{2}} \cdot \frac{\cancel{2} \cdot \cancel{2} \cdot 2}{\cancel{3} \cdot 5} = \frac{2}{5}$$

*Note:* Here, we have chosen to prime factorize the numbers, however, this is not necessary if you detect the common factors of the numerators and the denominators.

Exploit that the numerators and denominators have common factors, and calculate.

$$\text{a) } \frac{7}{9} : \frac{21}{12} \quad \text{b) } \frac{35}{24} : \frac{7}{18} \quad \text{c) } \frac{84}{55} : \frac{42}{77}$$

### Gruble 3

Apply [Rule 4.7](#) and [Rule 4.8](#) to insert the missing integers in place of " \_".

- a) Multiplying by  $\frac{1}{2}$  is the same as dividing by \_.
- b) Multiplying by  $\frac{1}{4}$  is the same as dividing by \_.
- c) Multiplying by  $\frac{1}{5}$  is the same as dividing by \_.

Look back at the answers of [Exercise 4.1.2a](#))-g). Fill inn the integer missing in place of " \_".

- d) Multiplying by 0,5 is the same as dividing by \_.
- e) Multiplying by 0,25 is the same as dividing by \_.
- f) Multiplying by 0,2 is the same as dividing by \_.
- g) Multiplying by 0,75 is the same as multiplying by \_ and dividing by \_.
- h) Multiplying by 0,4 is the same as multiplying by \_ and dividing by \_.
- i) Multiplying by 0,6 is the same as multiplying by \_ and dividing by \_.
- j) Multiplying by 0,8 is the same as multiplying by \_ and dividing by \_.



#### Gruble 4

Look up [Rule 4.10](#) and the answers of [Exercise 4.1.2a](#)) - g). Fill in the missing integers in place of " \_".

- (a) Dividing by 0,5 is the same as multiplying by \_.
- (b) Dividing by 0,25 is the same as multiplying by \_.
- (c) Dividing by 0,2 is the same as multiplying by \_.
- (d) Dividing by 0,75 is the same as multiplying by \_ and dividing by \_.
- (e) Dividing by 0,4 is the same as multiplying by \_ and dividing by \_.
- (f) Dividing by 0,6 is the same as multiplying by \_ and dividing by \_.
- (g) Dividing by 0,8 is the same as multiplying by \_ and dividing by \_.

#### Gruble 5

Calculate.


a)  $\frac{5}{204} + \frac{7}{198}$       b)  $\frac{11}{350} + \frac{17}{315}$

## Chapter 5

# Negative numbers

# 5.1 Introduction

Earlier we have seen that e.g. 5 on a number line is placed 5 units to the right of 0.




But what if we move in the other direction, that is to the left? The question is answered by introducing *negative numbers*.

## 5.1 Positive and negative numbers


On a number line, the following applies:

- Numbers placed to *the right* of 0 are positive numbers.
- Numbers placed to *the left* of 0 are negative numbers.



However, relying on the number line every time negative numbers are involved would be very inconvenient, therefore we use a symbol to indicate negative numbers. This is  $-$ , simply the same as the symbol of subtraction. From this it follows that 5 is a positive number, while  $-5$  is a negative number. On the number line,

- 5 is placed 5 units to *the right* of 0.
- $-5$  is placed 5 units to *the left* of 0.



Hence, the big difference between 5 and  $-5$  is on which side of 0 the numbers are placed. Since 5 and  $-5$  have the same distance from 0, we say that 5 and  $-5$  have equal *length*.

## 5.2 Length (absolute value/modulus/magnitude)

The length of a number is expressed by the symbol  $||$ .

The length of a positive number equals the value of the number.

The length of a negative number equals the value of the positive number with corresponding digits.

### Example 1

$$|27| = 27$$

### Example 2

$$|-27| = 27$$

### Sign

*Sign* is a collective name of  $+$  and  $-$ .  $+$  is the sign of 5 and  $-$  is the sign of  $-5$ .

# 5.2 The elementary operations

The introduction of negative numbers brings new aspects to the elementary operations. When adding, subtracting, multiplying or dividing by negative numbers, we'll frequently, for clarity, enclose negative numbers by parentheses. Then we'll write e.g.  $-4$  as  $(-4)$ .

## Addition

When adding in [Section 2.1](#)  $+$  implied moving to *the right*. Negative numbers bring an alternation of the interpretation of  $+$  :

$+$  "As long and in *the same* direction as"

Let's study the calculation


$$7 + (-4)$$

Our altered definition of  $+$  implies that

$$7 + (-4) = \text{"7 and as long and in the same direction as } (-4)\text{"}$$

$(-4)$  has length 4 and direction to *the left*. Hence, the calculation tells us to start at 7 and then move the length of 4 to *the left*.

$$7 + (-4) = 3$$



## 5.3 Addition involving negative numbers

Adding a negative number is the same as subtracting the number of equal magnitude.

### Example 1

$$4 + (-3) = 4 - 3 = 1$$

### Example 2

$$-8 + (-3) = -8 - 3 = -11$$

## Notice

Rule 2.1 declares that addition is commutative. This also applies after introducing negative numbers, for example is

$$7 + (-3) = 4 = -3 + 7$$

## Subtraction

In Section 2.2,  $-$  implied moving to *the left*. The interpretation of  $-$  also needs an alternation when working with negative numbers:

$-$  "As long and in the *opposite* direction as"

Let's study the calculation


$$2 - (-6)$$

Our altered definition of  $-$  implies that

$$2 - (-6) = \text{"2 and as long and in the } opposite \text{ direction as } (-6)\text{"}$$

$-6$  have length 6 and direction to *the left*. When moving an equal length, but in the *opposite* direction, we have to move the length of 6 to *the right*<sup>1</sup>. This is equivalent to adding 6:

$$2 - (-6) = 2 + 6 = 8$$



## 5.4 Subtraction involving negative numbers

Subtracting a negative number is the same as adding the positive number of equal magnitude.

### Example 1

$$11 - (-9) = 11 + 9 = 20$$

<sup>1</sup>Once again, recall that the red colored arrow indicates starting at the arrowhead, then moving to the other end.

### Example 2

$$-3 - (-7) = -3 + 7 = 4$$

## Multiplication

In [Section 2.3](#), multiplication by positive integers were introduced as repeated addition. By our altered interpretations of addition and subtraction we can now also alternate the interpretation of multiplication:

### 5.5 Multiplication by positive and negative integers

- Multiplication by a positive integer corresponds to repeated addition.
- Multiplication by a negative integer corresponds to repeated subtraction.

### Example 1

$$\begin{aligned} 2 \cdot 3 &= \text{"As long and in the same direction as 2, 3 times"} \\ &= 2 + 2 + 2 \\ &= 6 \end{aligned}$$

### Example 2

$$\begin{aligned} (-2) \cdot 3 &\equiv \text{"As long and in the same direction as } (-2), 3 \text{ times"} \\ &= -2 - 2 - 2 \\ &= -6 \end{aligned}$$

### Example 3

$$\begin{aligned} 2 \cdot (-3) &\equiv \text{"As long and in the opposite direction as 2, 3 times"} \\ &= -2 - 2 - 2 \\ &= -6 \end{aligned}$$

### Example 4

$$\begin{aligned} (-3) \cdot (-4) &\equiv \text{As long as in the opposite direction as } -3, 4 \text{ times} \\ &= 12 \end{aligned}$$

### Multiplication is commutative

*Example 2* and *Example 3* on page 86 illustrates that [Rule 2.2](#) also implies after introducing negative numbers:

$$(-2) \cdot 3 = 3 \cdot (-2)$$

It would be laborious to calculate multiplication by repeated addition/subtraction every time a negative number were involved, however, as a direct consequence of [Rule 5.5](#) we can make the two following rules:

### 5.6 Multiplication by negative numbers I

The product of a negative number and a positive number is a negative number.

The magnitude of the factors multiplied together yields the magnitude of the product.

### Example 1

Calculate  $(-7) \cdot 8$

#### Answer

Since  $7 \cdot 8 = 56$ , we have  $(-7) \cdot 8 = -56$

### Example 2

Calculate  $3 \cdot (-9)$ .

#### Answer

Since  $3 \cdot 9 = 27$ , we have  $3 \cdot (-9) = -27$



## 5.7 Multiplication by negative numbers II

The product of two negative numbers is a positive number.

The magnitude of the factors multiplied together yields the value of the product.

### Example 1

$$(-5) \cdot (-10) = 5 \cdot 10 = 50$$

### Example 2

$$(-2) \cdot (-8) = 2 \cdot 8 = 16$$

## Division

From the definition of division (see [Section 2.4](#)), combined with what we now know about multiplication involving negative numbers, it follows that

$-18 : 6 =$  "The number which yields  $-18$  when multiplied by  $6$  "

$$6 \cdot (-3) = -18, \text{ hence } -18 : 6 = -3$$

$42 : (-7) =$  "The number which yields  $42$  when multiplied by  $-7$ "

$$(-7) \cdot (-8) = 42, \text{ hence } 42 : (-7) = -8$$

$-45 : (-5) =$  "The number which yields  $-45$  when multiplied by  $-5$ "

$$(-5) \cdot 9 = -45, \text{ hence } -45 : (-5) = 9$$

## 5.8 Division involving negative numbers

Division between a positive number and a negative number yields a negative number.

Division between two negative numbers yields a positive number.

The magnitude of the dividend divided by the magnitude of the divisor yields the magnitude of the quotient.

### Example 1

$$-24 : 6 = -4$$

### Example 2

$$24 : (-2) = -12$$

### Example 3

$$-24 : (-3) = 8$$

### Example 4

$$\frac{2}{-3} = -\frac{2}{3}$$

### Example 5

$$\frac{-10}{7} = -\frac{10}{7}$$

## 5.3 Negative numbers as amounts

*Notice: This view of negative numbers will first come into use in [Section 9.2](#), a section a lot of readers can skip without loss of understanding.*

So far, we have studied negative number by the aid of number lines. Studying negative numbers as amounts is at first difficult because negative amounts makes no sense! To make an interpretation of negative numbers through the perspective of amounts, we'll use what we shall call the *weight principle*. Then we look upon the numbers as amounts of forces. The positive numbers are amounts of forces acting downwards while the negative numbers are amounts of forces working upwards<sup>1</sup>. In this way, the results of calculations involving positive and negative numbers can be looked upon as the result of weighing the amounts. Hence, a positive number and a negative number of equal magnitude will cancel each other.

### 5.9 Negative numbers as amounts

Negative numbers will be illustrated as a light blue amount:

$$\boxed{\text{light blue}} = -1$$

#### Example

$$1 + (-1) = 0$$

$$\boxed{\text{dark blue}} + \boxed{\text{light blue}} = 0$$

---

<sup>1</sup>From reality one can look upon the positive and the negative numbers as balloons filled with air and helium, respectively. Balloons filled with air acts with a force downwards (they fall), while balloons filled with helium acts with a force upwards (they rise).

## Exercises for Chapter 5

### 5.1.1

#### Example


8 is directed to the right, and has length 8.

$-7$  is directed to the left, and has length 7.

Fill in the missing words.

- a) 9 is directed to the ....., and has length ....
- b) 4 is directed to the ....., and has length ....
- c)  $-3$  is directed to the ....., and has length ....
- d) 12 is directed to the ....., and has length ....
- e)  $-11$  is directed to the ....., and has length ....
- f)  $-25$  is directed to the ....., and has length ....

### 5.1.2



draw the number line, and place the the numbers on it.

- a) 3      b)  $-4$       c)  $-8$       d) 7      e)  $-3$       f) 5      g)  $-5$

### 5.1.3

Look back at the numbers from [Exercise 5.1.2](#). Write down which numbers are

- a) positive numbers      b) negative numbers

### 5.2.1

Calculate.

- a)  $8 + (-7)$       b)  $12 + (-5)$       c)  $9 + (-3)$       d)  $7 + (-7)$   
e)  $-5 + 8$       f)  $-9 + 10$       g)  $-1 + 11$       h)  $-4 + 9$

### 5.2.2

Calculate.

- a)  $3 + (-19)$       b)  $7 + (-15)$       c)  $-20 + (-3)$       d)  $7 + (-7)$   
e)  $-4 + (-19)$       f)  $-2 + (-15)$       g)  $-8 + 5$       h)  $-6 + 6$

### 5.2.3

Calculate.

- a)  $8 - (-7)$       b)  $12 - (-5)$       c)  $9 - (-3)$       d)  $7 - (-7)$   
e)  $-5 - 8$       f)  $-9 - 10$       g)  $-1 - 11$       h)  $-4 - 9$

### 5.2.4

Calculate.

- a)  $3 - (-19)$       b)  $7 - (-15)$       c)  $-20 - (-3)$       d)  $7 - (-7)$   
e)  $-4 - (-19)$       f)  $-2 - (-15)$       g)  $-8 - 5$       h)  $-6 - 6$

### 5.2.5

Calculate.

- a)  $3 \cdot (-4)$       b)  $5 \cdot (-10)$       c)  $7 \cdot (-9)$       d)  $4 \cdot (-6)$   
e)  $(-7) \cdot 8$       f)  $(-3) \cdot 9$       g)  $(-1) \cdot 12$       h)  $(-10) \cdot 4$   
i)  $(-3) \cdot 7$       j)  $(-5) \cdot 5$       k)  $(-6) \cdot 2$       l)  $(-8) \cdot 9$

### 5.2.6

Calculate.

- a)  $(-32) : 8$       b)  $(-42) : 7$       c)  $(-30) : 6$       d)  $(-20) : 5$   
e)  $72 : (-9)$       f)  $63 : (-7)$       g)  $50 : (-10)$       h)  $25 : (-5)$   
i)  $(-72) : (-9)$       j)  $(-63) : (-7)$       k)  $(-50) : (-10)$

## Chapter 6

# Calculation methods

# 6.1 Addition

## Column addition

This method builds on the base-10 positional notation, in turn adding the ones, the tens, the hundreds and so on.

### Example 1

$$\begin{array}{r} 234 \\ + 612 \\ \hline = 846 \end{array}$$

### Example 2

$$\begin{array}{r} 273 \\ + 86 \\ \hline = 359 \end{array}$$

### Example 3

$$\begin{array}{r} 85 \\ + 79 \\ \hline = 164 \end{array}$$

### Example 4

$$\begin{array}{r} 397,2 \\ + 85,9 \\ \hline = 482,1 \end{array}$$

### Example 1 (explanation)

$$\begin{array}{r} 234 \\ + 612 \\ \hline \end{array}$$

(a)

$$\begin{array}{r} 234 \\ + 612 \\ \hline \end{array}$$

(b)

$$\begin{array}{r} 234 \\ + 612 \\ \hline = 846 \end{array}$$

(c)

a) We add the ones:  $4 + 2 = 6$

b) We add the tens:  $3 + 1 = 4$

c) We add the hundreds:  $2 + 6 = 8$

## Example 2 (explanation)

$$\begin{array}{|c|c|c|} \hline 2 & 7 & 3 \\ \hline + & 8 & 6 \\ \hline & & 9 \\ \hline \end{array}$$

(a)

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 7 & 3 \\ \hline + & 8 & 6 \\ \hline & 5 & 9 \\ \hline \end{array}$$

(b)

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 7 & 3 \\ \hline + & 2 & 7 & 3 \\ \hline = & 3 & 5 & 9 \\ \hline \end{array}$$

(c)

- a) We add the ones:  $3 + 6 = 9$
- b) We add the tens:  $7 + 8 = 15$ . Since 10 tens equals 100, we add 1 to the hundreds position, and write the remaining 5 tens at the tens position.
- c) We add the hundreds:  $1 + 2 = 3$ .

## The language box

Writing 1 on a place value to the left is called "carrying 1 over".



# 6.2 Subtraction

## Column subtraction

This method is founded on the base-10 positional notation, in turn subtracting the ones, the tens, the hundreds and so on. It is also based on the perspective of numbers as amounts, so it does not allow negative differences. (see the explanation of *Example 2*).

**Example 1**

$$\begin{array}{r}
 789 \\
 - 324 \\
 \hline
 = 465
 \end{array}$$

**Example 2**

$$\begin{array}{r}
 93 \\
 - 57 \\
 \hline
 = 36
 \end{array}$$

**Example 3**

$$\begin{array}{r}
 504 \\
 - 478 \\
 \hline
 = 86
 \end{array}$$

**Example 4**

$$\begin{array}{r}
 2061 \\
 - 3174 \\
 \hline
 = 1744
 \end{array}$$

### Example 1 (explanation)

$$\begin{array}{r}
 789 \\
 - 324 \\
 \hline
 \phantom{0} \phantom{0} 5
 \end{array}$$

(a)

$$\begin{array}{r}
 789 \\
 - 324 \\
 \hline
 \phantom{0} 6 \phantom{0} 5
 \end{array}$$

(b)

$$\begin{array}{r}
 789 \\
 - 324 \\
 \hline
 = 465
 \end{array}$$

(c)

- (a) We find the difference between the ones:  $9 - 4 = 5$
- (b) We find the difference between the tens:  $8 - 2 = 6$ .
- (c) We find the difference between the hundreds:  $7 - 3 = 4$ .

### Example 2 (explanation)

$\begin{array}{r} \overset{10}{\cancel{9}} \overset{10}{3} \\ - 5 \overset{10}{7} \\ \hline \phantom{0} 6 \end{array}$	$\begin{array}{r} \overset{10}{\cancel{9}} \overset{10}{3} \\ - \overset{10}{5} \overset{10}{7} \\ \hline = \overset{10}{3} \overset{10}{6} \end{array}$
(a)	(b)

- (a) We notice that 7 is larger than 3, and thus we take 1 ten from the 9 at the tens position. This is marked by drawing a line across 9. Then we find the difference between the ones:  $13 - 7 = 6$
- (b) Since we took 1 from the 9 tens, it is now only 8 tens. We find the difference between the tens:  $8 - 5 = 3$ .

### The table method

The table method takes advantage of subtraction being the inverse operation of addition. For example, the answer to the question "What is  $789 - 324$ ?" is the same as the answer to the question "How much must i add to 324 in order to get 789?". With the table method you can freely chose which numbers to add as long as you end up with the targeted number.

#### Example 1

$$789 - 324 = 465$$

	<span style="color: red;">324</span>
6	330
70	400
389	<span style="color: blue;">789</span>
<span style="color: blue;">465</span>	

#### Example 2

$$83 - 67 = 16$$

	<span style="color: red;">67</span>
3	70
13	<span style="color: blue;">83</span>
<span style="color: blue;">16</span>	

**Example 3**

$$564 - 478 = 86$$

	478
2	480
20	500
64	564
86	

**Example 4**

$$206,1 - 31,7 = 174,4$$

	31,7
0,3	32
70	102
104,1	206,1
174,4	

**Example 1 (explanation)**

$$789 - 324 = 465$$

	324

(a)

	324
6	330

(b)

	324
6	330
70	400

(c)

	324
6	330
70	400
389	789

(d)

	324
6	330
70	400
389	789
465	

(e)

- (a) We start at 324.
- (b) We add 6, and get  $324 + 6 = 330$
- (c) We add 70, and get  $70 + 330 = 400$
- (d) We add 389, and get  $389 + 400 = 789$ . Now we have reached 789.
- (e) We find the sum of the numbers we added:  
 $6 + 70 + 389 = 465$

## 6.3 Multiplication

Multiplying by 10, 100, 1 000 etc.

### 6.1 Å gange heltall med 10, 100 osv.

- When multiplying an integer by 10, the product can be found by adding the digit 0 behind the integer.
- When multiplying an integer by 100, the product can be found by adding the digits 00 behind the integer.
- The same pattern applies for the numbers 1 000, 10 000 etc.

#### Example 1

$$6 \cdot 10 = 60$$

$$79 \cdot 10 = 790$$

$$802 \cdot 10 = 8020$$

#### Example 2

$$6 \cdot 100 = 600$$

$$79 \cdot 100 = 7\,900$$

$$802 \cdot 100 = 80\,200$$

#### Example 3

$$6 \cdot 1\,000 = 6\,000$$

$$79 \cdot 10\,000 = 790\,000$$

$$802 \cdot 100\,000 = 80\,200\,000$$

## 6.2 Multiplying decimal numbers by 10, 100, etc.

- When multiplying an integer by 10, the product is found by moving the dot one position to the right.
- When multiplying an integer by 100, the product is found by moving the dot one position to the right.
- The same pattern applies for the numbers 1 000, 10 000 etc.

### Example 1

$$7.\dot{9} \cdot 10 = 79.\dot{ } = 79$$

$$38.\dot{0}2 \cdot 10 = 380.\dot{2} = 380.2$$

$$0.\dot{5}7 \cdot 10 = 05.\dot{7} = 5.7$$

$$0.\dot{1}94 \cdot 10 = 01.\dot{9}4 = 1.94$$

### Example 2

$$7.\dot{9} \cdot 100 = 790.\dot{ } = 790$$

$$38.\dot{0}2 \cdot 100 = 3802.\dot{ } = 3\,802$$

$$0.\dot{5}7 \cdot 100 = 057.\dot{ } = 57$$

$$0.\dot{1}94 \cdot 100 = 019.\dot{4} = 19.4$$

### Example 3

$$7.\dot{9} \cdot 1\,000 = 7900.\dot{ } = 7\,900$$

$$38.\dot{0}2 \cdot 10\,000 = 380020.\dot{ } = 380\,200$$

$$0.\dot{5}7 \cdot 100\,000 = 57000.\dot{ } = 57\,000$$

### Note

[Rule 6.1](#) is just a special case of [Rule 6.2](#). For example, applying [Rule 6.1](#) when calculating  $7 \cdot 10$  yields the same answer as when applying [Rule 6.2](#) when calculating  $7,0 \cdot 10$ .

## Multiplying by 10, 100 etc. (explanation)

The Base-10 positional notation is founded on groups of tens, hundreds, thousands etc., and tenths, hundredths, thousandths etc. (see [Rule 1.2](#)). When multiplying a number by 10, all the ones in the number will form a group of tens, all the tens will form a group of hundreds and so on. Hence, every digit is moved one position to the left. Similarly, every digit is moved one position to the left when multiplying by 100, three places when multiplying by 1 000 etc.

## Expanded form

Multiplication with expanded form can be applied on multi digit numbers. The method is based on the distributive law ([Rule 3.2](#)).

### Example 1

$$\begin{array}{|c|c|c|c|c|c|c|} \hline 2 & 4 & \cdot & 3 & = & 7 & 2 \\ \hline 20 & \cdot & 3 & = & 60 & \\ 4 & \cdot & 3 & = & 12 & \\ \hline & & & & 72 & \\ \hline \end{array}$$

### Example 2

$$279 \cdot 34 = 9486$$

$$\begin{array}{rcl} 200 \cdot 30 = 6000 & 200 \cdot 4 = 800 & 8370 \\ 70 \cdot 30 = 2100 & 70 \cdot 4 = 280 & 1116 \\ 9 \cdot 30 = 270 & 9 \cdot 4 = 36 & 9486 \\ \hline 8370 & 1116 & \end{array}$$

### Example 1 (explanation)

24 can be written as  $20 + 4$ , so

$$24 \cdot 3 = (20 + 4) \cdot 3$$

Moreover, by [Rule 3.2](#),

$$\begin{aligned} (20 + 4) \cdot 3 &= 20 \cdot 3 + 4 \cdot 3 \\ &= 60 + 12 \\ &= 72 \end{aligned}$$

We have

$$34 = 30 + 4$$

$$279 \cdot 34 = (200 + 70 + 9) \cdot (30 + 4)$$
$$\begin{aligned}(200 + 70 + 9) \cdot (30 + 4) &= 200 \cdot 30 + 70 \cdot 30 + 9 \cdot 30 + 200 \cdot 4 + 70 \cdot 4 + 9 \cdot 4 \\ &= 9486\end{aligned}$$

The compact method is based on the same principles as the expanded form method, only with a shorter way of writing.

$$279 \cdot 34 = 9486$$

$$\begin{array}{r} 23 \\ 886 \\ 22 \\ 617 \\ \hline 9486 \end{array}$$

### Example 1 (explanation)

First we multiply the digits of 279 by 4:

- $9 \cdot 4 = 36$ , so we write 6 at the ones position and carry over 3.
- $7 \cdot 4 = 28$ , so we write 8 at the tens position and carry over 2.
- $2 \cdot 4 = 8$ , so we write 8 at the hundreds position.

Then we multiply the digits of 279 by 30. This in turn can be simplified to multiplying by 3, as long as we shift the digits one position to the left, relative to when we multiplied by 4:

- $9 \cdot 3 = 27$ , so we write 7 at the tens position and carry over 2.
- $7 \cdot 3 = 21$ , so we write 1 at the hundreds position and carry over 2.
- $2 \cdot 3 = 6$ , so we write 6 at the thousands position.



## 6.4 Division

Division by 10, 100, 1 000 etc.

### 6.3 Deling med 10, 100, 1 000 osv.

When dividing a decimal number by 10, the quotient is found by moving the dot one position to the left.

When dividing a decimal number by 100, the quotient is found by moving the dot two positions to the left.

The same pattern applies for the numbers 1 000, 10 000 etc.

#### Example 1

$$\begin{aligned}200 : 10 &= 200.\textcolor{red}{0} : 10 \\&= 20.\textcolor{red}{00} \\&= 20\end{aligned}$$

$$\begin{aligned}45 : 10 &= 45.\textcolor{red}{0} : 10 \\&= 4.\textcolor{red}{50} \\&= 4.\textcolor{red}{5}\end{aligned}$$

#### Example 2

$$\begin{aligned}200 : 100 &= 200.\textcolor{red}{0} : 100 \\&= 2.\textcolor{red}{000} \\&= 2\end{aligned}$$

$$\begin{aligned}45 : 100 &= 45.\textcolor{red}{0} : 100 \\&= 0.\textcolor{red}{450} \\&= 0.\textcolor{red}{45}\end{aligned}$$

### Example 3

$$143.\dot{7} : 10 = 14.\dot{3}7$$

$$143.\dot{7} : 100 = 1.\dot{4}37$$

$$143.\dot{7} : 1\,000 = 0.14\dot{3}7$$

### Example 4

$$93.\dot{6} : 10 = 9.\dot{3}6$$

$$93.\dot{6} : 100 = 0.9\dot{3}6$$

$$93.\dot{6} : 1\,000 = 0.09\dot{3}6$$

### Division by 10, 100, 1 000 osv. (explanation)

The Base-10 positional notation is founded on groups of tens, hundreds, thousands etc., and tenths, hundredths, thousandths etc. (see [Rule 1.2](#)). When dividing a number by 10, all the ones in the number will form a group of tens, all the tens will form a group of ones and so on. Hence, every digit is moved one position to the right. Similarly, every digit is moved two positions to the right when multiplying by 100, three places when multiplying by 1 000 etc.

### Long division

Long division is based on the perspective of numbers as amount (see page 28).


#### Example 1

9	2	:	4	=	2	3
8						
1	2					
1	2					
	0					

#### Example 1


8	9	4	:	3	=	2	9	8
6								
2	9							
2	7							
	2	4						
	2	4						
		0						

### Example 1 (explanation)




The above figure illustrates the amount 92, which we shall divide into 4 equal groups.

- We start by distributing as many tens as possible. Of the 9 tens, each group can get 2. In total we have distributed  $2 \cdot 4 = 8$  tens.



- Now we are left with 1 ten and 2 ones, which equals 12 ones. Of the 12 ones, each group can get 3. In total we have distributed  $3 \cdot 4 = 12$  ones.



- The amount we started with, 92, is now equally distributed, and our calculation is done. In each group we got 23.

**The table method**

The table method takes advantage of division being the inverse operation of multiplication. For example, the answer to the question "What is  $76 : 4$ ?" is the same as the answer to the question "Which number yields 76 when multiplied by 4?". In the same way as with the table method for subtraction, the numbers can be chosen more ore less freely.

Example 1

$92 : 4 = 23$

· 4		
10	40	40
10	40	80
3	12	92
23		

Example 2

$894 : 3 = 298$

· 3		
200	600	600
30	90	690
30	90	780
30	90	870
8	24	894
298		

Example 3

$894 : 3 = 298$

· 3		
300	900	900
−2	−6	894
298		

Note: Same task as in

Example 2 but a different

calculation.

### Example 1 (explanation)

Since we are to divide 92 by 4, we multiply numbers by 4 until we reach 92.

·4		
10	40	40

(a)

·4		
10	40	40
10	40	80

(b)

·4		
10	40	40
10	40	80
3	12	92

(c)

·4		
10	40	40
10	40	80
3	12	92
23		

(d)

- (a) We multiply 10 by 4, which equals 40. So far we have reached 40.
- (b) We multiply 10 by 4, which equals 40. So far we have reached  $40 + 40 = 80$ .
- (c) We multiply 3 by 4, which equals 12. So far we have reached  $80 + 12 = 92$ , which was our target.
- (d) We add the numbers we multiplied by 4:  $10 + 10 + 3 = 23$ .

### Tip

It is wise to look back at calculations done with the table method and search for improvements. In *Example 1* on page 107, we could have started off multiplying by 20. This is almost as easy as multiplying by 10, and it takes us closer to the target.

## Divisjon med rest

The value of a division calculation is far from always an integer. One way of expressing the answer is by using the term **remainder**. What it means is best explained through an example:


### Example 1

Calculate  $11 : 4$  using remainder.

#### Answer

The largest integer we can multiply by 4 without having a product exceeding 11 is 2.  $2 \cdot 4 = 8$ , so we have  $11 - 8 = 3$  left to reach 11.

$$11 = 2 \cdot 4 + 3$$



This means that

$$11 : 4 = 2 \text{ and } 3 \text{ in remainder}$$


### Example 2

Calculate  $19 : 3$  using remainder.

#### Answer

The largest integer we can multiply by 3 without exceeding 19 is 6.  $6 \cdot 3 = 18$ , so we have  $19 - 18 = 1$  left to reach 19.

$$19 = 6 \cdot 3 + 1$$



This means that

$$19 : 3 = 6 \text{ and } 1 \text{ in remainder}$$

### Example 3

Calculate  $94 : 4$  using remainder.

#### Answer

Using long division

$$94 : 4 = 23 \text{ og } 2 \text{ i rest}$$

9	4	:	4	→	2	3
8						
1	4					
1	2					
	2					

Note: Since it's wrong to use `=` in the above calculation, we have chosen to use `→`.

Using the table method

$$94 : 4 = 23 \text{ and } 2 \text{ in remainder}$$

· 4			
20	80	80	
3	12	92	$94 - 92 = 2$
23			

### The language box

Performing a **modulo-operation**, we find the remainder of a division calculation. This is often abbreviated as `mod`. For example is

$$11 \bmod 4 = 3 \quad , \quad 19 \bmod 3 = 1$$

In addition to `mod`, `%` and `//` are common symbols for this operation in programming languages.

## Divisjon using mixed numbers

### Example 1

Calculate  $11 : 4$  using mixed numbers.

**Answer**

$$11 : 4 = 2 \text{ and } 3 \text{ as remainder} = 2 + \frac{3}{4}$$

### Example 2

Calculate  $19 : 3$  using mixed numbers.

**Answer**

$$19 : 3 = 6 \text{ and } 1 \text{ in remainder} = 6 + \frac{1}{3}$$

### Example 1 (explanation)

We start by noticing that  $4 = \frac{4}{1}$ . Hence

$$11 : 4 = 11 : \frac{4}{1}$$

From [Rule 4.10](#) it follows that

$$11 : \frac{4}{1} = 11 \cdot \frac{1}{4}$$

Moreover,  $11 = 2 \cdot 4 + 3$ , and thus

$$11 \cdot \frac{1}{4} = (2 \cdot 4 + 3) \cdot \frac{1}{4}$$

Now, by [Rule 3.2](#),

$$\begin{aligned} (2 \cdot 4 + 3) \cdot \frac{1}{4} &= 2 \cdot 4 \cdot \frac{1}{4} + 3 \cdot \frac{1}{4} \\ &= 2 + \frac{3}{4} \end{aligned}$$



### 6.4.1 Division using decimal numbers

#### Example 1

Calculate  $11 : 4$  using a decimal number.

#### Answer

With long division

$$11 : 4 = 2,75$$

$$\begin{array}{r|l} 11 : 4 = 2,75 \\ \hline \begin{array}{r} 8 \\ 30 \\ 28 \\ 20 \\ 20 \\ 0 \end{array} \end{array}$$

With the table method

$$11 : 4 = 2,75$$

· 4		
2	8	8
0,5	2	10
0,25	1	11
2,75		

#### Example 1; long division (explanation)

Since we divide 4, we separate 11 in 4 equal groups.

- We can equally distribute 8 of the 11 ones in 4 groups.  
Then there are 3 ones left. This equals 30 tenths.
- 28 of the 30 tenths can be equally distributed in 4 groups.  
Then there are 2 tenths left. This equals 20 hundredths.
- 20 of the 20 hundredths can be equally distributed in 4 groups.
- Now the whole amount of 11 is equally distributed, so our calculation is finished.

# 6.5 Calculation with time

Seconds, minutes and hours are organized in groups of 60:

$$1 \text{ minute} = 60 \text{ second}$$

$$1 \text{ hour} = 60 \text{ minute}$$

This means that *crossovers* arise in the calculations when we reach 60.

## Example 1

$$2 \text{ h } 25 \text{ min} + 10 \text{ h } 45 \text{ min} = 13 \text{ h } 10 \text{ min}$$

### Method 1

		10 h 45 min
15 min	15 min	11 h 00 min
10 min	25 min	11 h 10 min
2 h	2 h 25 min	13 h 10 min

### Method 2

		10:45
00:15	00:15	11:00
00:10	00:25	11:10
02:00	02:25	13:10

## Example 2

$$14 \text{ h } 18 \text{ min} - 9 \text{ h } 34 \text{ min} = 4 \text{ h } 44 \text{ min}$$

### Method 1

	9 h 34 min
26 min	10 h 00 min
18 min	10 h 18 min
4 h	14 h 00 min
4 h 44 min	

### Method 2

	09:34
00:26	10:00
00:18	10:18
04:00	14:18
04:44	

## 6.6 Rounding and estimates

### Rounding


When **rounding**, we decrease the amounts of digits other than 0 in a number. Also, we can round off to the *closest one*, *the closest ten* and such.

#### Example 1

When rounding off to the *closest ten* we round off

- 1, 2, 3 and 4 *down* to 0, because they are closer to 0 than to 10.
- 6, 7, 8 and 9 *up* to 10 because they are closer to 10 than to 0.

5 avrundes også opp til 10.



#### Example 2


- **63 rounded off to the closest 10 = 60**

Because 63 is closer to 60 than 70.




- **78 rounded off to the closest ten = 80**

Because 78 is closer to 80 than 70.



- **359 rounded off to the closest hundred = 400**

Because 359 is closer to 400 than 300.



- **11,8 rounded off to the closest one = 12**

Because 11,8 is closer to 12 than 11.



### 6.6.1 Estimate

Rather than knowing the exact answer to a calculation, at times there is more important to quickly determine the outcome *approximately*, preferably by mental arithmetic. When finding an approximate answer, the result is called an **estimate**. An estimate involves rounding off<sup>1</sup> numbers so that the calculation is easier to perform.

#### The language box

That something is "about the same as" is often written as "circa" ("ca."). The symbol for "circa" is  $\approx$ .

### Estimates when adding or multiplying

Let us estimate the calculation

$$98.2 + 24.6$$

We have  $98.2 \approx 100$ . If we write 100 instead of 98.2 in our calculation, the result will slightly exceed the exact answer. Therefore, if we are to round off 24.6 we should round it off downwards. 24.6 is pretty close to 20, we can write

$$98.2 + 24.6 \approx 100 + 20 = 120$$

As this example showcases, when making an estimate involving addition, we should try to round off one of the numbers (upwards) and the other number (downwards).

---

The same applies for estimates involving multiplication. Let us estimate

$$1\,689 \cdot 12$$

We round off 12 to 10. In that case, the estimate will result in an answer lower than the exact answer, so to account for this we round off 1 689 up to 1 700. Now

$$1\,689 \cdot 12 \approx 1\,700 \cdot 10 = 17\,000$$

---

<sup>1</sup>Note: Rounding off when performing estimates does not necessarily involve rounding off to the closest one, ten or such.

## Estimation when subtracting and dividing

Let us make an estimate of

$$186,4 - 28,9$$

If we round off 186,4 up to 190, the result will be slightly larger than the exact answer. To account for this, we should subtract a bit more than in the original calculation. That can be done by rounding off 28,9 up (to 30):

$$\begin{aligned} 186,4 - 28,9 &\approx 190 - 30 \\ &= 160 \end{aligned}$$

---

The same principle applies when an estimate involves division. Let us estimate

$$145 : 17$$

We round off 17 up to 20. This will make our result a bit smaller than the exact answer. Hence, we should also round off 145 upwards (to 150):

$$145 : 17 \approx 150 : 20 = 75$$

## Estimation, summary

### 6.4 Estimation

- Ved addisjon eller multiplikasjon mellom to tall, avrund gjerne et tall opp og et tall ned.
- Ved subtraksjon eller deling mellom to tall, avrund gjerne begge tall ned eller begge tall opp.

### Example

Estimate the calculations.

a)  $23,1 + 174,7$       b)  $11,8 \cdot 107,2$

c)  $37,4 - 18,9$       d)  $1054 : 209$

### Answer

a)  $32,1 + 174,7 \approx 30 + 170 = 200$

b)  $11,8 \cdot 107,2 \approx 10 \cdot 110 = 1\,100$

c)  $37,4 - 18,9 \approx 40 - 20 = 20$

d)  $1\,054 : 209 \approx 1\,000 : 200 = 5$

### Comment

There are no specific rules for what you *can* or *can not* do when making an estimate. Thereby, [Rule 6.4](#) is strictly speaking not a rule but rather a useful tip.

It is also natural to raise the question how far away from the exact answer an estimate is allowed to be. Neither in this case are there any rules to follow. However, an estimate and the exact answer should have the same **order of magnitude**. Simply put, if the exact answer has something to do with thousands, then so should also the estimate. To be more concise, the exact answer and the estimate shall have the same power of 10 when written in standard form<sup>1</sup>.

---

<sup>1</sup>See [Section 6.7](#)

## 6.7 Standard form

*Note: In this section it is taken for granted that the reader is familiar with powers, which we study in [Section 8.2](#).*

We can apply [Rule 6.2](#) and [Rule 6.3](#), and what we know about powers, to write numbers in **standard form**.

Let us look at the number 6 700. By [Rule 6.2](#),

$$6\,700 = 6.7 \cdot 1\,000$$

Since  $1000 = 10^3$ , we have

$$6\,700 = 6.7 \cdot 1\,000 = 6.7 \cdot 10^3$$

$6.7 \cdot 10^3$  or 6 700 written in standard form because

- 6.7 is larger than or equal to 1, and smaller than 10.
- $10^3$  is a power with base 10 and exponent 3, which is an integer.
- The product of 6.7 and  $10^3$  equals 6 700.

---

Let us look at the number 0.093. By [Rule 6.3](#),

$$0.093 = 9.3 : 100$$

Dividing by 100 is the same as multiplying by  $10^{-2}$ , so

$$0.093 = 9.3 : 100 = 9.3 \cdot 10^{-2}$$

$9.3 \cdot 10^{-2}$  is 0,093 written in standard form because

- 9.3 is larger than or equal to 1, and smaller than 10.
- $10^{-2}$  is a power with base 10 and exponent  $-2$ , which is an integer.
- The product of 9.3 and  $10^{-2}$  equals 0.093.

## 6.5 Standardform

A number written on the form

$$a \cdot 10^n$$

where  $1 \leq |a| < 10$  and  $n$  is an integer, is a number written in **standard form**.

### Example 1

Write 980 in standard form.

**Answer**

$$980 = 9.8 \cdot 10^2$$

### Example 2

Write 0.00671 in standardform.

**Answer**

$$0.00671 = 6.71 \cdot 10^{-3}$$

### Tip

To write numbers in standard form you can do the following:

1. Move the decimal separator so that you get a number between 1 and 10.
2. Multiply this number by a power of ten with exponent equal to the amount of places you moved the decimal separator. If you moved the decimal separator to the left/right, the exponent is positive/negative.



**Example 3**

Write 9 761 432 in standard form.

**Answer**

1. We move the decimal separator 6 places to the left, and get 9.761432
2. We multiply this number by  $10^6$ , and get

$$9\,761\,432 = 9.761432 \cdot 10^6$$

**Example 4**

Write 0.00039 in standard form

**Answer**

1. We move the decimal separator 4 places to the right, and get 3,9.
2. We multiply this number by  $10^{-4}$ , and get

$$0,00039 = 3,9 \cdot 10^{-4}$$

## Exercises for Chapter 6

### 6.1.1

Calculate.

- a)  $12 + 84$       b)  $36 + 51$       c)  $328 + 571$       d)  $242 + 56$

### 6.1.2

Calculate.

- a)  $19 + 84$       b)  $86 + 57$       c)  $529 + 471$       d)  $202 + 808$

### 6.2.1

Calculate.

- a)  $84 - 23$       b)  $286 - 52$       c)  $529 - 401$       d)  $782 - 131$

### 6.2.2

Calculate.

- a)  $78 - 19$       b)  $824 - 499$       c)  $731 - 208$       d)  $1078 - 991$

### 6.3.1

Calculate.

- a)  $12 \cdot 3$       b)  $28 \cdot 4$       c)  $76 \cdot 5$       d)  $43 \cdot 6$   
e)  $109 \cdot 7$       f)  $98 \cdot 8$       g)  $213 \cdot 9$

### 6.3.2

Calculate.

- a)  $29 \cdot 12$       b)  $83 \cdot 31$       c)  $91 \cdot 76$       d)  $14 \cdot 83$

### 6.3.3

Calculate.

- a)  $531 \cdot 56$       b)  $83 \cdot 701$       c)  $91 \cdot 673$       d)  $731 \cdot 67$

### 6.3.4

- a) Use a calculator to calculate  $27 \cdot 5$  og  $2.7 \cdot 5$ .
- b) Use a calculator to calculate  $247 \cdot 192$  og  $24.7 \cdot 19.2$ .
- c) Use a calculator to calculate  $928 \cdot 74$  og  $9.28 \cdot 7.4$ .
- d) Use a calculator to calculate  $134 \cdot 4\,249$  og  $1.34 \cdot 42.49$ .
- e) Compare the pairs of answers from a) - c), and make a rule for how to calculate multiplication involving decimal numbers.

### 6.3.5

Calculate.

- a)  $82.3 \cdot 5$       b)  $9.51 \cdot 7$       c)  $22.4 \cdot 1.7$

### 6.4.1

Calculate.

- a)  $98 : 2$       b)  $87 : 3$       c)  $92 : 4$       d)  $85 : 5$       e)  $72 : 6$

### 6.4.2

Calculate.

- a)  $378 : 2$       b)  $224 : 4$       c)  $495 : 5$
- e)  $133 : 7$       f)  $208 : 8$       g)  $873 : 9$

### 6.5.1

Write the number in standard form.

- a) 98 000      b) 167 000 000      c) 4 819      d) 21
- e) 9 132,27      f) 893.7      g) 18 002.1      h) 302.4

### 6.5.2

Write the number in standard form.

- a) 0.027      b) 0.0001901      c) 0.32      d) 0.00000020032

### 6.5.3

Given the calculation

$$900\,000\,000 \cdot 0.00007$$

- a) Explain why the calculation can be written as

$$9 \cdot 10^8 \cdot 7 \cdot 10^{-5}$$

- b) Use power rules (see [Section 8.2](#)) to find the product of the calculation from a).

### 6.5.4 (T1H21D1)

Calculate, and write answer in standard form.

$$\frac{6.2 \cdot 10^7 + 2.5 \cdot 10^8}{0.000002}$$

## Chapter 7

# Geometry

## 7.1 Terms


### Point

A given position is called a<sup>1</sup> *point*. We mark a point by drawing a dot, which we preferably name by a letter. Below we have drawn the points  $A$  and  $B$ .



### Line and segment

A straight dash with infinite length (!) is called a *line*. The fact that a line has infinite length, makes *drawing* a line impossible, we can only *imagine* a line. Imagining a line can be done by drawing a straight dash and think of its ends as wandering out in each direction.



A straight dash between two points is called a *segment*.



We write the segment between the points  $A$  and  $B$  as  $AB$ .


#### Notice

A segment is a part of a line, therefore a line and a segment have a lot of attributes in common. When writing about lines, it will be up to the reader to confirm whether the same applies for segments. Hence we avoid the need of writing "lines/segments".

---

<sup>1</sup>See also [Section 1.3](#).

## Segment or length?



The segments  $AB$  and  $CD$  have equal length, but they are not the same segment. Still we'll write  $AB = CD$ . That is, we'll use the same names for the line segments and their lengths (the same applies for angles and their values, see page 128-130).

We'll do this by the following reasons:

- The context will make it clear whether we are talking about a segment or a length.
- Finding it necessary to write e.g. "the length of  $AB$ " would make sentences less readable.


## Distance

There are infinite ways one can move from one point to another and some ways will be longer than others. When talking about a distance in geometry, we usually mean the *shortest* distance. For geometries studied in this book the shortest distance between two points will always equal the length of the segment (blue in the below figure) connecting them.




## Circle; center, radius and diameter

If we make an enclosed curve where all points on this curve have the same distance to a given point, we have a *circle*. The point which all the points on the curve have an equal distance to is the *center* of the circle. A segment between a point on the curve and the center is called a *radius*. A segment between two points on the curve, passing through the circle center, is called a *diameter*<sup>1</sup>.



## Arcs and sectors

A part of a circular curve is called an *arc*. The shape formed by an arc and two associated radii is called a *sector*. The below figure shows three different sectors.




---

<sup>1</sup>As mentioned, *radius* and *diameter* can just as well indicate the length of the segments.




## Parallel lines

Lines aligned in the same direction are *parallel*. The below figure shows two pairs of parallel lines.




We use the symbol  $\parallel$  to indicate that two lines are parallel.

$$AB \parallel CD$$




## Angles

Non-parallel lines will sooner or later intersect. The gap formed by two non-parallel lines is called an *angle*. We draw angles as small circular curves:




Lines creating an angle are called the *sides* of the angle. The intersection point of the lines are called the *vertex* of the angle. It is common to use the symbol  $\angle$  to underline the angle in question. In the below figure we have the following:

- the angle  $\angle BOA$  has angle sides  $OB$  and  $OA$  and vertex  $O$ .
- the angle  $\angle AOD$  has angle sides  $OA$  and  $OD$  and vertex  $O$ .




Measure of angles in degrees

When measuring an angle in degrees, we imagine a circular curve divided into 360 equally long pieces. We call one such piece 1 *degree*, indicated by the symbol  $^{\circ}$ .




Notice that an angle with measure  $90^{\circ}$  is indicated by the symbol  $\square$ . Such an angle is called a *right angle*. Lines which form right angles are said to be *perpendicular* to one another, indicated by the symbol  $\perp$ .




## What angle?

Strictly speaking, when two segments (or lines) intersect, they form two angles; the one larger than  $180^\circ$  and the other smaller than  $180^\circ$ . Usually it is the smaller angle we wish to study, therefore it is common to define  $\angle AOB$  as the *smaller* angle formed by the segments  $OA$  and  $OB$ .



As long as there are only two segments/lines present, it is common using only one letter to indicate the angle:



## 7.1 Vertical angles

Two opposite angles with a common vertex is called *vertical angles*. Vertical angles are of equal measure.



## 7.1 Vertical angles (explanation)



We have


$$\angle BOC + \angle DOB = 180^\circ$$

$$\angle AOD + \angle DOB = 180^\circ$$


Hence,  $\angle BOC = \angle AOD$ . Similarly,  $\angle COA = \angle DOB$ .

## Sides and vertices

When line segments form an enclosed shape, they form a *polygon*. The below figure shows, from left to right, a triangle (3-gon), a quadrilateral (4-gon) and a pentagon (5-gon).




The segments of a polygon are called *edges* or *sides*. The respective intersection points of the segments are the *vertices* of the polygon. That is, the triangle below has vertices  $A$ ,  $B$  and  $C$  and sides (edges)  $AB$ ,  $BC$  and  $AC$ .




### Note

Often we'll write a letter only to indicate a vertex of a polygon.




## Diagonals

Segments between two vertices not belonging to the same side of a polygon is called a *diagonal*. The below figure shows the diagonals  $AC$  and  $BD$ .




## Altitudes and base lines

In [Section 7.4](#), the terms *base* and *height* (*altitude*) play an important role. To find the height of a triangle, we choose one of the sides to be the base. In the below figure, let's start with  $AB$  as the base. Then the height is the segment from  $AB$  (potentially, as is the case here, the extension of  $AB$ ) to  $C$ , perpendicular to  $AB$ .



Since there are three sides which can be bases, a triangle has three heights.




### Notice

The terms altitude and base also applies to other polygons.

# 7.2 Attributes of triangles and quadrilaterals

In addition to having a certain number of sides and vertices, polygons have other attributes, such as sides or angles of equal measure, or parallel sides. There are specific names of polygons with special attributes, and these names can be put into an overview where some "inherit"<sup>1</sup> attributes from others.

## 7.2 Triangles



**Triangle**  
Have three sides and three vertices.



**Right triangle**  
Have an angle of  $90^\circ$ .



**Isosceles triangle**  
At least two sides are of equal length.  
At least two angles are of equal measure.



**Equilateral triangle**  
The sides are of equal length.  
Each of the angles equals  $60^\circ$ .

### Example

Since an equilateral triangle have three sides of equal length and three angles equal to  $60^\circ$ , it is also an isosceles triangle.

### The language box


The longest side of a right triangle is called the *hypotenuse* and the shorter sides are called *legs*.

<sup>1</sup>In Rule 7.2 and Rule 7.4 this is indicated by arrows.


### 7.3 The sum of angles in a triangle

In a triangle, the sum of the angles equals  $180^\circ$ .

$$\angle A + \angle B + \angle C = 180^\circ$$



### 7.3 The sum of angles in a triangle (explanation)



We draw a segment  $FG$  passing through  $C$  and parallel to  $AB$ . Moreover, we place  $E$  and  $D$  on the extension of  $AC$  and  $BC$ , respectively. Then  $\angle A = \angle GCE$  and  $\angle B = \angle DCF$ .  $\angle ACB = \angle ECD$  because they are vertical angles. Now

$$\angle DCF + \angle ECD = \angle GCE = 180^\circ$$


Hence

$$\angle CBA + \angle ACB + \angle BAC = 180^\circ$$




## 7.4 Quadrilaterals

Quadrilateral  $\longrightarrow$  Trapezoid  $\longrightarrow$  Parallelogram  $\begin{matrix} \nearrow & \text{Rhombus} \\ \searrow & \text{Rectangle} \end{matrix} \longrightarrow$  Square




### Quadrilateral

Have four sides and four vertices.




### Trapezoid

Have at least one pair of parallel sides.



### Parallelogram

Have two pairs of parallel sides.  
Have two pairs of equal angles.



### Rhombus

All sides are of equal length.



### Rectangle

All angles equals  $90^\circ$ .



### Square

### Example

The square is both a rhombus and a rectangle, which means it "inherits" their attributes. From this it follows that, in a square,

- all sides are of equal length.
- all angles equals  $90^\circ$ .

### 7.5 The sum of angles in a quadrilateral


In a quadrilateral, the sum of the angles equals  $360^\circ$ .

$$\angle A + \angle B + \angle C + \angle D = 360^\circ$$




### 7.5 The sum of angles in a quadrilateral (explanation)

The total sum of angles of  $\triangle ABD$  and  $\triangle BCD$  equals the sum of the angles in  $\square ABCD$ . By [Rule 7.3](#), the sum of angles of triangles  $180^\circ$ , therefore the sum of the angles of  $\square ABCD$  equals  $2 \cdot 180^\circ = 360^\circ$ .




# 7.3 Perimeter

When we measure the length around an enclosed shape, we find its *perimeter*. Let’s find the perimeter of this rectangle:



The rectangle has two sides of length 4 and two sides of length 5.




Hence

$$\begin{aligned} \text{The perimeter of the rectangle} &= 4 + 4 + 5 + 5 \\ &= 18 \end{aligned}$$

## 7.6 Perimeter

A perimeter is the length around a closed shape.

### Example



In figure (a) the perimeter equals  $5 + 2 + 4 = 11$ .

In figure (b) the perimeter equals  $4 + 5 + 3 + 1 + 6 + 5 = 24$ .

# 7.4 Area

Our surroundings are full of *surfaces*, for example on a floor or a sheet. When measuring surfaces, we find their *area*. The concept of area is the following:

We imagine a square with sides of length 1. We call this the *unit square*.




Then, regarding the surface for which we seek the area of, we ask:

”How many unit squares does this surface contain?”


## The area of a rectangle

Let’s find the area of a rectangle with baseline 3 and altitude 2.



Simply by counting, we find that the rectangle contains 6 unit squares:

The area of the rectangle = 6




Looking back at [Section 2.3](#), we notice that

$$\begin{aligned} \text{The area of the rectangle} &= 3 \cdot 2 \\ &= 6 \end{aligned}$$

## 7.7 The area of a rectangle

$$\text{Area} = \text{baseline} \cdot \text{altitude}$$




### Width and length

In a rectangle, the baseline and the altitude are also referred to as (in random order) the *width* and the *length*.

### Example 1

Find the area of the rectangle<sup>1</sup>.




**Answer**

$$\text{The area of the rectangle} = 4 \cdot 2 = 8$$

### Example 2

Find the area of the square.



**Answer**

$$\text{The area of the square} = 3 \cdot 3 = 9$$


*Note:* The lengths used in one figure will not necessarily correspond to the lengths in another figure. That is, a side of length 1 in one figure can might as well be shorter than a side of length 1 in another figure.

## The area of a triangle


Concerning triangles, there are three different cases to study:

1) *The baseline and the altitude have a common end point*

Let's find the area of a right triangle with baseline 5 and altitude 3.



We can make a rectangle by copying our triangle and joining the hypotenuses:



By [Rule 7.7](#), the area of the rectangle equals  $5 \cdot 3$ . The area of one of the triangles makes up half the area of the rectangle, so

$$\text{The area of the blue triangle} = \frac{5 \cdot 3}{2}$$

Regarding the blue triangle we have


$$\frac{5 \cdot 3}{2} = \frac{\text{baseline} \cdot \text{height}}{2}$$

2) The altitude is placed inside the triangle, but have no common end point with the baseline

The below triangle has baseline 5 and altitude 4.



We make a rectangle containing the blue triangle (split into the red triangle and the yellow triangle):



Observe that


- the area of the red triangle makes up half the area of the rectangle formed by the red and the yellow triangle.
- the area of the yellow triangle makes up half the area of the rectangle formed by the yellow and the green triangle.

It now follows that the sum of the areas of the yellow triangle and the red triangle makes up half the area of the rectangle formed by the four colored triangles. The area of this rectangle equals  $5 \cdot 4$ , and since our original triangle (the blue) includes the red triangle and the orange triangle, we have


$$\text{The area of the blue triangle} = \frac{5 \cdot 4}{2} = \frac{\text{baseline} \cdot \text{height}}{2}$$

2) *The altitude is placed outside the triangle*

The below triangle has baseline 4 and altitude 3.



We now make a rectangle containing the blue triangle:



Now we introduce the following names:

The area of the rectangle =  $R$

The area of the blue triangle =  $B$

The area of the orange triangle =  $O$

The area of the green triangle =  $G$

We have

$$R = 3 \cdot 10 = 30$$

$$O = \frac{3 \cdot 10}{2} = 15$$

$$G = \frac{3 \cdot 6}{2} = 9$$

Moreover,

$$\begin{aligned} B &= R - O - G \\ &= 30 - 15 - 9 \\ &= 6 \end{aligned}$$

Observe that we can write

$$6 = \frac{4 \cdot 3}{2}$$

Regarding the blue triangle we recognize this as

$$\frac{4 \cdot 3}{2} = \frac{\text{base} \cdot \text{height}}{2}$$




*The three cases summarized*

For a chosen baseline in a triangle, one of the cases discussed will always be valid. All cases resulted in the same expression for the area of the triangle.


### 7.8 The area of a triangle

$$\text{Area} = \frac{\text{base} \cdot \text{height}}{2}$$



#### Example 1

Find the area of the triangle.




**Answer**

$$\begin{aligned}\text{The area of the triangle} &= \frac{4 \cdot 3}{2} \\ &= 6\end{aligned}$$

**Example 2**

Find the area of the triangle.




**Answer**

$$\text{The area of the triangle} = \frac{6 \cdot 5}{2} = 15$$

**Example 3**

Find the area of the triangle.




**Answer**


$$\text{The area of the triangle} = \frac{7 \cdot 3}{2} = \frac{21}{2}$$

## 7.5 Tredimensjonal geometri

So far, we have studied two-dimensional shapes as triangles, quadrilaterals, circles etc. Every two-dimensional figure can be drawn in a coordinate system with two axes.




In order to draw **three-dimensional** shapes however, we need three axes:



While we have used the phrase *width/length* and *height* to describe a rectangle, we say that the box above has a width, a height *and* a length (depth).


The area lying on the "outside" of a three-dimensional shape is called the **surface**. The surface of the box above is made up by 6 rectangles. Polygons that are parts of a surface are called **lateral surfaces**.

## 7.9 Three-dimensional figures




### **Right prism**

Has two equal and four equal rectangles as lateral surfaces. All the surfaces in contact are perpendicular to each other.




### **Kube**

A right prism with squares as lateral surfaces.




### **Triangular prism**

Two of the lateral surfaces are congruent triangles parallel to each other. These triangles are connected by lateral surfaces that are rectangles.




### **Rectangular pyramid**

The surface is made up by one rectangle and four triangles.




### **Triangular pyramid**

The surface is made up by four triangles.



### **Cone**

One part of the surface is a circle, the other part is a folded sector.



### **Sphere**

All the points on the surface have equal distance to the center.


**Tip**

It is not that easy to imagine what a *folded sector* is, but try this:


Draw a sector on a sheet. Cut the sector, and join the two edges of the sector. Then you have a cone without bottom surface.

## 7.6 Volum


When we speak of how much space there is inside an object, we speak of it's **volume**. As a measure of volume, we imagine a cube with sides of length 1.



We shall call such a cube a 'unit cube'. Say we have a rectangular prism with width 3, length 4 and height 2.




Inside it, there is room for exactly 24 unit cubes.



Other than simply counting unit cubes, we can find this by calculating

$$3 \cdot 4 \cdot 2 = 24$$


### 7.10 The volume of a rectangular prism I



$$\text{volume} = \text{width} \cdot \text{length} \cdot \text{height}$$

## Base area


To calculate the volume of elementary shapes, it is useful to apply the term **base surface**. In the same way as with baselines<sup>1</sup>, it is our choice of base surface that decides what the height is. Regarding the prism on the preceding page, it is natural to choose the horizontally aligned surface as the base. To indicate the base we shall use the letter  $G$ :



The base surface has area  $3 \cdot 4 = 12$ , and the corresponding height is 2. The volume of the prism equals the base surface area multiplied by the height:

$$\begin{aligned} V &= 3 \cdot 4 \cdot 2 \\ &= G \cdot 2 \\ &= 24 \end{aligned}$$

### 7.11 Volumee of a rectangular prism II



$$\text{volume} = G \cdot \text{height}$$

### Grunnflata eller grunnflatearealet?


In the above text, we first let  $G$  denote the base surface, and later we let  $G$  denote the *base surface area*. In this book the term *base surface* is so strongly connected to the *base surface area* that we will let the same symbol denote them both, leaving the reader to differ them from the context of use.

<sup>1</sup>See page ??.


## Exercises for Chapter 7

### 7.1.1


Find the perimeter of the colored figure.



(a)




(b)




(c)

### 7.1.2


Calculate the area of the rectangles.



(a)



(b)



(c)

### 7.1.3

Find the width and height of the rectangle, given that

- a) the area is 16 and the perimeter is 20.
- b) the area is 12 and the perimeter is 14.
- c) the area is 18 and the perimeter is 18.




### 7.1.4

- a) Find the area of a square with perimeter 16.
- b) Give three examples of rectangles with perimeter 36. Provide the answer by stating the width, height and area of the rectangle.



### 7.1.5

Find the area of the triangle.




### 7.1.6

A prism has length 9, width 10 and height 8.

- Find the base area of the prism.
- Find the volume of the prism.

### 7.2.1




Shift the triangle with the vector shown.

a)




b)




### 7.2.2

Mirror the triangle about the dashed line.


a)



b)




c)




### 7.2.3

Find the angle and point of rotation.

a)



b)



### Gruble 6


- a) Explain why the perimeter of a rectangle is always an even number.
- b) "If both the width and height of a rectangle is an odd number, there is no way the area and the perimeter of the rectangle can have the same value."

Explain why the proposition is correct/incorrect.

- c) What is the side length of the only square with area and perimeter of equal value?

### Gruble 7

$\square ABCD$  is a square and  $\triangle DEC$  is equilateral. Find the value of  $v$ .



## Part II

# Algebra and geometry

## Chapter 8

# Algebra

# 8.1 Introduction

Simply said, **algebra** is mathematics where letters represent numbers. This makes it easier working with *general* cases. For example,  $3 \cdot 2 = 2 \cdot 3$  and  $6 \cdot 7 = 7 \cdot 6$  but these are only two of the infinitely many examples of the commutative property of multiplication! One of the aims of algebra is giving *one* example that explains *all* cases, and since our digits (0-9) are inevitably connected to specific numbers, we apply letters to reach this target.

The value of the numbers represented by letters will often vary, in that case we call the letter-numbers *variables*. If letter-numbers on the other hand have a specific value, they are called *constants*.

In [Part I](#), we studied calculations through examples with specific numbers, however, most of these rules are *general*; they are valid for all numbers. On page 159-162, many of these rules are reproduced in a general form. A good way of getting acquainted with algebra is comparing the rules here presented by the way they are expressed in<sup>1</sup> [Part I](#).

## 8.1 Addition is commutative (2.1)

$$a + b = b + a$$

### Example

$$7 + 5 = 5 + 7$$

## 8.2 Multiplication is commutative (2.2)

$$a \cdot b = b \cdot a$$

### Example 1

$$9 \cdot 8 = 8 \cdot 9$$

### Example 2

$$8 \cdot a = a \cdot 8$$

---

<sup>1</sup>The number of the rules as found in [Part I](#) are written inside parentheses.



## Multiplication involving letters

When multiplication involves letters, it is common to omit the symbol of multiplication. If a specific number and a letter are multiplied together, the specific number is written first. For example,

$$a \cdot b = ab$$

and

$$a \cdot 8 = 8a$$

We also write

$$1 \cdot a = a$$

In addition, it is common to omit the symbol of multiplication when an expression with parentheses is involved:

$$3 \cdot (a + b) = 3(a + b)$$

## 8.3 Fractions as rewriting of division (4.1)

$$a : b = \frac{a}{b}$$

### Example

$$a : 2 = \frac{a}{2}$$

## 8.4 Fractions multiplied by fractions (4.8)

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$$

### Example 1

$$\frac{2}{11} \cdot \frac{13}{21} = \frac{2 \cdot 13}{11 \cdot 21} = \frac{26}{231}$$

### Example 2

$$\frac{3}{b} \cdot \frac{a}{7} = \frac{3a}{7b}$$

### 8.5 Division by fractions (4.10)

$$\frac{a}{b} : \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c}$$

#### Example 1

$$\frac{1}{2} : \frac{5}{7} = \frac{1}{2} \cdot \frac{7}{5}$$

#### Example 2

$$\begin{aligned}\frac{a}{13} : \frac{b}{3} &= \frac{a}{13} \cdot \frac{3}{b} \\ &= \frac{3a}{13b}\end{aligned}$$

### 8.6 Distributive law (3.2)

$$(a + b)c = ac + bc$$

#### Example 1

$$(2 + a)b = 2b + ab$$

#### Example 2

$$a(5b - 3) = 5ab - 3a$$

### 8.7 Multiplication by negative numbers I (5.6)

$$a \cdot (-b) = -(a \cdot b)$$

#### Example 1

$$\begin{aligned}3 \cdot (-4) &= -(3 \cdot 4) \\ &= -12\end{aligned}$$

**Example 2**

$$\begin{aligned}(-a) \cdot 7 &= -(a \cdot 7) \\ &= -7a\end{aligned}$$

**8.8 Multiplication by negative numbers II (5.7)**

$$(-a) \cdot (-b) = a \cdot b$$

**Example 1**

$$\begin{aligned}(-2) \cdot (-8) &= 2 \cdot 8 \\ &= 16\end{aligned}$$

**Example 2**

$$(-a) \cdot (-15) = 15a$$

## Extensions of the rules

One of the strengths of algebra is that we can express compact rules which are easily extended to apply for other cases. Let's, as an example, find another expression of

$$(a + b + c)d$$

Rule 8.6 does not directly imply how to calculate between the expression inside the parentheses and  $d$ , but there is no wrongdoing in defining  $a + b$  as  $k$ :

$$a + b = k$$

Then

$$(a + b + c)d = (k + c)d$$

Now, by Rule 8.6, we have

$$(k + c)d = kd + cd$$

Inserting the expression for  $k$ , we have

$$kd + cd = (a + b)d + cd$$

By applying Rule 8.6 once more we can write

$$(a + b)d + cd = ad + bc + cd$$

Then

$$(a + b + c)d = ad + bc + cd$$

*Notice! This example is not meant to show how to handle expressions not directly covered by Rule 8.1 - Rule 8.8, but to emphasize why it's always sufficient to write rules with the least amount of terms, factors etc. Usually you apply extension of the rules without even thinking about it, and surely not in such meticulous manner as here provided.*

## 8.2 Powers

$$\text{base} \rightarrow 2^3 \leftarrow \text{exponent}$$

A power is composed by a *base* and an *exponent*. For example,  $2^3$  is a power with base 2 and exponent 3. An exponent which is a positive integer indicates the amount of instances of the base to be multiplied together, that is

$$2^3 = 2 \cdot 2 \cdot 2$$

### 8.9 Powers

$a^n$  is a power with base  $a$  and exponent  $n$ .

If  $n$  is a natural number,  $a^n$  corresponds to  $n$  instances of  $a$  multiplied together.

*Note:*  $a^1 = a$

#### Example 1

$$\begin{aligned} 5^3 &= 5 \cdot 5 \cdot 5 \\ &= 125 \end{aligned}$$

#### Example 2

$$c^4 = c \cdot c \cdot c \cdot c$$

#### Example 3

$$\begin{aligned} (-7)^2 &= (-7) \cdot (-7) \\ &= 49 \end{aligned}$$

## The language box

Common ways of saying  $2^3$  are<sup>1</sup>

- "2 (raised) to the power of 3"
- "2 to the third power"

In computer programming, the symbol `^` or the symbols `**` is usually written between the base and the exponent.

Raising a number to the power of 2 is called "squaring" the number.

---

<sup>1</sup>Attention! The examples illustrate a paradox in the English language; *power* is also a synonym for *exponent*.

## Note

The next pages declares rules concerning powers with corresponding explanations. Even though one wish to have these explanations as general as possible, we choose to use, mostly, specific numbers as exponents . Using variables as exponents would lead to less reader-friendly expressions, and it is our claim that the general cases are well illustrated by the specific cases.

## 8.10 Multiplication by powers

$$a^m \cdot a^n = a^{m+n} \quad (8.1)$$

### Example 1

$$\begin{aligned} 3^5 \cdot 3^2 &= 3^{5+2} \\ &= 3^7 \end{aligned}$$

### Example 2

$$\begin{aligned} b^4 \cdot b^{11} &= b^{3+11} \\ &= b^{14} \end{aligned}$$

**Example 3**

$$\begin{aligned}
 a^5 \cdot a^{-7} &= a^{5+(-7)} \\
 &= a^{5-7} \\
 &= a^{-2}
 \end{aligned}$$

(See [Rule 8.13](#) regarding how powers with negative exponents can be interpreted.)

**8.10 Multiplication by powers (explanation)**

Let's study the case

$$a^2 \cdot a^3$$

We have

$$a^2 = 2 \cdot 2$$

$$a^3 = 2 \cdot 2 \cdot 2$$

Hence we can write

$$\begin{aligned}
 a^2 \cdot a^3 &= \overbrace{a \cdot a}^{a^2} \cdot \overbrace{a \cdot a \cdot a}^{a^3} \\
 &= a^5
 \end{aligned}$$

**8.11 Division by powers**

$$\frac{a^m}{a^n} = a^{m-n}$$

**Example 1**

$$\frac{3^5}{3^2} = 3^{5-2} = 3^3$$

**Example 2**

$$\begin{aligned}
 \frac{2^4 \cdot a^7}{a^6 \cdot 2^2} &= 2^{4-2} \cdot a^{7-6} \\
 &= 2^2 a \\
 &= 4a
 \end{aligned}$$

### 8.11 Division by powers (explanation)

Let's examine the fraction  $\frac{a^5}{a^2}$ . Expanding the powers, we get

$$\begin{aligned}\frac{a^5}{a^2} &= \frac{a \cdot a \cdot a \cdot a \cdot a}{a \cdot a} \\ &= \frac{\cancel{a} \cdot \cancel{a} \cdot a \cdot a \cdot a}{\cancel{a} \cdot \cancel{a}} \\ &= a \cdot a \cdot a \\ &= a^3\end{aligned}$$

The above calculations are equivalent to writing

$$\begin{aligned}\frac{a^5}{a^2} &= a^{5-2} \\ &= a^3\end{aligned}$$

### 8.12 The special case of $a^0$

$$a^0 = 1$$

#### Example 1

$$1000^0 = 1$$

#### Example 2

$$(-b)^0 = 1$$

### 8.12 The special case of $a^0$ (explanation)

A number divided by itself always equals 1, therefore

$$\frac{a^n}{a^n} = 1$$

From this, and [Rule 8.11](#), it follows that

$$\begin{aligned}1 &= \frac{a^n}{a^n} \\ &= a^{n-n} \\ &= a^0\end{aligned}$$



### 8.13 Powers with negative exponents

$$a^{-n} = \frac{1}{a^n}$$

#### Example 1

$$a^{-8} = \frac{1}{a^8}$$

#### Example 2

$$(-4)^{-3} = \frac{1}{(-4)^3} = -\frac{1}{64}$$

### 8.13 Powers with negative exponents (explanation)

By [Rule 8.12](#), we have  $a^0 = 1$ . Thus

$$\frac{1}{a^n} = \frac{a^0}{a^n}$$

By [Rule 8.11](#), we obtain

$$\begin{aligned}\frac{a^0}{a^n} &= a^{0-n} \\ &= a^{-n}\end{aligned}$$

### 8.14 Fractions as base

$$\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m} \quad (8.2)$$

#### Example 1

$$\left(\frac{3}{4}\right)^2 = \frac{3^2}{4^2} = \frac{9}{16}$$

#### Example 2

$$\left(\frac{a}{7}\right)^3 = \frac{a^3}{7^3} = \frac{a^3}{343}$$

### 8.14 Fractions as base (explanation)

Let's study

$$\left(\frac{a}{b}\right)^3$$

We have

$$\begin{aligned}\left(\frac{a}{b}\right)^3 &= \frac{a}{b} \cdot \frac{a}{b} \cdot \frac{a}{b} \\ &= \frac{a \cdot a \cdot a}{b \cdot b \cdot b} \\ &= \frac{a^3}{b^3}\end{aligned}$$

### 8.15 Factors as base

$$(ab)^m = a^m b^m \quad (8.3)$$

**Example 1**

$$\begin{aligned}(3a)^5 &= 3^5 a^5 \\ &= 243a^5\end{aligned}$$

**Example 2**

$$(ab)^4 = a^4 b^4$$

### 8.15 Factors as base (explanation)

Let's use  $(a \cdot b)^3$  as an example. We have

$$\begin{aligned}(a \cdot b)^3 &= (a \cdot b) \cdot (a \cdot b) \cdot (a \cdot b) \\ &= a \cdot a \cdot a \cdot b \cdot b \cdot b \\ &= a^3 b^3\end{aligned}$$

## 8.16 Powers as base

$$(a^m)^n = a^{m \cdot n} \quad (8.4)$$

### Example 1

$$\begin{aligned} (c^4)^5 &= c^{4 \cdot 5} \\ &= c^{20} \end{aligned}$$

### Example 2

$$\begin{aligned} \left(3^{\frac{5}{4}}\right)^8 &= 3^{\frac{5}{4} \cdot 8} \\ &= 3^{10} \end{aligned}$$

## 8.16 Powers as base (explanation)

Let's use  $(a^3)^4$  as an example. We have

$$(a^3)^4 = a^3 \cdot a^3 \cdot a^3 \cdot a^3$$

By [Rule 8.10](#), we get

$$\begin{aligned} a^3 \cdot a^3 \cdot a^3 \cdot a^3 &= a^{3+3+3+3} \\ &= a^{3 \cdot 4} \\ &= a^{12} \end{aligned}$$

### 8.17 $n$ -root

$$a^{\frac{1}{n}} = \sqrt[n]{a}$$

The symbol  $\sqrt{\phantom{x}}$  is called the *radical sign*. In the case of an exponent equal to  $\frac{1}{2}$ , it is common to omit 2 from the radical:

$$a^{\frac{1}{2}} = \sqrt{a}$$

#### Example

By [Rule 8.16](#), we have

$$\begin{aligned}\left(a^b\right)^{\frac{1}{b}} &= a^{b \cdot \frac{1}{b}} \\ &= a\end{aligned}$$

For example is

$$9^{\frac{1}{2}} = \sqrt{9} = 3, \text{ since } 3^2 = 9$$

$$125^{\frac{1}{3}} = \sqrt[3]{125} = 5, \text{ since } 5^3 = 125$$

$$16^{\frac{1}{4}} = \sqrt[4]{16} = 2, \text{ since } 2^4 = 16$$

#### The language box

$\sqrt{9}$  is called "the square (the 2nd) root of 9"

$\sqrt[3]{8}$  is called "the cube (the 3th) root of 8"

## 8.3 Irrational numbers

### 8.18 Irrational numbers

A number which is *not* a rational number, is an irrational number<sup>1</sup>.

The value of an irrational number are decimal numbers with infinite digits in a non-repeating manner.

#### Example 1

$\sqrt{2}$  is an irrational number.

$$\sqrt{2} = 1.414213562373\dots$$

---

<sup>1</sup>Strictly speaking, irrational numbers are all *real* numbers which are not rational numbers. But to explain what *real* numbers are, we have to mention *imaginary* numbers, and this we choose not to do in this book.

## Exercises for Chapter 8

### 8.1.1

Exploit the relation between repeated addition and multiplication (see [Rule 2.3](#) and [Rule 5.5](#)) to simplify the expressions.

a)  $a + a + a$       b)  $a + a + a + a$       c)  $a + a + a + a + a + a + a$

d)  $-b - b$       e)  $-b - b - b - b - b$       f)  $-k - k - k$

### 8.1.2

Simplify the expressions.

a)  $2a + b - a$       b)  $-4a + 2b + 3a$       c)  $7b - 3a + 2b$

### 8.1.3

Simplify the expressions.

a)  $4c + 2b - 5a - 3c$       b)  $-9a - 3c + 3b + 3c$       c)  $9b - 3a + 2b$

### 8.1.4

Use [Rule 3.2](#) to write the expression without parentheses.

a)  $7(a + 2)$       b)  $9(b + 3)$       c)  $8(b - 3c)$       d)  $(-2)(3a + 5b)$

e)  $(9a + 2)$       f)  $(3b + 8)a$       g)  $(b - 3c)(-a)$

h)  $2(a + 3b + 4c)$       i)  $9(3b - c + 7a)$       j)  $(3b - c + 7a)(-2)$

### 8.1.5

Use [Rule 3.2](#) to factorize the expression.

a)  $2a + 2b$       b)  $4ab + 5b$       c)  $9bc - c$       d)  $4ac - 2a$

### 8.1.6

Prove that

a)  $(a + b)^2 = a^2 + 2ab + b^2$

b)  $(a - b)^2 = a^2 - 2ab + b^2$

c)  $(a + b)(a - b) = a^2 - b^2$

### 8.1.7 (GV21D1)

a) Simplify the expression.

$$\frac{a + a + a + a}{4a}$$

b) What value will the expression  $\frac{y^2 - 2y}{y^2}$  attain if  $x = 4$  and  $y = -2$ ?

### 8.1.8 (E22)

Given the expression  $(a + b)^2 = 16$ . Decide whether the below alternatives makes the expression valid.

- $a = 2$  and  $b = 2$
- $a = 8$  and  $b = 4$
- $a = 8$  and  $b = -4$

### 8.2.1

Write as a power.

a)  $3 \cdot 3 \cdot 3 \cdot 3$       b)  $5 \cdot 5$       c)  $7 \cdot 7 \cdot 7 \cdot 7 \cdot 7 \cdot 7$

d)  $a \cdot a \cdot a$       e)  $b \cdot b$       f)  $(-c)(-c)(-c)(-c)$

### 8.2.2

Find the value of the power.

a)  $8^2$       b)  $2^5$       c)  $4^3$       d)  $(-2)^3$       e)  $(-3)^5$       f)  $(-4)^4$

### 8.2.3

Write the expression as a power.

a)  $2^7 \cdot 2^9$       b)  $3^4 \cdot 3^7$       c)  $9 \cdot 9^5$       d)  $6^8 \cdot 6^{-3}$       e)  $5^3 \cdot 5^{-7}$

f)  $10^8 \cdot 10^{-3} \cdot 10^6$       g)  $a^9 \cdot a^7$       h)  $k^5 \cdot k^2$       i)  $x^5 \cdot x^{-2}$

k)  $x^{-4} \cdot x^5$       l)  $a^{-5} \cdot a \cdot a^4$       m)  $a^3 \cdot b^5 \cdot a^2 \cdot b^{-8}$

### 8.2.4

Calculate..

a)  $\sqrt{25}$       b)  $\sqrt{100}$       c)  $\sqrt{144}$

d)  $\sqrt[3]{27}$       e)  $\sqrt[3]{729}$       f)  $\sqrt[5]{100000}$



### Gruble 8

(1TH21D1)

Simplify the expression

$$\frac{9^{\frac{1}{2}} \cdot 3^{-1} + 9^0}{8^{\frac{3}{4}}}$$

### Gruble 9

By adding the digits of a number, we find the **digit sum** of the number. For example is the digit sum of 14 equal to  $1 + 4 = 5$ , and the digit sum of 918 equals  $9 + 1 + 8 = 18$ . Prove that if the digit sum of a 3-digit integer is divisible by 3, then the number is also divisible by 3.

*Note:* Generalizing the 3-digit case is quite easy, thus proving that the rule is valid for an integer with any number of digits.

## Comment (for the particularly interested)

Mathematics is *axiomatically* founded. This means we declare<sup>1</sup> some propositions to be true, and these are called *axioms* or *postulates*. For the subject of calculations we have 12 axioms<sup>2</sup>, but in this book we have confined ourselves to explicitly mention the following 6:

### Axioms

For the numbers  $a$ ,  $b$  and  $c$  we have

$$a + (b + c) = (a + b) + c \quad (\text{A1})$$

$$a + b = b + a \quad (\text{A2})$$

$$a(bc) = (ab)c \quad (\text{A3})$$

$$ab = ba \quad (\text{A4})$$

$$a(b + c) = ab + ac \quad (\text{A5})$$

$$a \cdot \frac{1}{a} = 1 \quad (a \neq 0) \quad (\text{A6})$$

- 
- (A1) Associative law for addition
  - (A2) Commutative law for addition
  - (A3) Associative law for multiplication
  - (A4) Commutative law for multiplication
  - (A5) Distributive law
  - (A6) Existence of a multiplicative identity

By applying axioms, we can derive more complex contexts which we call *theorems*. In this book we chose to let *rules* be the collective name for definitions, theorems and axioms. This is because alle three, in practice, draws up guidelines (rules) inside the mathematical system in which we wish to operate.

---

<sup>1</sup>Preferably, as few as possible.

<sup>2</sup>The number can slightly vary, depending on how the axioms are expressed.

In [Part I](#) we have tried to present the *motivation* behind the axioms, because (obviously) they are not randomly selected. The train of thoughts that leads us to them is the following:

1. Vi define positive numbers as representations of either an amount or a placement on a number line.
2. We define what addition, subtraction, multiplication and division entail for positive integers (and 0).
3. From the marks above, it's as good as self-evident that (A1)-(A6) is valid for all positive integers.
4. We define also fractions as representations of either an amount or a placement on a number line. What the elementary operations entail for fractions rests upon what is valid for the positive integers.
5. From the remarks above, we conclude that (A1)-(A6) is valid for all rational numbers.
6. We introduce negative numbers and an extended interpretation of addition and subtraction. This in turn leads to the interpretations of multiplication and division involving negative numbers.
7. (A1)-(A6) is still valid after the introduction of negative integers. Deriving that they are also valid for negative rational numbers is a formality (omitted in the book).
8. We can never write the value of an irrational number exact, but it can be approximated by a rational number<sup>1</sup>. Therefore, all calculations involving irrational numbers is, in practice, calculations involving rational numbers, and in this way we can conclude that<sup>2</sup> (A1)-(A6) is also valid for irrational numbers.

A similar train of thoughts can be applied concerning the power-rules found in [Section 8.2](#).

---

<sup>1</sup>For example, we can write  $\sqrt{2} = 1.414213562373... \approx \frac{1414213562373}{1000000000000}$

<sup>2</sup>*Attention!* This explanation is good enough for the aim of this book but is a rather extreme simplification. Irrational numbers are a very complex subject, in fact, many books presenting advanced mathematics utilize several chapters to cover the subject in full depth .

## Chapter 9

# Equations

## 9.1 Introduction

Even though every mathematical expression involving  $=$  is an *equation*, the word is, traditionally, closely linked to the presence of an *unknown* number.

Say we want to find the number which when added by 4 results in 7. The name of this unknown number is free to choose but commonly it's called  $x$ . Our equation can now be written as

$$x + 4 = 7$$

The  $x$ -value<sup>1</sup> which results in the same values on each side of the equal sign is the *solution* of the equation. It is nothing wrong done by simply observing what the value of  $x$  must be. Probably you have already realized that  $x = 3$  is the solution of the equation, since

$$3 + 4 = 7$$

However, most equations are difficult to solve simply by observing, and it is therefore vice to take the advantage of more general methods. In reality, there is only one principle to follow:

We can always carry out a mathematical operation on one of the sides of the equal sign, as long as we carry out the operation on the other side too.

The mathematical operations presented in this book is the four elementary operations. Concerning these the principal sounds:

We can always add, subtract, multiply or divide by a number on one side of the equal sign, as long as we also do it on the other side.


The principle follows from the meaning of  $=$ . When two expressions are of equal value, their values are necessarily still equal as long as we carry out identical mathematical operations on them. Anyways, in the coming section we'll specify this principle for every single elementary operation. If you already feel things make sense, you can, without no great loss of insight, skip to [Section 9.3](#).

---

<sup>1</sup>In other cases it can be several values.

# 9.2 Solving with the elementary operations

In the figures of this section we'll understand equations from what we call the weight principle. In that case,  $=$  indicates<sup>1</sup> there is equal weight (equal value) on the left and the right side.




## Addition and subtraction; moving terms

### First example


We have already found the solution of this equation, but let's now solve it in a different way<sup>2</sup>:

$$x + 4 = 7$$




The value of  $x$  becomes clear if  $x$  is alone on one of the sides, and we can isolate  $x$  on the left side by removing 4. But if we are to remove 4 from the left side, we must also remove 4 from the right side, in order to preserve equal values on both sides.

$$x + 4 - 4 = 7 - 4$$



Since  $4 - 4 = 0$  and  $7 - 4 = 3$ , we get

$$x = 3$$



<sup>1</sup>  $\neq$  symbols "not equal".

<sup>2</sup>Note: In earlier figures, there have been a correspondence between the size of the boxes and the (absolute) value of the number they represent. This does not apply to the boxes representing  $x$ .

In a more abbreviated way this can be written as


$$\begin{aligned}x + 4 &= 7 \\x &= 7 - 4 \\x &= 3\end{aligned}$$

Between the first and second line it is common to say that *4 has shifted side and therefore also sign (from + to -)*.

## Second example


Let's move on to a somehow more complex equation<sup>1</sup>:

$$4x - 2 = 3x + 5$$




To get an expression with  $x$  exclusively on one side, we remove  $3x$  on both sides:

$$4x - 2 - 3x = 3x + 5 - 3x$$




Now,

$$x - 2 = 5$$




To isolate  $x$  we add 2 on the left side. Then we must also add 2 on the right side:

$$x - 2 + 2 = 5 + 2$$



<sup>1</sup>Notice that the figure illustrates  $4x + (-2)$  (see [Section 5.3](#)) on the left side. However,  $4x + (-2)$  equals  $4x - 2$  (see [Section 5.2](#)).

Hence

$$x = 7$$


The steps we have made can be summarized in this way:

$4x - 2 = 3x + 5$	1. figure
$4x - \textcolor{red}{3x} - 2 = 3x - \textcolor{red}{3x} + 5$	2. figure
$x - 2 = 5$	3. figure
$x - 2 + \textcolor{blue}{2} = 5 + \textcolor{blue}{2}$	4. figure
$x = 7$	5. figure

In a more abbreviated way we can write

$$\begin{aligned}
 4x - 2 &= 3x + 5 \\
 4x - \textcolor{red}{3x} &= 5 + 2 \\
 x &= 7
 \end{aligned}$$

## 9.1 Moving numbers across the equal sign

To solve an equation, we gather all  $x$ -terms and all known terms on respective sides of the equal sign. A term which shifts side, also shifts sign.

### Example 1

Solve the equation

$$3x + 5 = 2x + 9$$

**Answer**

$$\begin{aligned}
 3x - 2x &= 9 - 5 \\
 x &= 4
 \end{aligned}$$

### Example 2

Solve the equation

$$-4x - 3 = -5x + 12$$

**Answer**

$$\begin{aligned}
 -4x + 5x &= 12 + 3 \\
 x &= 15
 \end{aligned}$$



# Multiplication and division

## Division

So far we have studied equations resulting in a single instance of  $x$  on one side of the equal sign. Often there are several instances of  $x$ , as, for example, in the equation

$$3x = 6$$

$x$

$x$

$x$

=

If we separate the left side into three equal groups, we get a single  $x$  in each group. And by separating the right side into three equal groups, all groups present are of equal value

$$\frac{3x}{3} = \frac{6}{3}$$

$x$

$\vdots$

$x$

$\vdots$

$x$

=

$\vdots$

$\vdots$

$\vdots$

Therefore

$$x = 2$$

$x$

=

Let's summarize our calculations:

- $3x = 6$

1. figure
- $\frac{3x}{3} = \frac{6}{3}$

2. figure
- $x = 2$

3. figure

## 9.2 Division on both sides of an equation

We can divide both sides of an equation by the same number.

### Example 1

Solve the equation

$$4x = 20$$

**Answer**

$$\begin{array}{r} \cancel{4}x = \frac{20}{\cancel{4}} \\ x = 5 \end{array}$$

### Example 2

Solve the equation

$$2x + 6 = 3x - 2$$

**Answer**

$$\begin{array}{r} 2x - 3x = -2 - 6 \\ -x = -8 \\ \cancel{-1}x = \frac{-8}{\cancel{-1}} \quad (-x = -1x) \\ x = 8 \end{array}$$

## Multiplication

Let's solve the equation

$$\frac{x}{3} = 4$$

$$\boxed{\frac{x}{3}} = \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array}$$

We can get a unit  $x$  on the left side if we add two more instances of  $\frac{x}{3}$ . The equation informs that  $\frac{x}{3}$  equals 4, this implies that for every instance of  $\frac{x}{3}$  we add to the left side, we must add 4 to the right side, in order to keep the balance.


$$\frac{x}{3} + \frac{x}{3} + \frac{x}{3} = 4 + 4 + 4$$

$$\boxed{\frac{x}{3}} + \boxed{\frac{x}{3}} + \boxed{\frac{x}{3}} = \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array} + \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array} + \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array}$$

Now we notice that  $\frac{x}{3} + \frac{x}{3} + \frac{x}{3} = \frac{x}{3} \cdot 3$  and that  $4 + 4 + 4 = 4 \cdot 3$ :

$$\frac{x}{3} \cdot 3 = 4 \cdot 3$$


$\frac{x}{3}$	$\frac{x}{3}$	$\frac{x}{3}$
---------------	---------------	---------------

=


Since  $\frac{x}{3} \cdot 3 = x$  and  $4 \cdot 3 = 12$ , we have

$$x = 12$$

$x$
-----

=


Our steps can be summarized in the following way:

- $$\frac{x}{3} = 4$$

$$\frac{x}{3} + \frac{x}{3} + \frac{x}{3} = 4 + 4 + 4$$

$$\frac{x}{3} \cdot 3 = 4 \cdot 3$$

$$x = 12$$

1. figure

2. figure

3. figure

4. figure

In a more abbreviated form this can be written as

$$\frac{x}{3} = 4$$

$$\frac{x}{\cancel{3}} \cdot \cancel{3} = 4 \cdot 3$$

$$x = 12$$

### 9.3 Multiplication on both sides of an equation

We can multiply both sides of an equation by the same number.

#### Example 1

Solve the equation

$$\frac{x}{5} = 2$$

**Answer**

$$\begin{aligned}\frac{x}{\cancel{5}} \cdot \cancel{5} &= 2 \cdot 5 \\ x &= 10\end{aligned}$$

#### Example 2

Solve the equation

$$\frac{7x}{10} - 5 = 13 + \frac{x}{10}$$

**Answer**

$$\frac{7x}{10} - \frac{x}{10} = 13 + 5$$

$$\frac{6x}{10} = 18$$

$$\frac{6x}{\cancel{10}} \cdot \cancel{10} = 18 \cdot 10$$

$$6x = 180$$

$$\frac{\cancel{6}x}{\cancel{6}} = \frac{180}{6}$$

$$x = 30$$

## 9.3 Solving with elementary operations summarized

### 9.4 Solving methods with elementary operations

We can always

- add or subtract both sides of an equation by the same number. This is equivalent to shifting a term from one side to the other, also shifting the terms sign.
- multiply or divide both sides of an equation by the same number.

#### Example 1

Solve the equation

$$3x - 4 = 6 + 2x$$

**Answer**

$$\begin{aligned} 3x - 2x &= 6 + 4 \\ x &= 10 \end{aligned}$$

#### Example 2

Solve the equation

$$9 - 7x = -8x + 3$$

**Answer**

$$\begin{aligned} 8x - 7x &= 3 - 9 \\ x &= -6 \end{aligned}$$

### Example 3

Solve the equation

$$10x - 20 = 7x - 5$$

**Answer**

$$10x - 7x = 20 - 5$$

$$3x = 15$$

$$\frac{\cancel{3}x}{\cancel{3}} = \frac{15}{3}$$

$$x = 5$$

### Example 4

Solve the equation

$$15 - 4x = x + 5$$

**Answer**

$$15 - 5 = x + 4x$$

$$10 = 5x$$

$$\frac{10}{5} = \frac{\cancel{5}x}{\cancel{5}}$$

$$2 = x$$

*Note:* In the other examples, we have chosen to collect the instances of  $x$  on the left side of the equation but we can just as well collect them on the right side. By doing that here, we have avoided calculations with negative numbers.

### Example 5

Solve the equation

$$\frac{4x}{9} - 20 = 8 - \frac{3x}{9}$$

**Answer**

$$\frac{4x}{9} + \frac{3x}{9} = 20 + 8$$

$$\frac{\cancel{7}x}{9 \cdot \cancel{7}} = \frac{28}{7}$$

$$\frac{x}{9} \cdot \cancel{9} = 4 \cdot 9$$

$$x = 36$$

### Example 6

Solve the equation

$$\frac{1}{3}x + \frac{1}{6} = \frac{5}{12}x + 2$$

#### Answer

To avoid fractions, we multiply both sides by the common denominator 12:

$$\left(\frac{1}{3}x + \frac{1}{6}\right) 12 = \left(\frac{5}{12}x + 2\right) 12 \quad (9.1)$$

$$\frac{1}{3}x \cdot 12 + \frac{1}{6} \cdot 12 = \frac{5}{12}x \cdot 12 + 2 \cdot 12 \quad (*)$$

$$4x + 2 = 5x + 24 \quad (9.2)$$

$$4x - 5x = 24 - 2 \quad (9.3)$$

$$-x = 22 \quad (9.4)$$

$$\frac{\cancel{1}x}{\cancel{1}} = \frac{22}{-1} \quad (9.5)$$

$$x = -22 \quad (9.6)$$

### Tip

Some like to make the rule that "we can multiply or divide all terms by the same number". In that case, we could have jumped to the second line in the calculations of the example above.

### Example 7

Solve the equation

$$3 - \frac{6}{x} = 2 + \frac{5}{2x}$$

#### Answer

We multiply both sides by the common denominator  $2x$ :

$$2x \left(3 - \frac{6}{x}\right) = 2x \left(2 + \frac{5}{2x}\right)$$

$$6x - 12 = 4x + 5$$

$$6x - 4x = 5 + 12$$

$$2x = 17$$

$$x = \frac{17}{2}$$

## 9.4 Power equations

Let's solve the equation

$$x^2 = 9$$

This is called a *power equation*. In general, power equations are difficult to solve applying the four elementary operations only. Applying power-rules, we raise both sides to the power of the inverse of the exponent associated with  $x$ :

$$\left(x^2\right)^{\frac{1}{2}} = 9^{\frac{1}{2}}$$

By [Rule 8.16](#), we have

$$\begin{aligned}x^{2 \cdot \frac{1}{2}} &= 9^{\frac{1}{2}} \\x &= 9^{\frac{1}{2}}\end{aligned}$$

Since  $3^2 = 9$ , we have  $9^{\frac{1}{2}} = 3$ . Now observe this:

*The principle stated on page 180 says we can, like we just did, carry out a mathematical operation on both sides of an equation. However, sticking to this principle does not guarantee that all solutions are found.*

Concerning our equation, we know that  $x = 3$  is a solution. For the sake of it, we can confirm this by the calculation

$$3^2 = 3 \cdot 3 = 9$$

But we also have

$$(-3)^2 = (-3)(-3) = 9$$

Hence,  $-3$  is also a solution of our original equation!

### 9.5 Power equations

An equation which can be written as

$$x^a = b$$

where  $a$  and  $b$  are constants, is a *power equation*.

The equation has  $a$  distinct solutions.



### Example 1

Solve the equation

$$x^2 + 5 = 21$$

**Answer**

$$x^2 + 5 = 21$$

$$x^2 = 21 - 5$$

$$x^2 = 16$$

Since  $4 \cdot 4 = 16$  and  $(-4) \cdot (-4) = 16$ , we have

$$x = 4 \quad \vee \quad x = -4$$

### Example 2

Solve the equation

$$3x^2 + 1 = 7$$

**Answer**

$$3x^2 = 7 - 1$$

$$3x^2 = 6$$

$$\frac{3x^2}{3} = \frac{6}{3}$$

$$x^2 = 2$$

Hence,

$$x = \sqrt{2} \quad \vee \quad x = -\sqrt{2}$$

### Note

Although the equation

$$x^a = b$$

has  $a$  solutions, they are not necessarily all *real*<sup>1</sup>. Concerning this book, it means we settle with finding all rational or irrational numbers which solves the equation. For example,

$$x^3 = 8$$

has 3 solutions, but we settle with the solution  $x = 2$ .

---

<sup>1</sup>As earlier mentioned, *real* and *imaginary* numbers lie outside the scope of this book.

## Exercises for Chapter 9

### 9.2.1

Solve the equation.

- a)  $x + 8 = 18$       b)  $x - 3 = 2$       c)  $x - 8 = 1$   
d)  $x + 12 = 14$       e)  $x - 1 = 2$       f)  $x - 3 = 1$   
g)  $21 = x + 11$       h)  $24 = x + 16$       i)  $4 = x - 6$

### 9.2.2

Solve the equation.

- a)  $16x - 20 = 15x + 17$       b)  $18x - 11 = 17x + 18$   
c)  $17x - 15 = 16x + 8$       d)  $4x - 9 = 6 + 3x$   
e)  $12x - 6 = 11x + 2$       f)  $2x + 10 = 3x - 1$   
g)  $5 + 8x = 9x - 18$       h)  $15 + 2x = 3x - 4$   
i)  $9x + 8 = 10x - 2$       j)  $17x + 9 = 18x - 19$

### 9.2.3

Solve the equation.

- a)  $3x = 12$       b)  $10x = 50$       c)  $7x = 63$       d)  $2x = 30$

### 9.2.4

Solve the equation.

- a)  $\frac{x}{4} = 2$       b)  $\frac{x}{9} = 8$       c)  $\frac{x}{7} = 7$       d)  $\frac{x}{15} = 10$

### 9.2.5

Solve the equation.

- a)  $18x - 27 = 9x + 36$       b)  $7x - 27 = 4x + 3$   
c)  $15x - 16 = 7x + 32$       d)  $13x - 42 = 7x + 12$   
e)  $4 + 9x = 13x - 32$       f)  $7x + 8 = 11x - 24$   
g)  $5x + 4 = 8x - 11$       h)  $7 + 10x = 14x - 9$

**9.2.6**

Given  $\triangle ABC$  with  $\angle C = 90^\circ$ . Prove that

$$\angle A = 90^\circ - \angle B$$

**9.2.7 (E22)**

Solve the equation

$$3 \cdot 24 \cdot 9 = 4 \cdot 9 \cdot x$$

**9.5.1 (1TV21D1)**

Solve the equation set.

$$2x - y = 4$$

$$x - 2y = 5$$

## Gruble 10

a) Prove that

$$0,2626... = \frac{26}{99}$$

Given

$$a = b \left( \frac{1}{10^c} + \frac{1}{10^{2c}} + \frac{1}{10^{3c}} + \dots \right)$$

where  $b$  is a number with  $c$  digits.

b) Prove that if  $b = 26$ , then  $a = 0,2626... \ .$

c) Prove that


$$a = \frac{b}{10^c - 1}$$

# Chapter 10

## Functions

# 10.1 Introduction

Variables are values that change. A value which changes in compliance with a variable is called a *function*.



In the above figure, the amount of boxes follows a specific pattern. Mathematically we can describe the pattern like this:

Number of boxes in Figure 1 =  $2 \cdot 1 + 1 = 3$   
Number of boxes in Figure 2 =  $2 \cdot 2 + 1 = 5$   
Number of boxes in Figure 3 =  $2 \cdot 3 + 1 = 7$   
Number of boxes in Figure 4 =  $2 \cdot 4 + 1 = 9$

Hence, for a figure of a random number  $x$ , we have

Number of boxes in Figure  $x = 2x + 1$

The amount of boxes changes in compliance with the change of  $x$ , in this case we say that

"Number of boxes in Figure  $x$ " is a function of  $x$ .

$2x + 1$  is the expression of the function "Number of boxes in Figure  $x$ ".

### General expressions

If we were to continue working with the function just studied, writing "Number of boxes in *Figure x*" all the time would be very cumbersome. It is common to let letters indicate functions and to write the associated variable inside parentheses. Let's rename "Number of boxes in *Figure x*" to  $a(x)$ . Then

$$\text{Number of boxes in Figure } x = a(x) = 2x + 1$$


If we write  $a(x)$ , but substitute  $x$  by a specific number, we substitute  $x$  by this number in the expression of our function:

$$a(1) = 2 \cdot 1 + 1 = 3$$

$$a(2) = 2 \cdot 2 + 1 = 5$$


$$a(3) = 2 \cdot 3 + 1 = 7$$

$$a(4) = 2 \cdot 4 + 1 = 9$$



### Example

Let the number of boxes in the below pattern be given by  $a(x)$ .



- a) Find the expression of  $a(x)$ .  
b) How many boxes are there when  $x = 10$ ?  
c) What is the value of  $x$  when  $a(x) = 628$ ?

### Answer

a) We notice that

- When  $x = 1$ , there are  $1 \cdot 1 + 3 = 4$  boxes.
- When  $x = 2$ , there are  $2 \cdot 2 + 3 = 7$  boxes.
- When  $x = 3$ , there are  $3 \cdot 3 + 3 = 12$  boxes.
- When  $x = 4$ , there are  $4 \cdot 4 + 3 = 17$  boxes.

Therefore

$$a(x) = x \cdot x + 3 = x^2 + 3$$

b)

$$a(10) = 10^2 + 3 = 100 + 3 = 103$$

When  $x = 10$ , there are 103 boxes.

c) We have the equation

$$x^2 + 3 = 628$$

$$x^2 = 625$$

Hence

$$x = 15 \quad \vee \quad x = -15$$

Since we seek a positive value of  $x$ , we have  $x = 15$ .



## 10.2 Linear functions and graphs

When a variable  $x$  and a function  $f(x)$  are present, we have two values; the value of  $x$  and the associated value of  $f(x)$ . These pairs of values can be put into a coordinate system (see [Section 1.3](#)) to form the *graph* of  $f(x)$ .

Let's use the function

$$f(x) = 2x - 1$$

as an example. We have

$$f(0) = 2 \cdot 0 - 1 = -1$$

$$f(1) = 2 \cdot 1 - 1 = 1$$

$$f(2) = 2 \cdot 2 - 1 = 3$$

$$f(3) = 2 \cdot 3 - 1 = 5$$

These pairs of values can be put into a table:

$x$	0	1	2	3
$f(x)$	-1	1	3	5

The above table yields the points


$$(0, -1) \quad (1, 1) \quad (2, 3) \quad (3, 5)$$

Now we place these points into a coordinate system (see the figure on page 201). Concerning functions, it is common to name the horizontal and the vertical axis the  $x$ -axis and the  $y$ -axis, respectively. Now the graph of  $f(x)$  is the curve passing through all the infinite many points we can create by the  $x$ -values and their associated  $f(x)$ -values. Our function is a *linear* function, which means its graph is a straight line. Hence, the graph is created by drawing the line going through the points we found.

As earlier mentioned, we can never draw a line, only a part of it. This also applies to graphs. In the figure on page 201 we have drawn the graph of  $f(x)$  for  $x$ -values in the range  $-2$  to  $4$ . That  $x$  is included in this *interval* we write as<sup>1</sup>  $-2 \leq x \leq 4$  or  $x \in [-2, 4]$ .

---

<sup>1</sup>Consult the list of symbols on page 4.



## 10.1 Linear functions

A function with the expression


$$f(x) = ax + b$$

where  $a$  and  $b$  are constants, is a *linear* function with *slope*  $a$  and *intercept*  $b$ .

The graph of a linear function is a straight line passing through the point  $(0, b)$ .

For two distinct  $x$ -values,  $x_1$  and  $x_2$ , we have

$$a = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$



### Example 1

Find the slope and the intercept of the functions.

$$f(x) = 2x + 1$$

$$g(x) = -3 + \frac{7}{2}$$

$$h(x) = \frac{1}{4}x - \frac{5}{6}$$

$$j(x) = 4 - \frac{1}{2}x$$

### Answer

- $f(x)$  have slope 2 and intercept 1.
- $g(x)$  have slope  $-3$  and intercept  $\frac{7}{2}$ .
- $h(x)$  have slope  $\frac{1}{4}$  and intercept  $-\frac{5}{6}$ .
- $j(x)$  have slope  $-\frac{1}{2}$  and intercept 4.

## Example 2

Draw the graph of

$$f(x) = \frac{3}{4}x - 2$$

for  $x \in [-5, 6]$ .

### Answer

To draw the graph of a linear function, we only need to know two points lying on it. The points are free to choose, therefore, in order to make calculations as simple as possible, we start off by finding the point where  $x = 0$ :

$$f(0) = \frac{3}{4} \cdot 0 - 2 = -2$$


Further on, we choose  $x = 4$ , since this also results in easy calculations:

$$f(4) = \frac{3}{4} \cdot 4 - 2 = 1$$

Now we have all the information we need and for the record we put it into a table:


$x$	0	4
$f(x)$	-2	1

Now we place the points in a coordinate system and draw the line passing through them:



### Example 3

Find the respective expressions of  $f(x)$  and  $g(x)$ .




### Answer

Firstly, we find the expression of  $f(x)$ . The point  $(0, 3)$  lies on the graph of  $f(x)$  (also see the figure on the next page). It then follows that  $f(0) = 3$ , and hence 3 is the intercept of  $f(x)$ . Moreover, we observe that  $(1, 2)$  also lies on the graph of  $f(x)$ . The slope of  $f(x)$  is then expressed by the fraction

$$\frac{2 - 3}{1 - 0} = -1$$

Therefore

$$f(x) = -x + 3$$



We now move our attention to finding the expression of  $g(x)$ . The point  $(0, -1)$  lies on the graph of  $g(x)$ . It then follows that  $f(0) = -1$ , and hence  $-1$  is the intercept of  $g(x)$ . Moreover, we observe that  $(5, 2)$  also lies on the graph of  $g(x)$ . The slope  $g(x)$  is then expressed by the fraction

$$\frac{2 - (-1)}{5 - 0} = \frac{3}{5}$$

Therefore

$$g(x) = \frac{3}{5}x + 1$$

## 10.1 Linear functions (explanation)

### The expression of $a$

Given a linear function

$$f(x) = ax + b$$

For two distinct  $x$ -values,  $x_1$  and  $x_2$ , we have

$$f(x_1) = ax_1 + b \quad (10.1)$$


$$f(x_2) = ax_2 + b \quad (10.2)$$

Subtracting (10.1) from (10.2), we get

$$\begin{aligned} f(x_2) - f(x_1) &= ax_2 + b - (ax_1 + b) \\ f(x_2) - f(x_1) &= ax_2 - ax_1 \\ f(x_2) - f(x_1) &= a(x_2 - x_1) \\ \frac{f(x_2) - f(x_1)}{x_2 - x_1} &= a \end{aligned} \quad (10.3)$$

### The graph of a linear function is a straight line

Given a linear function  $f(x) = ax + b$  and two distinct  $x$ -values  $x_1$  and  $x_2$ . Let  $A = (x_1, b)$ ,  $B = (x_2, b)$ ,  $C = (b, f(x_1))$ ,  $D = (0, f(x_2))$  and  $E = (0, b)$ .



By (10.3), we obtain

$$\begin{aligned} \frac{f(x_1) - f(0)}{x_1 - 0} &= a \\ \frac{ax_1 + b - b}{x_1} &= a \\ \frac{ax_1}{x_1} &= a \end{aligned} \quad (10.4)$$

Similarly,

$$\frac{ax_2}{x_2} = a \quad (10.5)$$

Moreover,

$$AC = f(x_1) - b = ax_1$$

$$BD = f(x_2) - b = ax_2$$

$$EA = x_1$$

$$EB = x_2$$

From (10.4) and (10.5) it follows that

$$\frac{ax_1}{x_1} = \frac{ax_2}{x_2}$$

Hence

$$\frac{AC}{BD} = \frac{EA}{EB}$$




In addition,  $\angle A = \angle B$ , so  $\triangle EAC$  and  $\triangle EBD$  satisfy term iii from [Rule 11.12](#), and hence the triangles are similar. Consequently,  $C$  and  $D$  lies on the same line, which must be the graph of  $f(x)$ .



## Exercises for Chapter 10

### 10.1.1




Let the number of boxes in the below figures be given by  $f(x)$ .



- Find an expression for  $f(x)$ .
- How many boxes are there when  $x = 100$ ?
- What is  $x$  when  $f(x) = 24$ ?

### 10.1.2


Let the number of boxes in the below figures be given by  $a(x)$ .



- Find an expression for  $a(x)$ .
- How many boxes are there when  $x = 20$ ?
- What is  $x$  when  $a(x) = 405$ ?

### 10.1.3


Let the number of boxes in the below figures be given by  $b(x)$ .



- Find an expression for  $b(x)$ .
- How many boxes are there when  $x = 20$ ?
- What is  $x$  when  $b(x) = 80$ ?

### 10.1.4 (EGV22D1)

The below figure shows the three first figures of a pattern. The figures are made up by triangles and squares.



How many triangles and how many squares will be present in figure number 10?

### 10.1.5

Let  $x$  be a positive integer.

- Make a function  $p(x)$  yielding the value of the  $x$ th positive, even number.
- Make a function  $o(x)$  yielding the value of the  $x$ th positive odd number.

### 10.2.1

Find the slope and the intercept of the function.

a)  $f(x) = 5x + 10$       b)  $g(x) = 3x - 12$

c)  $h(x) = -\frac{1}{7}x - 9$       d)  $i(x) = \frac{3}{2}x - \frac{1}{4}$

### 10.2.2

Draw the function on the interval  $x \in [-5, 5]$ :

a)  $f(x) = 2x - 1$       b)  $g(x) = -3x + 5$

### 10.3.1

Given the equation set

$$x - y = 5 \quad (\text{I})$$

$$x + y = 9 \quad (\text{II})$$

- a) Explain why the solutions of this set yield the interception point of the functions


$$f(x) = x - 5$$

$$g(x) = 9 - x$$

- b) Solve the equation set.


10.3.2

Find the expression of  $f(x)$




10.3.3

Find the expression of  $f(x)$



### 10.3.4

Find the expression of  $f(x)$  and  $g(x)$ .



### Gruble 11

Use the formulas from [Exercise 10.1.5](#) to prove that

- a) the sum/difference of two even numbers is an even number.
- b) sum/difference of two odd numbers is an even number.
- c) the sum/difference of an even number and an odd number is an odd number.

### Gruble 12

- a) A linear function  $f(x)$  has slope 3, and the point  $(2, 1)$  lies on the graph of  $f$ . Find the expression of  $f$ .
- b) A linear function  $f(x)$  has slope  $a$ , and the point  $(x_1, y_1)$  lies on the graph of  $f$ . Prove that

$$f(x) = a(x - x_1) + y_1$$

(This equation is called the **point-slope equation** .)

### Gruble 13

Given the functions  $f(x)$  and  $g(x)$ , where the graph of  $g$  is the line passing through  $A = (a, f(a))$  and  $B = (b, f(b))$ . Prove that

$$f - g = f(x) - \frac{f(b) - f(a)}{b - a}(x - a) + f(a)$$

## Chapter 11

# Geometry

## 11.1 Formulas of area and perimeter

A *formula* is an equation where (usually) one variable is isolated on one side of the equal sign. In [Section 7.4](#) we have already looked at the formulas for the area of rectangles and triangles, but there using words instead of symbols. Here we shall reproduce these two formulas, followed by other classical formulas for area and perimeter.

### 11.1 The area of a rectangle (7.4)


The area  $A$  of a rectangle with base  $g$  and height  $h$  is

$$A = gh$$



#### Example 1

Find the area of the rectangle.




#### Answer

The area  $A$  of the rectangle is

$$A = b \cdot 2 = 2b$$

#### Example 2

Find the area of the square.



#### Answer

The area  $A$  of the square is

$$A = a \cdot a = a^2$$



## 11.2 The area of a triangle (7.4)


The area  $A$  of a triangle with base  $g$  and height  $h$  is

$$A = \frac{gh}{2}$$



### Example

Which one of the triangles have the largest area?



### Answer

Let  $A_1$ ,  $A_2$  and  $A_3$  denote the areas of, respectively, the triangle to the left, in the middle and to the right. Then

$$A_1 = \frac{4 \cdot 3}{2} = 6$$

$$A_2 = \frac{2 \cdot 3}{2} = 3$$


$$A_3 = \frac{2 \cdot 5}{2} = 5$$

Hence, it is the triangle to the left which has the largest area.

### 11.3 The area of a parallelogram

The area  $A$  of a parallelogram with base  $g$  and height  $h$  is

$$A = gh$$



#### Example

Find the area of the parallelogram




#### Answer

The area  $A$  of the parallelogram is

$$A = 5 \cdot 2 = 10$$

### 11.3 The area of a parallelogram (explanation)

From a parallelogram we can always, by drawing one of its diagonals, form two triangles which both have base  $g$  and height  $h$ .




Hence, both triangles have an area equal to  $\frac{gh}{2}$ . Therefore, the area  $A$  of the parallelogram is

$$\begin{aligned} A &= \frac{gh}{2} + \frac{gh}{2} \\ &= g \cdot h \end{aligned}$$

### 11.4 The area of a trapezoid


The area  $A$  of a trapezoid with parallel sides  $a$  and  $b$  and height  $h$  is

$$A = \frac{h(a + b)}{2}$$



#### Example

Find the area of the trapezoid.



#### Answer

The area  $A$  of the trapezoid is


$$\begin{aligned} A &= \frac{3(6 + 4)}{2} \\ &= \frac{3 \cdot 10}{2} \\ &= 15 \end{aligned}$$

#### Note

In respect of a base and a height, the area formulas for a parallelogram and a rectangle are identical. Applying [Rule 11.4](#) on a parallelogram also results in an expression equal to  $gh$ . This follows from the fact that a parallelogram is just a special case of a trapezoid (and a rectangle is just a special case of a parallelogram).

### 11.4 The area of a trapezoid (explanation)

In a trapezoid, we can, by drawing one of the diagonals, create two triangles:



In the above figure we have

$$\text{The area of the blue triangle} = \frac{ah}{2}$$

$$\text{The area of the green triangle} = \frac{bh}{2}$$


Therefore, the area  $A$  of the trapezoid is

$$\begin{aligned} A &= \frac{ah}{2} + \frac{bh}{2} \\ &= \frac{h(a+b)}{2} \end{aligned}$$

### 11.5 The perimeter of a circle (and the value of $\pi$ )

The perimeter (the circumference)  $O$  of a circle with radius  $r$  is

$$O = 2\pi r$$



$$\pi = 3.141592653589793....$$

#### Example 1

Find the circumference of the circle.



#### Answer


The circumference  $O$  is

$$\begin{aligned} O &= 2\pi \cdot 3 \\ &= 6\pi \end{aligned}$$

## 11.6 The area of a circle


The area  $A$  of a circle with radius  $r$  is

$$A = \pi r^2$$



### Example

Find the area of the circle.




### Answer

The area  $A$  of the circle is

$$A = \pi \cdot 5^2 = 25\pi$$


## 11.6 The area of a circle (explanation)

In the below figure, we have divided a circle into 4, 10 and 50 (equal-sized) sectors, and placed them consecutively.



In each case, the arcs make up the circumference of the circle. If the circle has radius  $r$ , the sum of the arcs equals  $2\pi r$ . And when there are equally many sectors turned upwards as downwards, the total length of the arcs equals  $\pi r$  on both the bottom and the top.

The more sectors the circle is divided into, the more the composition takes the form of a rectangle (in the figure below there are 100 sectors). The base  $g$  of this "rectangle" approximately equals  $\pi r$ , while the height  $h$  approximately equals  $r$ .





Hence, the area  $A$  of the "rectangle", that is, the circle, is

$$A \approx gh \approx \pi r \cdot r = \pi r^2$$

## 11.7 Pythagoras's theorem


In a right triangle, the area of the square formed by the hypotenuse equals the sum of the areas of the squares formed by the legs.

$$a^2 + b^2 = c^2$$



### Example 1

Find the length of  $c$ .



### Answer

We know that

$$c^2 = a^2 + b^2$$

where  $a$  and  $b$  are the legs of the right triangle. Therefore

$$\begin{aligned} c^2 &= 4^2 + 3^2 \\ &= 16 + 9 \\ &= 25 \end{aligned}$$

Hence,

$$c = 5 \quad \vee \quad c = -5$$

Since  $c$  is a length,  $c = 5$ .




## 11.7 Pythagoras's theorem (explanation)

The below figure shows equal-sized squares divided into different shapes.



We observe the following:

1. The area of the red square is  $a^2$ , the area of the purple square is  $b^2$  and the area of the blue square is  $c^2$ .
2. The area of an orange rectangle is  $ab$  and the area of a green triangle is  $\frac{ab}{2}$ .
3. If we remove the two orange rectangles and the four green triangles, the remaining area to the left equals the remaining area to the right (by remark 2).



Hence

$$a^2 + b^2 = c^2 \quad (11.1)$$


Given a triangle with sides of length  $a, b$  and  $c$ , of which  $c$  is the longest. As long as the triangle is right, we can always form two squares with sides of length  $a + b$ , as in the initial figure. Therefore, (11.1) is valid for alle right triangles.

## 11.2 Congruent and similar triangles

### 11.8 Unique construction of triangles


A triangle  $\triangle ABC$ , as shown in the below figure, can be uniquely constructed if one of the following terms are satisfied:

- i)  $c, \angle A$  and  $\angle B$  are known.
- ii)  $a, b$  and  $c$  are known.
- iii)  $b, c$  and  $\angle A$  are known.



### 11.9 Congruent triangles

Two triangles of equal shape and size are congruent.




The congruence in the above figure is written

$$\triangle ABC \cong \triangle DEF$$

### 11.10 Similar triangles

Similar triangles constitute three pairs of equal angles.




The similarity in the above figure is written

$$\triangle ABC \sim \triangle DEF$$

### Corresponding sides

When studying similar triangles, *corresponding* sides plays an important role. Corresponding sides are sides in similar triangles adjacent to the same angle.



Regarding the similar triangles  $\triangle ABC$  and  $\triangle DEF$  we have

In  $\triangle ABC$  is

- $BC$  adjacent to  $u$ .
- $AC$  adjacent to  $v$
- $AB$  adjacent to  $w$ .

In  $\triangle DEF$  is

- $FE$  adjacent to  $u$ .
- $FD$  adjacent to  $v$
- $ED$  adjacent to  $w$ .


This means that these are corresponding sides:

- $BC$  and  $FE$
- $AC$  and  $FD$
- $AB$  and  $ED$

#### 11.11 Ratios in similar triangles

If two triangles are similar, the ratios of corresponding sides are equal<sup>1</sup>.

$$\frac{AB}{DE} = \frac{AC}{DF} = \frac{BC}{EF}$$



<sup>1</sup>Here, we take it for granted that corresponding sides are apparent from the figure.


### Notice

From [Rule 11.11](#) it follows that

$$\frac{AB}{BC} = \frac{DE}{EF} \quad , \quad \frac{AB}{AC} = \frac{DE}{DF} \quad , \quad \frac{BC}{AC} = \frac{EF}{DF}$$

### Example

The triangles are similar. Find the length of  $EF$ .



### Answer

We observe that  $AB$  corresponds to  $DE$ ,  $BC$  to  $EF$  and  $AC$  to  $DF$ . Therefore

$$\frac{DE}{AB} = \frac{EF}{BC}$$

$$\frac{10}{5} = \frac{EF}{3}$$

$$2 \cdot 3 = \frac{EF}{3} \cdot 3$$

$$6 = EF$$


### 11.12 Terms of similar triangles

Two triangles  $\triangle ABC$  and  $\triangle DEF$  are similar if one of these terms are satisfied:

i) They constitute two pairs of equal angles.


ii)  $\frac{AB}{DE} = \frac{AC}{DF} = \frac{BC}{EF}$

iii)  $\frac{AB}{DE} = \frac{AC}{DF}$  and  $\angle A = \angle D$ .



#### Example 1

$\angle ACB = 90^\circ$ . Show that  $\triangle ABC \sim \triangle ACD$ .




#### Answer

$\triangle ABC$  and  $\triangle ACD$  are both right and they have  $\angle DAC$  in common. Hence, the triangles satisfy term *i* from [Rule 11.12](#), and therefore they are similar.

*Note:* Similarly it can be shown that  $\triangle ABC \sim \triangle CBD$ .

### Example 2

Examine whether the triangles are similar.



### Answer

We have


$$\frac{AC}{FD} = \frac{18}{12} = \frac{3}{2}, \quad \frac{BC}{FE} = \frac{9}{6} = \frac{3}{2}, \quad \frac{AB}{DE} = \frac{12}{10} = \frac{6}{5}$$

$$\frac{AC}{IG} = \frac{18}{12} = \frac{3}{2}, \quad \frac{BC}{IH} = \frac{9}{6} = \frac{3}{2}, \quad \frac{AC}{IG} = \frac{18}{12} = \frac{3}{2}$$

Hence,  $\triangle ABC$  and  $\triangle GHI$  satisfy term *ii* from [Rule 11.12](#), and therefore they are similar. (Hence,  $\triangle GHI$  and  $\triangle FED$  are not similar.)

### Example 3

Examine whether the triangles are similar.



### Answer

We have  $\angle BAC = \angle EDF$ . Also,

$$\frac{ED}{AB} = \frac{8}{4} = 2, \quad \frac{FD}{AC} = \frac{14}{7} = 2$$


Hence, term *iii* from [Rule 11.12](#) is satisfied, and therefore the triangles are similar.

## 11.3 Explanations

### 11.5 The perimeter of a circle (and the value of $\pi$ ) (explanation)


*Here we shall use regular polygons along the path to our wanted result. In regular polygons, all sides are of equal length. Since all polygons here to be mentioned are regular, we'll mention them simply as polygons.*

We'll start off by examining some approximations of the circumference  $O_1$  of a circle with radius 1.



#### Upper and lower boundary

When seeking a value, it is a good habit to conclude how large or small you expect it to be. With this target, we enclose the circle by a square with sides of length 2:




Clearly, the circumference of the circle is smaller than the perimeter of the square, therefore

$$\begin{aligned} O_1 &< 2 \cdot 4 \\ &< 8 \end{aligned}$$

Now we inscribe a 6-gon (hexagon). The hexagon can be divided into 6 equilateral triangles with, necessarily, sides of length 1. The circumference of the circle must be larger than the perimeter of the hexagon, so


$$\begin{aligned} O_1 &> 6 \cdot 1 \\ &> 6 \end{aligned}$$




Now advancing to a more sophisticated hunt for the circumference, we know that we seek a value between 6 and 8.

### Increasingly better approximations

The idea of inscribing polygons carries on. We let the below figures work as a sufficient prove of the fact that the more sides of the polygon, the better estimate its perimeter makes of the circumference of the circle.




(a) 6-gon



(b) 12-gon

Since a 6-gon has sides of length 1, it is tempting to examine if this can help us find the side lengths of other polygons. By inscribing both a 6-gon and a 12-gon (and drawing a triangle) we have a figure like this:





(a) A 6-gon and a 12-gon together with a triangle formed by the circle center and one side of the 12-gon.



(b) The triangle from figure (a).

Let  $s_{12}$  and  $s_6$  denote the side lengths of the 12-gon and the 6-gon, respectively. Moreover, we observe that both  $A$  and  $C$  lies on the circular arc and that both  $\triangle ABC$  and  $\triangle BSC$  are right-angled (explain to yourself why!). We have

$$\begin{aligned} SC &= 1 \\ BC &= \frac{s_6}{2} \\ SB &= \sqrt{SC^2 - BC^2} \\ BA &= 1 - SB \\ AC &= s_{12} \\ s_{12}^2 &= BA^2 + BC^2 \end{aligned}$$

To find  $s_{12}$ , we need to know  $BA$ , and to find  $BA$  we need to know  $SB$ . Hence, we start off finding  $SB$ . Since  $SC = 1$  and  $BC = \frac{s_6}{2}$ ,

$$\begin{aligned} SB &= \sqrt{1 - \left(\frac{s_6}{2}\right)^2} \\ &= \sqrt{1 - \frac{s_6^2}{4}} \end{aligned}$$

Now we focus on finding  $s_{12}$ :

$$\begin{aligned} s_{12}^2 &= (1 - SB)^2 + \left(\frac{s_6}{2}\right)^2 \\ &= 1^2 - 2SB + SB^2 + \frac{s_6^2}{4} \end{aligned}$$

At first, it looks like the expression to the right cannot be simplified, but a small operation can change this. If  $-1$  was a term present, we could have combined  $-1$  and  $\frac{s_6^2}{4}$  to become  $-SB^2$ . We obtain  $-1$  by both adding and subtracting it on the right side of the equation:

$$\begin{aligned}
 s_{12}^2 &= 1 - 2SB + SB^2 + \frac{s_6^2}{4} - 1 + 1 \\
 &= 2 - 2SB + SB^2 - \left(1 - \frac{s_6^2}{4}\right) \\
 &= 2 - 2SB + SB^2 - SB^2 \\
 &= 2 - 2SB \\
 &= 2 - 2\sqrt{1 - \frac{s_6^2}{4}} \\
 &= 2 - \sqrt{4} \sqrt{1 - \frac{s_6^2}{4}} \\
 &= 2 - \sqrt{4 - s_6^2}
 \end{aligned}$$

Hence


$$s_{12} = \sqrt{2 - \sqrt{4 - s_6^2}}$$

Even though we have derived a relation between the side lengths  $s_{12}$  and  $s_6$ , this relation is valid for all pairs of side lengths where one is the side length of a polygon with twice as many sides as the other. Now let  $s_n$  and  $s_{2n}$ , respectively, denote the side lengths of a polygon and a polygon with twice as many sides. Then


$$s_{2n} = \sqrt{2 - \sqrt{4 - s_n^2}} \quad (11.2)$$

The perimeter of a polygon inscribed in the circle is an estimate of the circumference. Applying (11.2), we can successively find the side length of a polygon with twice as many sides as the previous. The below table shows the side length and the associated estimate of the circumference up to a 96-gon:


<i>Side length formula</i>	<i>Side length</i>	<i>Estimate, circumference</i>
	$s_6 = 1$	$6 \cdot s_6 = 6$
$s_{12} = \sqrt{2 - \sqrt{4 - s_6^2}}$	$s_{12} = 0.517\dots$	$12 \cdot s_{12} = 6.211\dots$
$s_{24} = \sqrt{2 - \sqrt{4 - s_{12}^2}}$	$s_{24} = 0.261\dots$	$24 \cdot s_{24} = 6.265\dots$
$s_{48} = \sqrt{2 - \sqrt{4 - s_{24}^2}}$	$s_{48} = 0.130\dots$	$48 \cdot s_{48} = 6.278\dots$
$s_{96} = \sqrt{2 - \sqrt{4 - s_{48}^2}}$	$s_{96} = 0.065\dots$	$96 \cdot s_{96} = 6.282\dots$




(a) 6-gon




(b) 12-gon



(c) 24-gon



(d) 48-gon



(e) 96-gon

In fact, the mathematician [Archimedes](#) reached as far as the above calculation approximately 250 b.c!

A computer has no problems performing calculations<sup>1</sup> on a polygon with extremely many sides. Calculating the perimeter of a 201 326 592-gon yields


Circumference of a circle with radius 1 = 6.283185307179586...

(With the aid of more advanced mathematics it can be proved that the circumference of a circle with radius 1 is an irrational number, but that the digits shown above are correct, thereby the equal sign.)


## The formula and $\pi$

We shall now derive the famous formula for the circumference of any circle. Here as well, we take it for granted that the perimeter of an inscribed polygon yields an estimate of the circumference which gets more accurate the more sides the polygon has.

For the sake of simplicity, we shall use inscribed squares to illustrate the outline. We draw two circles of random size, but the one larger than the other, and inscribe a square in both. Let  $R$  and  $r$  denote the radius of the larger and the smaller circle, respectively. Also, let  $K$  and  $k$  denote the side length of the larger and the smaller square, respectively.



Both squares can be divided into four isosceles triangles:



Since these triangles are similar,

$$\frac{K}{R} = \frac{k}{r} \quad (11.3)$$

Let  $\tilde{O} = 4K$  and  $\tilde{o} = 4k$  denote the estimated circumferences of the larger and the smaller circle, respectively. Multiplying both

sides of (11.3) by 4 yields

$$\frac{4A}{R} = \frac{4a}{r} \quad (11.4)$$

$$\frac{\tilde{O}}{R} = \frac{\tilde{o}}{r} \quad (11.5)$$

Now we observe this:

*If we were to inscribe polygons with 4, 100 or any number of sides, the polygons could still be divided into triangles obeying (11.3). And in the same way as we did in the above example, we could then rewrite (11.3) into (11.5).*

Let's therefore imagine polygons with such a large number of sides that we accept their respective perimeters as equal to the respective circumferences of the circles. Letting  $O$  and  $o$  denote the circumferences of the larger and smaller circle respectively, we have

$$\frac{O}{R} = \frac{o}{r}$$

Since the circles are randomly chosen, we conclude that *all circles have the same ratio of the circumference to the radius*. An equivalent statement is that *all circles have the same ratio of the circumference to the diameter*.

The ratio of the circumference  $O$  to the diameter  $d$  in a circle is named  $\pi$  (pronounced "pi"):

$$\frac{O}{d} = \pi$$

The above equation yields the formula for the circumference of a circle with diameter  $d$  and radius  $r$ :

$$\begin{aligned} O &= \pi d \\ &= 2\pi r \end{aligned}$$

Earlier we found that the circumference of a circle with radius 1 (and diameter 2) equals 6.283185307179586... Hence


$$\begin{aligned} \pi &= \frac{6.283185307179586...}{2} \\ &= 3.141592653589793... \end{aligned}$$

---

<sup>1</sup>For those interested in computer programming; the iteration algorithm must be altered in order to avoid instabilities when the number of sides are large.

### 11.11 Ratios in similar triangles (explanation)

Here, we shall write the area of a triangle  $\triangle ABC$  as  $ABC$ .



In the figure above, we have  $BB' \parallel CC'$ . With  $BB'$  as base,  $HB'$  is the height of both  $\triangle CBB'$  and  $\triangle C'B'B'$ . Therefore

$$CBB' = C'BB' \quad (11.6)$$

Moreover,

$$ABB' = AB \cdot HB'$$

$$CBB' = BC \cdot HB'$$

Hence

$$\frac{ABB'}{CBB'} = \frac{AB}{BC} \quad (11.7)$$

Similarly,

$$\frac{ABB'}{C'BB'} = \frac{AB'}{B'C'} \quad (11.8)$$

From (11.6), (11.7) and (11.8) it follows that

$$\frac{AB}{BC} = \frac{ABB'}{CBB'} = \frac{ABB'}{C'BB'} = \frac{AB'}{B'C'} \quad (11.9)$$

For the similar triangles  $\triangle ACC'$  and  $\triangle ABB'$ ,

$$\begin{aligned}\frac{AC}{AB} &= \frac{AB + BC}{AB} \\ &= 1 + \frac{BC}{AB}\end{aligned}$$


$$\begin{aligned}\frac{AC'}{AB'} &= \frac{AB' + B'C'}{AB'} \\ &= 1 + \frac{B'C'}{AB'}\end{aligned}$$

By (11.9), the ratio of corresponding sides in the two triangles are equal.

## Note

In the following explanations of term *ii* and *iii* from [Rule 11.8](#) we assume this:

- Two circles intersect in maximum two points.
- Given a coordinate system placed in the center of one of the circles, such that the horizontal axis passes through both circle centers. If  $(a, b)$  is one of the intersection points,  $(a, -b)$  is the other.




The remarks above are quite easy to prove, but since they are largely intuitively true, we hold them as granted. This implies that the triangle formed by the two centers and one of the intersection points is congruent to the triangle formed by the two centers and the other intersection point. By this, we can study attributes of triangles with the aid of semi-circles.

## 11.8 Unique construction of triangles (explanation)


### Term i

Given a length  $c$  and two angles  $u$  and  $v$ . We make a segment  $AB$  with length  $c$ . Then we draw two angle sides, such that  $\angle A = u$  and  $\angle B = v$ . As long as these angle sides are not parallel, they must intersect in one, and one only, point ( $C$  in the figure). Together with  $A$  and  $B$ , this point will form a triangle uniquely determined by  $c$ ,  $u$  and  $v$ .



### Term ii

Given three lengths  $a$ ,  $b$  and  $c$ . We make a segment  $AB$  with length  $c$ . Then we make two semi-circles with respective radii  $a$  and  $b$  and centers  $B$  and  $A$ . If a triangle  $\triangle ABC$  is to have sides of length  $a$ ,  $b$  and  $c$ ,  $C$  must lie on both of the semi-circles. Since the semi-circles intersect in one point only,  $\triangle ABC$  is uniquely determined by  $a$ ,  $b$  and  $c$ .




### Term iii

Given two lengths  $b$  and  $c$  and an angle  $u$ . We begin as follows:

1. We make a segment  $AB$  with length  $c$ .
2. In  $A$  we draw a semi-circle with radius  $b$ .

By placing  $C$  randomly on the arc of the semi-circle, we get all instances of a triangle  $\triangle ABC$  with sides of length  $AB = c$  and  $AC = b$ . Specifically placing  $C$  on the arc of the semi-circle is equivalent to setting a specific value of  $\angle A$ . Now it remains to show that every placement of  $C$  implies a unique length of  $BC$ .





Let  $C_1$  and  $C_2$  denote two potential placements of  $C$ , where  $C_2$ , along the semicircle, lies closer to  $E$  than  $C_1$ . Now we dot a circular arc with radius  $BC_1$  and center  $B$ . Since the dotted arc and the semi-circle only intersects in  $C_1$ , other points will either lie inside or outside the dotted arc. Necessarily,  $C_2$  lies outside the dotted arc, and therefore  $BC_2$  is longer than  $BC_1$ . From this we can conclude that the length of  $BC$  increases as  $C$  moves against  $E$  along the semi-circle. Therefore, specifying  $\angle A = u$  yields a unique value of  $BC$ , and hence a unique triangle  $\triangle ABC$  where  $AC = b$ ,  $c = AB$  and  $\angle BAC = u$ .

## 11.12 Terms of similar triangles (explanation)

### Term i

Given two triangles  $\triangle ABC$  and  $\triangle DEF$ . By [Rule 7.3](#),


$$\angle A + \angle B + \angle C = \angle D + \angle E + \angle F$$

If  $\angle A = \angle D$  and  $\angle B = \angle E$ , it follows that  $\angle C = \angle F$ .

### Term ii

Given two triangles  $\triangle ABC$  and  $\triangle DEF$ , where

$$\frac{AC}{DF} = \frac{BC}{EF} \quad , \quad \angle C = \angle F \quad (11.10)$$



Let  $a = BC$ ,  $b = AC$ ,  $d = EF$  and  $e = DF$ . We place  $D'$  and  $E'$  on  $AC$  and  $BC$ , respectively, such that  $D'C = e$  and  $AB \parallel D'E'$ . Then  $\triangle ABC \sim \triangle D'E'C$ , and hence

$$\frac{E'C}{BC} = \frac{D'C}{AC}$$

$$E'C = \frac{ae}{b}$$

By (11.10),

$$EF = \frac{ae}{b}$$

Hence  $E'C = EF$ . From term ii of [Rule 11.8](#) it now follows that  $\triangle D'E'C \cong \triangle DEF$ . This implies that  $\triangle ABC \sim \triangle DEF$ .

### Term iii

Given two triangles  $\triangle ABC$  and  $\triangle DEF$ , where

$$\frac{AB}{DE} = \frac{AC}{DF} = \frac{BC}{EF} \quad (11.11)$$

We place  $D'$  and  $E'$  on  $AC$  and  $BC$ , respectively, such that  $D'C = e$  and  $E'C = d$ . From term i of [Rule 11.12](#) we have  $\triangle ABC \sim \triangle D'E'C$ . Therefore


$$\frac{D'E'}{AB} = \frac{D'C}{AC}$$

$$D'E' = \frac{ae}{c}$$

By (11.11),

$$f = \frac{ae}{c}$$


Hence, the side lengths of  $\triangle D'E'C$  and  $\triangle DEF$  are pairwise equal, and then, from term i of [Rule 11.8](#), they are congruent. This implies that  $\triangle ABC \sim \triangle DEF$ .



# Exercises for Chapter 11


## 11.1.1

The triangles are similar. Find the value of  $\angle ACB$ .




## 11.1.2

The triangles are similar. Find the three pairs of corresponding sides.




## 11.1.3

The triangles are similar. Find the length of  $EF$  and the length of  $DF$ .




### 11.1.4

The triangles are similar. Find the length of  $AC$  and the length of  $DF$ .



### 11.1.5


Find all the similar triangles defined by  $A, B, C$  og  $D$ .



### 11.1.6

$\triangle ABC$  and  $\triangle DEF$  are similar.

- What is the ratio of the area of  $\triangle DEF$  to the area of  $\triangle ABC$  if  $h_1 = 2$  and  $h_2 = 6$ ?
- Given a number  $a$ . What is the ratio of the area of  $\triangle DEF$  to the area of  $\triangle ABC$  if  $h_2 = ah_1$ ?



### 11.1.7

En kjegle has radius 10 and height 4.

- a) Find the surface area of the cone.
- b) Find the volume of the cone.


### 11.1.8

- a) Two spheres have radii 2 and 6, respectively. What is the ratio of the volume of the larger sphere to the volume of the smaller sphere?
- b) A sphere has radius  $r$ , and another sphere has radius  $ar$ , where  $a > 1$ . What is the ratio of the volume of the larger sphere to the volume of the smaller sphere?

### Gruble 14

(GV21D1)

The polygon is a regular<sup>1</sup> hexagon. Find the value of  $v$ .




### Gruble 15

Given an isosceles triangle  $\triangle ABC$ , with  $AC = BC$ . Prove that<sup>2</sup> the bisector of  $\angle ACB$  is the perpendicular bisector of  $AB$ .

### Gruble 16

$\triangle ABC$  is equilateral and have sides of length  $s$ .



- Prove that in a triangle with angles  $30^\circ$ ,  $60^\circ$ ,  $90^\circ$ , the hypotenuse is twice as long as the shortest leg.
- Prove that the height of  $\triangle ABC$  is  $\frac{\sqrt{3}}{2}s$ .

---


<sup>1</sup>Regular polygons have sides of equal length.

<sup>2</sup>The definition of the bisector of an angle and the perpendicular bisector of a segment is found in [TM1](#).

### Gruble 17


Prove that the double area of  $\triangle ABC$  can be written as

$$AE \cdot BD + CE \cdot AD$$



### Gruble 18

A **median** of a triangle is a segment joining a vertex to midpoint of the opposite side.



Given  $\triangle ABC$  with medians  $AE$ ,  $BF$  and  $CD$ .


- Prove that  $AE$ ,  $BF$  and  $CD$  intersect at exactly one point ( $G$  in the figure).
- Prove that

$$\frac{GC}{DG} = \frac{GB}{FG} = \frac{GA}{EG} = 2$$

Note: Part b) is probably easier than part a).

**Gruble 19**

The three circles have radius 2, and  $A$ ,  $B$  and  $C$  lie on the same line. Find the area of the red surface.




Hint: You could get use of the fact that a sector with angle  $v$  makes up  $\frac{v}{360^\circ}$  of the area of a circle with equal radius.

**Gruble 20**


The colored surfaces makes up a square, and  $F$ ,  $G$ ,  $H$  and  $I$  are, respectively, the square's midpoints.

Prove that the area of the blue surface equals the area of the green surface.



**Gruble 21**


The square has sides of length 4. Find the radius of the circle.






Gruble 22

a) Prove that  $\frac{a}{b} = \sqrt{2} + \sqrt{6}$ .



*Note:* To solve this problem you may (bot far from necessarily) need the  $abc$ -formula found in [TM1](#).


b)  $AD = BC$ . Find the value of  $\angle A$ .



### Gruble 23


*Note:* This exercise deals with results that at first seems obvious, but turns out rather har to prove.

- a) Prove that if  $AC = BC$ , then  $\angle A = \angle B$ .




*Note:* Earlier on, we have declared that an isosceles triangle has two equal angles, but strictly we have to prove it.

- b) Prove that if  $AC > BC$ , then  $\angle B > \angle C$ .



- c) Given  $\triangle ABC$ , where  $AB$  is the longest side. Prove that when  $AB$  is the base, the height lies inside the triangle.
- d) In the figure below,  $c$  is the longest side of the triangle.



Prove that

$$c > a + b \quad , \quad b + c > a \quad , \quad a + c > b$$

*Note:* These three inequalities are called the **triangle inequality**.

## Comment (for the particularly interested)


Also in geometry, axioms (see comment on page 177) lays the ground of the mathematical system we create, but the axiomatic structure of geometry is quite extensive and intricate. In addition, some theorems are such intuitively true that it, at least in a book like this, would be more confusing than clarifying to explain them all in detail.

However, it is worth noticing that [Rule 11.8](#) states three terms regarding the unique construction in a triangle, and [Rule 11.9](#) states a term regarding congruence. In more advanced texts on geometry, chances are that you will recognize the content of these rules as axioms and theorems of congruence:

### Congruence

Two triangles  $\triangle ABC$  and  $\triangle DEF$  are congruent if one of the following terms are satisfied:

- i)  $AB = DE$ ,  $BC = EF$  and  $\angle A = \angle D$ .
- ii)  $\angle A = \angle D$ ,  $\angle B = \angle E$  and  $AB = DE$ .
- iii)  $AB = DE$ ,  $BC = EF$  and  $AC = FD$ .
- iv)  $\angle A = \angle D$  and  $\angle B = \angle E$  and, in addition,  $AB = DE$  or  $BC = EF$  or  $AC = FD$ .



- 
- i) The Side-angle-side (SAS) axiom
  - ii) The Angle-side-angle (ASA) theorem
  - iii) The Side-side-side (SSS) theorem
  - iv) The Side-angle-angle (SAA) theorem

In the text-box on the previous side, term i) - iii) brings sufficient information regarding the unique construction of a triangle. However, in this book we have chosen to separate the concepts of congruence and unique construction. This is done under the presumption that most people will have a good intuition about congruent triangles, while having more difficulties stating terms of unique construction — and it is not necessarily easy to observe this directly from the terms of congruence.

Also, observe that iv) is just term ii) in a wider sense, but it cannot be used as a term of unique construction. Therefore, this term is not found in either [Rule 11.8](#) or [Rule 11.9](#).

# Exercise answers

## Chapter1

- 1.1.1 a) 22      b) 13      c) 36  
1.1.2 a) 17      b) 29      c) 11  
1.1.3 a) 1,7      b) 2,3  
1.1.4 a) 13,9      b) 32,8      c) 0,7      d) 2,4  
1.1.5 a)  
1.1.6 c)  
1.1.7 b)  
1.1.8 a)  
1.1.7 c)

## Chapter6

- 2.1.1 *Note:* Multiple possible answers. a)  $4 = 1+3$       b)  $5 = 2+3$       c)  $6 = 2+4$   
d)  $7 = 1+6$       e)  $8 = 3+5$       f)  $9 = 2+7$   
2.1.2 a)  $5 = 2+2+1$       b)  $6 = 1+3+2$       c)  $7 = 3+2+2$       d)  $8 = 1+1+6$   
e)  $9 = 3+3+3$       f)  $2+5+3$   
2.1.3  
1) a) 8      b) 7      c) 6      d) 5  
2) Fordi de er svarene i oppgave 1a), 1b) og 1c).  
2.1.4 a) 1)      b) 1)      c) 2)  
2.2.1 *Note:* Multiple possible answers. a)  $2 = 7-5$       b)  $3 = 10-7$       c)  $4 = 5-1$   
d)  $5 = 10-5$       e)  $6 = 9-3$       f)  $7 = 9-2$       g)  $8 = 10-2$   
2.2.2 a) 1)      b) 1)      c) 2)  
2.3.1  
a)  $2+2+2 = 2 \cdot 4 = 4+4$   
b)  $3+3+3+3+3+3 = 3 \cdot 6 = 6+6+6$   
c)  $4+4 = \cdot 4 \cdot 2 = 2+2+2+2$   
d)  $5+5+5+5+5+5+5+5+5+5 = 5 \cdot 10 = 10+10+10+10+10$   
e)  $6+6+6+6 = 6 \cdot 5 = 5+5+5+5+5$   
f)  $7+7+7+7 = 7 \cdot 4 = 4+4+4+4+4+4+4$

**2.3.2** a) 20      b) 24      c) 18      d) 30      e) 56

**2.3.3** a) partall      b) partall, 0      c) oddetall, 5

## Chapter??

**3.2.1** *Note:* Multiple possible answers. a)  $5 \cdot 20$       b)  $3 \cdot 10$       c)  $2 \cdot 20$

d)  $2 \cdot 35$

e)  $7 \cdot 6$       f)  $8 \cdot 4$       g)  $42 \cdot 2$       h)  $3 \cdot 30$

**3.2.2** a)  $2 \cdot 2 \cdot 2 \cdot 2 \cdot 3$       b)  $2 \cdot 2 \cdot 3 \cdot 5$       c)  $2 \cdot 5$

**3.2.4** 28

### Gruble 1

**Gruble 2** See the solutions manual.

**3.1.1** a) 34      b) 177      c) 100      d) 664      e) 2943

## Chapter??

**4.1.1** a) 6      b) 5      c) 2      d) 7      e) 9      f) 4

**4.1.2** a) 0,5      b) 0,25      c) 0,2      d) 0,75      e) 0,4      f) 0,6  
g) 0,8      h) 1,5      i) 0,33...      j) 2,5      k) 0,833...      l) 1,4      m) 2,75  
n) 0,7

**4.1.3** a)  $\frac{3}{4}$       b)  $\frac{2}{7}$       c)  $\frac{2}{5}$

**4.1.4** a)  $\frac{14}{3}$       b)  $\frac{13}{2}$       c)  $\frac{11}{5}$

**4.2.1** a)  $\frac{20}{6}$       b)  $\frac{9}{12}$       c)  $\frac{12}{28}$       d)  $\frac{45}{40}$       e)  $\frac{54}{30}$       f)  $\frac{77}{28}$

**4.2.2** a)  $\frac{9}{4}$       b)  $\frac{9}{5}$       c)  $\frac{2}{7}$

**4.3.1** a)  $\frac{10}{3}$       b)  $\frac{14}{4}$       c)  $\frac{11}{6}$       d)  $\frac{10}{7}$       e) 1

*Note:* Éin av brøkane kan forkortast.

**4.3.2** a)  $\frac{22}{3}$       b)  $\frac{8}{5}$       c)  $\frac{17}{7}$

**4.3.3** a)  $\frac{1}{3}$       b)  $\frac{2}{4}$       c)  $\frac{9}{6}$       d) 1      e) 0

*Note:* To av brøkane kan forkortast.

**4.3.4** a)  $\frac{6}{5}$       b)  $\frac{5}{7}$       c) 4

**4.3.5** a)  $\frac{9}{10}$       b)  $\frac{73}{63}$       c)  $\frac{101}{24}$       d)  $\frac{73}{20}$       e)  $\frac{5}{6}$

**4.3.6** a)  $\frac{1}{10}$       b)  $\frac{29}{36}$       c)  $\frac{71}{72}$       d)  $\frac{11}{20}$       e)  $\frac{5}{6}$

**4.3.7** a)  $\frac{5}{12}$       b)  $\frac{157}{30}$       c)  $\frac{229}{56}$

**4.4.1** a)  $\frac{20}{3}$       b)  $\frac{40}{7}$       c)  $\frac{54}{10}$       d)  $\frac{80}{7}$       e)  $\frac{21}{2}$       f)  $\frac{28}{3}$       g)  $\frac{35}{3}$   
h)  $\frac{30}{7}$       i)  $\frac{5}{11}$       j)  $\frac{72}{17}$

**4.5.1** a)  $\frac{4}{15}$       b)  $\frac{5}{56}$       c)  $\frac{9}{60}$       d)  $\frac{8}{70}$       e)  $\frac{3}{14}$       f)  $\frac{9}{110}$       g)  $\frac{1}{60}$   
h)  $\frac{9}{290}$       i)  $\frac{8}{459}$       j)  $\frac{4}{158}$

**4.6.1** a)  $\frac{20}{27}$       b)  $\frac{7}{32}$       c)  $\frac{18}{21}$       d)  $\frac{60}{5}$       e)  $\frac{21}{10}$       f)  $\frac{10}{21}$       g)  $\frac{16}{21}$   
h)  $\frac{80}{9}$       i)  $\frac{36}{35}$       j)  $\frac{35}{12}$

**4.7.1** a)  $\frac{15}{40}$       b)  $\frac{153}{32}$       c)  $\frac{46}{32}$       d)  $\frac{21}{648}$       e)  $\frac{203}{328}$

**4.8.1** a)  $\frac{4}{11}$       b)  $\frac{35}{8}$       c)  $\frac{1}{9}$       d) 4

**4.8.2** a)  $\frac{7}{4}$       b)  $\frac{3}{7}$       c)  $\frac{2}{3}$       d)  $\frac{8}{7}$       e)  $\frac{1}{2}$       f)  $\frac{7}{2}$

- 4.8.3** a) 49      b) 54      c) 70      d) 16      e) 30      f) 12      g) 25  
h) 14      i) 7      j) 63
- 4.9.3** a)  $\frac{14}{15}$       b)  $\frac{24}{45}$       c)  $\frac{30}{21}$       d)  $\frac{7}{20}$       e)  $\frac{66}{15}$
- 4.9.4** a)  $\frac{4}{9}$       b)  $\frac{15}{4}$       c)  $\frac{14}{5}$

**Gruble 3** a) 2      b) 4      c) 5      d) 2      e) 4      f) 5      g) 3, 4  
h) 2, 5      i) 3, 5      j) 4, 5

**Gruble 4** a) 2      b) 4      c) 5      d) 4, 3      e) 5, 2      f) 5, 3  
g) 5, 4


**Gruble 5** a)  $\frac{403}{6732}$       b)  $\frac{269}{3150}$

## Chapter 5

### 5.1.1

- a) 9 har retning mot højre og længde 9.  
b) 4 har retning mot højre og længde 4  
c) -3 har retning mot venstre og længde 3  
d) 12 har retning mot højre og længde 12  
e) -11 har retning mot venstre og længde 11  
f) -25 har retning mot venstre og længde 25

### 5.1.2



**5.1.3** a) 9, 4 og 12.      b) -3, -11 og -25.

**5.2.1** a) 1      b) 8      c) 6      d) 0

e) 3      f) 1      g) 10      h) 5

**5.2.2** a) -16      b) -8      c) -23      d) 0

e) -23      f) -17      g) -3      h) 0

**5.2.3** a) 15      b) 17      c) 12      d) 14

e) -12      f) -19      g) -12      h) -13

**5.2.4** a) 22      b) 22      c) -17      d) 14      e) 15      f) 13      g) -13  
h) -12

**5.2.5** a) -12      b) -50      c) -63      d) -24      e)  $(-7) \cdot 8$       f)  $(-3) \cdot 9$   
g)  $(-1) \cdot 12$       h)  $(-10) \cdot 4$       i) -21      j) -25      k) -12      l) -72

**5.2.6** a) -4      b) -6      c) -5      d) -4      e) -8      f) -9      g) -5  
h) -5      i) 8      j) 9      k) 5

## Chapter ??

**7.1.1** a) 14      b) 20      c) 24

**7.1.2** a) 2      b) 16      c) 27

**7.1.3** a) 2 og 8      b) 3 og 4      c) 3 og 6

**7.1.4** a) 81      b) 1) Et åpenbart eksempel er kvadratet fra oppgave a), som har bredde og høyde 9, og areal 81. 2) Bredde 15 og høyde 3, areal 45. 3) Bredde 12 og høyde 6, areal 72. *Note:* Multiple possible answers.

**7.1.5** a) 3      b) 10      c) 6      d) 6      e) 10      f) 6      g) 4  
h) 3      i) 28

**7.1.6** a) 90. *Note:* Grunnflaten kan også være 72 eller 80, avhengig av hvilken side man velger ut som grunnflate.

**7.2.1** See the solutions manual.

**7.2.2** See the solutions manual.

**Gruble 6** See the solutions manual.

**Gruble 7** See the solutions manual.

**Gruble 20** See the solutions manual.

## Chapter??

**8.1.1** a)  $3a$       b)  $4a$       c)  $7a$       d)  $-2b$       e)  $-5b$       f)  $-3k$

**8.1.2** a)  $a + b$       b)  $a + 2b$       c)  $9b - 3a$

**8.1.3** a)  $2b - 5a + c$       b)  $3b - 9a$       c)  $11b - 3a$

**8.1.4** a)  $7a + 14$       b)  $9b + 27$       c)  $8b - 24c$       d)  $-6a - 10b$       e)  $(9a + 2)$   
f)  $(3b + 8)a$       g)  $3ac - ab$       h)  $2a + 6b + 8c$       i)  $27b - 9c + 63a$   
j)  $2c - 6b - 14a$

**8.1.5** Bruk [Rule 3.2](#) til å faktorisere uttrykket.

a)  $2(a + b)$       b)  $b(4a + 5)$       c)  $c(9b - 1)$       d)  $2a(2c - 1)$

**8.1.6**

**8.1.7** a)  $\frac{1}{4}$       b)  $\frac{1}{2}$  når  $x = 4$  og 2 når  $x = -2$ .

**8.1.8** Uttrykkene gitt av a) og c) stemmer.

**8.2.1** a)  $3^4$       b)  $5^2$       c)  $7^6$       d)  $a^3$       e)  $b^2$       f)  $(-c)^4$  *Note:*  
 $(-c)^4 = c^4$

**8.2.2** a) 64      b) 32      c) 64      d) -8      e) -243      f) 256

**8.2.3** a)  $2^{16}$       b)  $3^{11}$       c)  $9^6$       d)  $6^5$       e)  $5^{-4}$       f)  $10^{11}$       g)  $a^{16}$   
h)  $k^7$       i)  $x^3$       j)  $x$       k) 1      l)  $a^5 \cdot b^{-3}$

**8.2.4** a) 5      b) 10      c) 12      d) 3      e) 9      f) 10

**Gruble 8**  $2^{-\frac{5}{4}}$

## Chapter9

**9.2.1** a)  $x = 10$       b)  $x = 5$       c)  $x = 9$       d)  $x = 2$       e)  $x = 3$   
f)  $x = 4$       g)  $x = 10$       h)  $x = 8$       i)  $x = 10$

**9.2.2** a)  $x = 37$       b)  $x = 29$       c)  $x = 23$       d)  $x = 15$       e)  $x = 8$   
f)  $x = 11$       g)  $x = 23$       h)  $x = 19$       i)  $x = 10$       j)  $x = 28$

**9.2.3**

a)  $x = 4$       b)  $x = 5$       c)  $x = 9$       d)  $x = 15$



- 9.2.4** a)  $x = 8$       b)  $x = 72$       c)  $x = 49$       d)  $x = 150$   
**9.2.5** a)  $x = 7$       b)  $x = 10$       c)  $x = 6$       d)  $x = 9$       e)  $x = 9$   
 f)  $x = 8$       g)  $x = 5$       h)  $x = 2$

**9.2.6** See the solutions manual.

**9.2.7**  $x = 18$

**9.5.1**  $x = 1, y = -2$

**Gruble 10** See the solutions manual.

## Chapter10

**10.1.1** a)  $f(x) = 2x + 4$ .      b) 204      c)  $x = 24$ .

**10.1.2** a)  $a(x) = 5x^2$ .      b)  $x = 2000?$       c)  $x = 9?$

**10.1.3** a)  $b(x) = x^2 + 2x$       b) 440?      c)  $x = 8?$

**10.1.4** 22 trekanter og 10 firkanter.

**10.1.5** La  $x$  være et positivt heltall. a)  $p(x) = 2n$       b)  $o(x) = 2n - 1$

**10.2.1**

- a) Stigningstall 5 og konstantledd 10.      b) Stigningstall 3 og konstantledd  $-12$ .  
 c) Stigningstall  $-\frac{1}{7}$  og konstantledd  $-9$ .      d) Stigningstall  $\frac{3}{2}$  og konstantledd  $-\frac{1}{4}$ .

**10.2.2** See the solutions manual.

**10.3.1** a) (I) og (II) gir hver for seg en ligning som bekrirer en rett linje. Disse linjene kan også representeres ved  $f$  og  $g$  slik som de er definert.      b)  $x = 7$  og  $y = 2$ .

**10.3.2**  $f(x) = -8x + 16$

**10.3.3**  $f(x) = 4x + 3$

**10.3.4**  $f(x) = x - 3$  og  $g(x) = \frac{1}{4}x + 1$

**Gruble 11** See the solutions manual.

## Chapter??

**11.1.1**  $80^\circ$

**11.1.2**  $AC$  og  $DE$ ,  $BC$  og  $EF$ ,  $AB$  og  $DF$ .

**11.1.3**  $EF = 3$  og  $DF = 2$ .

**11.1.4**  $AC = 15$  og  $DF = 6$ .

**11.1.6** a) 9      b)  $a^2$

**11.1.7** a)  $100\pi$       b)  $400\pi$

**11.1.8** a) 27      b)  $a^3$

**Gruble 15** Se side ?? i [TM1](#).

**Gruble ??** See the solutions manual.

**Gruble ??** See the solutions manual.

**Gruble 17** See the solutions manual.

**Gruble 18** See the solutions manual.

**Gruble 19** See the solutions manual.

**Gruble 14** See the solutions manual.

# Literature

Kiselev, A. (2006). *Kiselev's Geometry: Book 1. Planimetry* (A. Givental, Overs.). Sumizdat. (Originally published 1892).

Lindstrøm, T. (2006). *Kalkulus* (2nd ed.). Oslo, Universitetsforlaget AS.

Moise, E. E. (1974). *Elementary geometry from an advanced standpoint*. Reading, Addison-Wesley Publishing Company.

Spivak, M. (1994). *Calculus* (3rd ed.). Cambridge, Cambridge University Press

*Note: The text, at least a very similar one, about Pythagoras's theorem on page 224 was first printed in Skage Hansen's book Tempelgeometri (2020).*

# Indeks

$\pi$ , 236

absolute value, 83

algebra, 159

altitude, 133

angle, 128

    right, 129

    side, 128

    vertex, 128

    vertical, 131

arc, 127

area, 139

areal

    of a circle, 221

    of a rectangle, 215

    of a trapezoid, 218

base, 133, 164

base surface, 150

cancellation, 64

circle, 127

    center, 127

circumference, 220

common denominator, 57

constant, 159

coordinate system, 16

degree, 129

denominator, 48

diameter, 127

difference, 23

digits, 12

dividend, 28

divisor, 28

edge, 132

equal sign, 10

equation, 180

exponent, 164

factor, 25

factorization, 42

formula, 215

fraction, 48

    expansion of, 52

    inverted, 69

    simplifying of, 52

function, 197

    graph, 200

    linear, 200

height, 133

hypotenuse, 134

intercept, 202

interval, 200

legs, 134

length, 82, 140

line, 125

magnitude, 83

number

    irrational, 172

    length of, 83

- natural, 11
- negative, 82
- positive, 82
- prime, 42
- rational, 71
- numbers
  - even, 15
  - mixed, 71
  - odd, 15
- numerator, 48
- of a parallelogram, 217
- of a triangle, 216
- parallel, 128
- perimeter, 138
  - of a circle, 220
- point, 16, 125
- point-slope equation, 213
- polygon, 132
  - vertices of, 132
- positive integers, 11
- power equation, 191
- prime factorization, 43
- product, 25
- quadrilateral, 132

- quotient, 28
- radical sign, 171
- radius, 127
- ratio, 28
- remainder, 109
- sector, 127
- segment, 125
  - perpendicular, 129
- side
  - corresponding, 226
  - of polygon, 132
- sign, 83
- slope, 202
- sum, 21
- surface, 139
- tal, 10
- term, 21, 23
- triangle, 132
  - congruent, 225
  - similar, 225
- value, 12
- variable, 159
- width, 140

## **The author**

Sindre Sogge Heggen earned a master degree of Applied mathematics attending the University of Oslo.