



# The Fractional Knapsack

# The Knapsack Problem...



Capacity of knapsack:  $K = 4$


# The Knapsack Problem...



Capacity of knapsack:  $K = 4$

**Fractional** Knapsack Problem:


# The Knapsack Problem...



Capacity of knapsack:  $K = 4$

**Fractional** Knapsack Problem:  
Can take a **fraction** of an item.

# The Knapsack Problem...




Capacity of knapsack:  $K = 4$

**Fractional** Knapsack Problem:  
Can take a **fraction** of an item.

Solution:

2 pd A \$100	2 pd C \$80
--------------------	-------------------

# The Knapsack Problem...



Capacity of knapsack:  $K = 4$


**Fractional** Knapsack Problem:  
Can take a **fraction** of an item.

Solution:

2 pd A \$100	2 pd C \$80
--------------------	-------------------

**0-1** Knapsack Problem:  
Can only **take or leave** item. You  
can't take a fraction.

# The Knapsack Problem...



Capacity of knapsack:  $K = 4$

**Fractional** Knapsack Problem:  
Can take a **fraction** of an item.

Solution:

2 pd A \$100	2 pd C \$80
--------------------	-------------------

**0-1** Knapsack Problem:  
Can only **take or leave** item. You  
can't take a fraction.

Solution:

3 pd C \$120	
--------------------	--

# The Fractional Knapsack Problem: Formal Definition

Given  $K$  and a set of  $n$  items:

weight	$w_1$	$w_2$	$\dots$	$w_n$
value	$v_1$	$v_2$	$\dots$	$v_n$

Find:  $0 \leq x_i \leq 1, i = 1, 2, \dots, n$  such that

$$\sum_{i=1}^n x_i w_i \leq K$$

and the following is maximized:

$$\sum_{i=1}^n x_i v_i$$



The Knapsack problem.


Algorithm for the fractional knapsack problem?

# Greedy Solution for Fractional Knapsack

Sort items by **decreasing** value-per-pound


# Greedy Solution for Fractional Knapsack

Sort items by **decreasing** value-per-pound



# Greedy Solution for Fractional Knapsack

Sort items by **decreasing** value-per-pound



If knapsack holds  $K = 5$  pd, solution is:

1	pd	A
3	pd	B
1	pd	C

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

Let the sorted item sequence be  $1, 2, \dots, i, \dots, n$ ,

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

Let the sorted item sequence be  $1, 2, \dots, i, \dots, n$ , and the corresponding value-per-pound and weight be  $\rho_i$  and  $w_i$  respectively.



# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

Let the sorted item sequence be  $1, 2, \dots, i, \dots, n$ , and the corresponding value-per-pound and weight be  $\rho_i$  and  $w_i$  respectively.

Let  $k$  be the current weight limit (Initially,  $k = K$ ).

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

Let the sorted item sequence be  $1, 2, \dots, i, \dots, n$ , and the corresponding value-per-pound and weight be  $\rho_i$  and  $w_i$  respectively.

Let  $k$  be the current weight limit (Initially,  $k = K$ ).

In each iteration, we choose item  $i$  from the head of the unselected list.

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

Let the sorted item sequence be  $1, 2, \dots, i, \dots, n$ , and the corresponding value-per-pound and weight be  $\rho_i$  and  $w_i$  respectively.

Let  $k$  be the current weight limit (Initially,  $k = K$ ).

In each iteration, we choose item  $i$  from the head of the unselected list.

- If  $k \geq w_i$ ,

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

Let the sorted item sequence be  $1, 2, \dots, i, \dots, n$ , and the corresponding value-per-pound and weight be  $\rho_i$  and  $w_i$  respectively.

Let  $k$  be the current weight limit (Initially,  $k = K$ ).

In each iteration, we choose item  $i$  from the head of the unselected list.

- If  $k \geq w_i$ , set  $x_i = 1$  (we take item  $i$ ), and reduce  $k = k - w_i$ , then consider the next unselected item.

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

Let the sorted item sequence be  $1, 2, \dots, i, \dots, n$ , and the corresponding value-per-pound and weight be  $\rho_i$  and  $w_i$  respectively.

Let  $k$  be the current weight limit (Initially,  $k = K$ ).

In each iteration, we choose item  $i$  from the head of the unselected list.

- If  $k \geq w_i$ , set  $x_i = 1$  (we take item  $i$ ), and reduce  $k = k - w_i$ , then consider the next unselected item.
- If  $k < w_i$ ,

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

Let the sorted item sequence be  $1, 2, \dots, i, \dots, n$ , and the corresponding value-per-pound and weight be  $\rho_i$  and  $w_i$  respectively.

Let  $k$  be the current weight limit (Initially,  $k = K$ ).

In each iteration, we choose item  $i$  from the head of the unselected list.

- If  $k \geq w_i$ , set  $x_i = 1$  (we take item  $i$ ), and reduce  $k = k - w_i$ , then consider the next unselected item.
- If  $k < w_i$ , set  $x_i = k/w_i$  ( we take a **fraction**  $k/w_i$  of item  $i$ ),

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

Let the sorted item sequence be  $1, 2, \dots, i, \dots, n$ , and the corresponding value-per-pound and weight be  $\rho_i$  and  $w_i$  respectively.

Let  $k$  be the current weight limit (Initially,  $k = K$ ).

In each iteration, we choose item  $i$  from the head of the unselected list.

- If  $k \geq w_i$ , set  $x_i = 1$  (we take item  $i$ ), and reduce  $k = k - w_i$ , then consider the next unselected item.
- If  $k < w_i$ , set  $x_i = k/w_i$  ( we take a **fraction**  $k/w_i$  of item  $i$ ), Then the algorithm terminates.

# Greedy Solution for Fractional Knapsack

Calculate the value-per-pound  $\rho_i = \frac{v_i}{w_i}$  for  $i = 1, 2, \dots, n$ .

Sort the items by decreasing  $\rho_i$ .

Let the sorted item sequence be  $1, 2, \dots, i, \dots, n$ , and the corresponding value-per-pound and weight be  $\rho_i$  and  $w_i$  respectively.

Let  $k$  be the current weight limit (Initially,  $k = K$ ).

In each iteration, we choose item  $i$  from the head of the unselected list.

- If  $k \geq w_i$ , set  $x_i = 1$  (we take item  $i$ ), and reduce  $k = k - w_i$ , then consider the next unselected item.
- If  $k < w_i$ , set  $x_i = k/w_i$  ( we take a **fraction**  $k/w_i$  of item  $i$ ), Then the algorithm terminates.

Running time:  $O(n \log n)$ .



# Greedy Solution for Fractional Knapsack

Observe that the algorithm may take a fraction of an item.

# Greedy Solution for Fractional Knapsack

Observe that the algorithm may take a fraction of an item.  
This can **only** be the **last** selected item.

# Greedy Solution for Fractional Knapsack

Observe that the algorithm may take a fraction of an item.  
This can **only** be the **last** selected item.

We claim that the total value for this set of items is the **optimal** value.

Introduction

The Knapsack problem.

Algorithm for the fractional knapsack problem

Correctness

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

# Correctness

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

$x_i$  indicates fraction of item  $i$  taken



# Correctness

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

$x_i$  indicates fraction of item  $i$  taken (all  $x_i = 1$ , except possibly for  $i = k$ ).

# Correctness

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

$x_i$  indicates fraction of item  $i$  taken (all  $x_i = 1$ , except possibly for  $i = k$ ).

Consider any optimal solution  $O = \langle y_1, y_2, \dots, y_n \rangle$

# Correctness

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

$x_i$  indicates fraction of item  $i$  taken (all  $x_i = 1$ , except possibly for  $i = k$ ).

Consider any optimal solution  $O = \langle y_1, y_2, \dots, y_n \rangle$

$y_i$  indicates fraction of item  $i$  taken in  $O$

# Correctness

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

$x_i$  indicates fraction of item  $i$  taken (all  $x_i = 1$ , except possibly for  $i = k$ ).

Consider any optimal solution  $O = \langle y_1, y_2, \dots, y_n \rangle$

$y_i$  indicates fraction of item  $i$  taken in  $O$  (for all  $i$ ,  $0 \leq y_i \leq 1$ ).

# Correctness

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

$x_i$  indicates fraction of item  $i$  taken (all  $x_i = 1$ , except possibly for  $i = k$ ).

Consider any optimal solution  $O = \langle y_1, y_2, \dots, y_n \rangle$

$y_i$  indicates fraction of item  $i$  taken in  $O$  (for all  $i$ ,  $0 \leq y_i \leq 1$ ).

Knapsack must be full in both  $G$  and  $O$ :

$$\sum_{i=1}^n x_i w_i = \sum_{i=1}^n y_i w_i = K.$$

# Correctness

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

$x_i$  indicates fraction of item  $i$  taken (all  $x_i = 1$ , except possibly for  $i = k$ ).

Consider any optimal solution  $O = \langle y_1, y_2, \dots, y_n \rangle$

$y_i$  indicates fraction of item  $i$  taken in  $O$  (for all  $i$ ,  $0 \leq y_i \leq 1$ ).

Knapsack must be full in both  $G$  and  $O$ :

$$\sum_{i=1}^n x_i w_i = \sum_{i=1}^n y_i w_i = K.$$

Consider the first item  $i$  where the two selections differ.

# Correctness

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

$x_i$  indicates fraction of item  $i$  taken (all  $x_i = 1$ , except possibly for  $i = k$ ).

Consider any optimal solution  $O = \langle y_1, y_2, \dots, y_n \rangle$

$y_i$  indicates fraction of item  $i$  taken in  $O$  (for all  $i$ ,  $0 \leq y_i \leq 1$ ).

Knapsack must be full in both  $G$  and  $O$ :

$$\sum_{i=1}^n x_i w_i = \sum_{i=1}^n y_i w_i = K.$$

Consider the first item  $i$  where the two selections differ.

By definition, solution  $G$  takes a greater amount of item  $i$  than solution  $O$

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

$x_i$  indicates fraction of item  $i$  taken (all  $x_i = 1$ , except possibly for  $i = k$ ).

Consider any optimal solution  $O = \langle y_1, y_2, \dots, y_n \rangle$

$y_i$  indicates fraction of item  $i$  taken in  $O$  (for all  $i$ ,  $0 \leq y_i \leq 1$ ).

Knapsack must be full in both  $G$  and  $O$ :

$$\sum_{i=1}^n x_i w_i = \sum_{i=1}^n y_i w_i = K.$$

Consider the first item  $i$  where the two selections differ.

By definition, solution  $G$  takes a greater amount of item  $i$  than solution  $O$  (because the greedy solution always takes as much as it can).



# Correctness

Given a set of  $n$  items  $\{1, 2, \dots, n\}$ .

Assume items sorted by per-pound values:  $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$ .

Let the greedy solution be  $G = \langle x_1, x_2, \dots, x_k \rangle$

$x_i$  indicates fraction of item  $i$  taken (all  $x_i = 1$ , except possibly for  $i = k$ ).

Consider any optimal solution  $O = \langle y_1, y_2, \dots, y_n \rangle$

$y_i$  indicates fraction of item  $i$  taken in  $O$  (for all  $i$ ,  $0 \leq y_i \leq 1$ ).

Knapsack must be full in both  $G$  and  $O$ :

$$\sum_{i=1}^n x_i w_i = \sum_{i=1}^n y_i w_i = K.$$

Consider the first item  $i$  where the two selections differ.

By definition, solution  $G$  takes a greater amount of item  $i$  than solution  $O$  (because the greedy solution always takes as much as it can). Let  $x = x_i - y_i$ .

## Correctness...

Consider the following new solution  $O'$  constructed from  $O$ :

## Correctness...

Consider the following new solution  $O'$  constructed from  $O$ :

For  $j < i$ , keep  $y'_j = y_j$ .

Set  $y'_i = x_i$ .

In  $O$ , remove items of total weight  $xw_i$  from items  $i + 1$  to  $n$ , resetting the  $y'_j$  appropriately.

## Correctness...

Consider the following new solution  $O'$  constructed from  $O$ :

For  $j < i$ , keep  $y'_j = y_j$ .

Set  $y'_i = x_i$ .

In  $O$ , remove items of total weight  $xw_i$  from items  $i + 1$  to  $n$ , resetting the  $y'_j$  appropriately.

This is always doable because 
$$\sum_{j=1}^n x_j w_j = \sum_{j=1}^n y_j w_j = K$$

The total value of solution  $O'$  is greater than or equal to the total value of solution  $O$  (why?)

## Correctness...

Consider the following new solution  $O'$  constructed from  $O$ :

For  $j < i$ , keep  $y'_j = y_j$ .

Set  $y'_i = x_i$ .

In  $O$ , remove items of total weight  $xw_i$  from items  $i + 1$  to  $n$ , resetting the  $y'_j$  appropriately.

This is always doable because 
$$\sum_{j=1}^n x_j w_j = \sum_{j=1}^n y_j w_j = K$$

The total value of solution  $O'$  is greater than or equal to the total value of solution  $O$  (why?)

Since  $O$  is largest possible solution and value of  $O'$  cannot be smaller than that of  $O$ ,  $O$  and  $O'$  must be equal.

## Correctness...

Consider the following new solution  $O'$  constructed from  $O$ :

For  $j < i$ , keep  $y'_j = y_j$ .

Set  $y'_i = x_i$ .

In  $O$ , remove items of total weight  $xw_i$  from items  $i + 1$  to  $n$ , resetting the  $y'_j$  appropriately.

This is always doable because 
$$\sum_{j=1}^n x_j w_j = \sum_{j=1}^n y_j w_j = K$$

The total value of solution  $O'$  is greater than or equal to the total value of solution  $O$  (why?)

Since  $O$  is largest possible solution and value of  $O'$  cannot be smaller than that of  $O$ ,  $O$  and  $O'$  must be equal.

Thus solution  $O'$  is also optimal.

## Correctness...

Consider the following new solution  $O'$  constructed from  $O$ :

For  $j < i$ , keep  $y'_j = y_j$ .

Set  $y'_i = x_i$ .

In  $O$ , remove items of total weight  $xw_i$  from items  $i + 1$  to  $n$ , resetting the  $y'_j$  appropriately.

This is always doable because 
$$\sum_{j=1}^n x_j w_j = \sum_{j=1}^n y_j w_j = K$$

The total value of solution  $O'$  is greater than or equal to the total value of solution  $O$  (why?)

Since  $O$  is largest possible solution and value of  $O'$  cannot be smaller than that of  $O$ ,  $O$  and  $O'$  must be equal.

Thus solution  $O'$  is also optimal.

By repeating this process, we will eventually convert  $O$  into  $G$ , without changing the total value of the selection.

# Correctness...

Consider the following new solution  $O'$  constructed from  $O$ :

For  $j < i$ , keep  $y'_j = y_j$ .

Set  $y'_i = x_i$ .

In  $O$ , remove items of total weight  $xw_i$  from items  $i + 1$  to  $n$ , resetting the  $y'_j$  appropriately.

This is always doable because 
$$\sum_{j=1}^n x_j w_j = \sum_{j=1}^n y_j w_j = K$$

The total value of solution  $O'$  is greater than or equal to the total value of solution  $O$  (why?)

Since  $O$  is largest possible solution and value of  $O'$  cannot be smaller than that of  $O$ ,  $O$  and  $O'$  must be equal.

Thus solution  $O'$  is also optimal.

By repeating this process, we will eventually convert  $O$  into  $G$ , without changing the total value of the selection.

**Therefore  $G$  is also optimal!**